

Abstract

The NorESM1-M simulation results for CMIP5 (<http://cmip-pcmdi.llnl.gov/cmip5/index.html>) are described and discussed. Together with the accompanying paper by Bentsen et al. (2012), this paper documents that NorESM1-M is a valuable global climate model for research and for providing complementary results to the evaluation of possible man made climate change. NorESM is based on the model CCSM4 operated at NCAR on behalf of many contributors in USA. The ocean model is replaced by a developed version of MICOM and the atmospheric model is extended with on-line calculations of aerosols, their direct effect, and their indirect effect on warm clouds. Model validation is presented in a companion paper (Bentsen et al., 2012). NorESM1-M is estimated to have equilibrium climate sensitivity slightly smaller than 2.9 K, a transient climate response just below 1.4 K, and is less sensitive than most other models. Cloud feedbacks damp the response, and a strong AMOC reduces the heat fraction available for increasing near surface temperatures, for evaporation, and for melting ice. The future projections based on RCP scenarios yield global surface air temperature increase almost one standard deviation lower than a 15-model average. Summer sea-ice is projected to decrease considerably by 2100, and completely for RCP8.5. The AMOC is projected to reduce by 12 %, 15–17 %, and 32 % for the RCP2.6, 4.5, 6.0 and 8.5 respectively. Precipitation is projected to increase in the tropics, decrease in the subtropics and in southern parts of the northern extra-tropics during summer, and otherwise increase in most of the extra-tropics. Changes in the atmospheric water cycle indicate that precipitation events over continents will become more intense and dry spells more frequent. Extra-tropical storminess in the Northern Hemisphere is projected to shift northwards. There are indications of more frequent spring and summer blocking in the Euro-Atlantic sectors and that ENSO events weaken but appear more frequent. These indications are uncertain because of biases in the model's representation of present-day conditions. There are indications that positive phase PNA and negative phase NAO become less frequent under the RCP8.5 scenario, but also this result is considered uncertain.

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Single-forcing experiments indicate that aerosols and greenhouse gases produce similar geographical patterns of response for near surface temperature and precipitation. These patterns tend to have opposite sign, with important exceptions for precipitation at low latitudes. The asymmetric aerosol effects between the two hemispheres leads to a southward displacement of ITCZ. Both forcing agents thus tend to reduce northern hemispheric subtropical precipitation.

1 Introduction

Simulations of the Earth's climate are presented using a version of the Norwegian Earth System Model (NorESM1-M) with online calculations of aerosols and their direct effect and the first and second indirect effects of warm clouds. Explicit description of the interactive carbon cycle is not included in the present version, but is included in a version (NorESM1-ME) that is described by Tjiputra et al. (2012). In the companion paper by Bentsen et al. (2012) the NorESM1-M model system is described in technical detail and validated through evaluation of its conservative properties and by comparing simulation results with observationally based data for the historical period since 1850.

The present paper focuses on the simulated response of NorESM1-M to a selection of experiments, including projections of the future global climate based on scenarios defined in the fifth phase of the Coupled Model Intercomparison Project (CMIP5) (Taylor et al., 2012). A range of climate models and climate model versions participate in CMIP5, thereby providing input to the fifth Assessment Report (AR5) of the Intergovernmental Panel on Climate Change (IPCC) for inclusion in the fifth Assessment scheduled for publication in 2013. All data produced by the participating models, including NorESM1-M, can be downloaded from the CMIP5 multi-model data archive (<http://cmip-pcmdi.llnl.gov/cmip5/index.html>).

As elaborated in more detail by Bentsen et al. (2012), NorESM1-M is to a large extent based on the fourth version of the Community Climate System Model (CCSM4) developed in the Community Earth System Model project centred at the US National Center

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for Atmospheric Research (NCAR) but in collaboration with many partners (Gent et al., 2011). For NorESM, the ocean model in CCSM4 is replaced by a further developed version of the Miami Isopycnic Community Ocean Model (MICOM) and adapted for multi-century simulations in coupled mode (Assmann et al., 2010; Otterå et al., 2010).

5 An earlier version of this ocean model was also used for this purpose in the Bergen Climate Model (BCM), which was used to provide data for CMIP3 and the AR4 of the IPCC (Furevik et al., 2003; Otterå et al., 2009). This part of the development work for NorESM1-M is predominantly based at the Bjerknes Centre in Bergen, Norway. All the extensions of the ocean model since the original MICOM are summarized by Bentsen
10 et al. (2012). Important extensions since the BCM version include improved parameterization of diapycnal mixing, thickness and isopycnal eddy diffusion, and the mixed layer depth.

The atmospheric model in NorESM1 (both M and ME) is based on the version of the original CAM4 that was publicly released in April 2010 (Neale et al., 2010, 2012).
15 Over the later 15 yr research and modelling groups at the University of Oslo and (later) the Norwegian Meteorological Institute in Oslo have used a range of model versions from NCAR to develop representations of aerosols and their interactions with radiation and warm cloud microphysics in order to study the direct and indirect aerosol effects on climate (Iversen and Seland, 2002, 2003; Kirkevåg and Iversen, 2002; Kristjánsson,
20 2002; Storelvmo et al., 2006; Seland et al., 2008; Hoose et al., 2009; Struthers, et al., 2011).

For studies of climate response of the aerosol processes, earlier versions of the atmospheric model were run coupled to a slab ocean (Kristjánsson et al., 2005; Kirkevåg et al., 2008a,b). In the fully coupled NorESM1 it was natural to base the aerosol work on
25 the atmospheric model CAM4. This version is thus named CAM4-Oslo, and its properties related to the modelled aerosols are thoroughly discussed by Kirkevåg et al. (2012). We use the finite volume dynamical core for transport calculations (Rasch et al., 2006) with horizontal resolution 1.9° latitude times 2.5° longitude (in short: 2 degrees) and with 26 levels in the vertical with a hybrid sigma-pressure co-ordinate. The horizontal grid

**NorESM1-M –
climate response and
scenario projections**T. Iversen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mesh size is double of the standard version used in CCSM4, although Gent et al. (2011) also discuss a 2 degree version. The stratiform cloud parameterization is based on Rasch and Kristjánsson (1998), and the parameterization of deep convective clouds follows Zhang and McFarlane (1995) extended with the plume dilution and Convective Momentum Transport also used in CCSM4 (Richter and Rasch, 2007; Neale et al., 2008). Plume dilution influences the vertical distribution of aerosols (Kirkevåg et al., 2012) and water vapour (Gent et al., 2012), and improves the tropical deep convection in a favourable way for modelling the Madden-Julian Oscillation (MJO) (Subramanian et al., 2011). The favourable MJO properties are also diagnosed for NorESM1-M by Bentsen et al. (2012). NorESM1-M accounts for the radiative effects of deposited light-absorbing mineral dust and black carbon on snow (Flanner and Zender, 2006) and sea-ice.

A schematic of the CMIP5-experiments with NorESM1-M is shown by Bentsen et al. (2012) in their Fig. 1. Throughout this paper, we use “piControl” to identify the 500 yr control simulation with constant external forcing prescribed at 1850 conditions, which starts in year 700 after a spin-up with the same forcing. The spin-up is done to reduce trends in the piControl after tuning of parameters as discussed by Bentsen et al. (2012). Three ensemble members were branched off from the control at years 700, 730 and 760 for simulations “Historic1”, “Historic2” and “Historic3”. From 1850 to 2005, natural variations of solar radiation (Lean, 2000; Wang et al., 2005) and stratospheric sulphate aerosol concentrations from explosive volcanoes (Ammann et al., 2003), as well as anthropogenic GHG concentrations, aerosol emissions, and land-cover changes, were prescribed using the data from <http://cmip-pcmdi.llnl.gov/cmip5/forcing.html>. Historical forcing experiments are “GHG only”, “Aerosol only”, and “Natural forcing only”, where all but the single forcing identified by the name are kept constant as in piControl. From 2005 onwards, the representative concentration pathway (RCP) scenarios (van Vuuren et al., 2011) were the basis for climate projections until 2100: RCP2.6, RCP4.5, RCP6.0 and RCP8.5, where the numbers are the expected TOA forcing in W m^{-2} by 2100. An extension of RCP4.5 until 2300 was run with NorESM1-M.

The historical simulations have been extended to 2012 using RCP8.5 for the years 2006–2012.

Bentsen et al. (2012) present a thorough validation analysis of trends in piControl along with comparisons of the historical runs with data that are observationally based or from global re-analyses. In summary, the average radiative heat flux at the top of the atmosphere (TOA) in piControl is positive but smaller than 0.1 W m^{-2} . More than 99 % of this excess heat is transferred to the oceans, which experience a statistically significant temperature increase. There are also small negative trends in ocean salinity, in winter maxima sea-ice area in both hemispheres, and in the Atlantic meridional overturning circulation (AMOC). Other climatologically important parameters have insignificant global trends during the 500 yr of the piControl, including surface air temperature, cloudiness, precipitation and evaporation. The difference between global evapotranspiration and precipitation ($E - P$) averaged over a few decades, is not significantly different from zero in any of the experiments, including piControl.

In summary, by the end of the 20th century the surface air temperature is simulated too low by about 0.8–0.9 K globally and 1.0–1.1 K over land. The global precipitation is estimated to be up to about 0.15 mm d^{-1} too high, the evaporation from oceans is over-estimated with ca. 4 %, and the net flux between oceans and continents are ca 8 % over-estimated. The intensity of the water-cycle is therefore slightly overestimated while the atmospheric lifetime of water vapour is close to correct (compared to Trenberth et al., 2011). These properties can be linked to the fact that the model underestimates the global cloud fraction considerably by 15–25 %, while the tropospheric liquid water is over-estimated (Jiang et al., 2012). The double ITCZ is less pronounced in NorESM1-M than in CCSM4 with the same resolution.

The model simulates flow patterns that can be associated with observations, such as the Madden-Julian Oscillation (MJO), which was simulated with skill already in CCSM4 (Subramanian, 2012), ENSO, and the northern and southern annular modes. The AMOC strength is in the upper range found in models contributing to CMIP3 and above the range estimated from synthesized observational data (Medhaug and

GMDD

5, 2933–2998, 2012

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Furevik, 2011). Whilst the sea-ice extent is overestimated in both hemispheres in summer and in the southern winter, it is underestimated during northern winter. Kirkevåg et al. (2012) used NorESM's atmospheric model CAM4-Oslo to estimate the direct and indirect forcing of aerosol changes between the years 1850 and 2000 (2006) to be -0.08 and -0.91 W m^{-2} (1.2 W m^{-2}) respectively. The estimated indirect forcing in warm clouds is modest compared to many other models, without assuming artificial lower thresholds in the number of aerosols and cloud droplets (Hoose et al., 2009). However, the modelled aerosol loadings are at the high end in the free troposphere (Myhre et al., 2012).

The main purpose of this paper is to establish that results from the CMIP5 experiments with NorESM1-M are valuable for the climate system science and the evaluation of possible anthropogenic influence on the global climate. After discussing climate sensitivity, response, and gross feedbacks in Sect. 2, the present paper addresses aspects of the historical simulations and the RCP scenarios produced with NorESM1-M. Section 3 discusses model simulated time-developments of global variables from 1850 to 2005 ("Historic") and onwards for future RCP projections. In Sect. 4 the single forcing experiments for 1850–2005 are addressed, whilst further discussions of the RCP scenario projections are done in Sect. 5. After an analysis of various regional climate patterns are done in Sect. 6, conclusions are drawn in Sect. 7.

2 Equilibrium climate sensitivity and transient response

Global climate models are useful for diagnosing a range of characteristics for how the global climate may respond to a standard specified forcing. This facilitates the comparison of climate change properties across different climate models. This section discusses results of two such experiments under the CMIP5 protocol with NorESM1-M integrated over 150 and 140 yr respectively, and initiated in year 700 (i.e. from the start, Bentsen et al., 2012) of the piControl. These are referred to as "abrupt $4 \times \text{CO}_2$ " (quadrupling of atmospheric CO_2 concentrations at $t=0$) and "gradual $4 \times \text{CO}_2$ " (1 % increase

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



per year until quadrupling). Results are presented in Tables 1–4 as well as in Fig. 1. Since we have not applied any proper method for estimating changes in single climate elements (e.g. cloud cover) in response to temperature increase while other elements are kept unchanged, the feedback factors we present (e.g. for clouds) are termed gross feedback factors, since they can be influenced by simultaneous changes in other elements than the temperature (e.g. snow cover). See Gettelman et al. (2012) for estimates of proper feedback factors. Our analysis is comparable to Andrews et al. (2012).

The Equilibrium Climate Sensitivity (ECS) is the change in global mean near-surface air temperature when a new climate equilibrium is reached after an abrupt increase of the atmospheric CO₂ concentrations is introduced to a climate already in equilibrium. To calculate the ECS from first principles requires a full climate model run over several thousands years (Boer and Yu, 2003). ECS is therefore frequently approximated as the difference, ΔT_{eq} , between equilibrium near surface air temperatures obtained from two runs over a few decades, but with a model version where the deep ocean model is replaced by a thermodynamic slab. Bitz et al. (2012) used a slab ocean model for which the deep ocean heat fluxes were calibrated with data from runs with the full CCSM4. With 1 degree atmospheric resolution they estimated $\Delta T_{\text{eq}} = 3.20\text{K}$ after doubling of CO₂, while 3.13 K was estimated for the 2 degree version. This is close to the value 3.14 K for the previous CAM3-based version of CAM-Oslo coupled to a slab ocean (Kirkevåg et al., 2008a).

Estimates of ΔT_{eq} for NorESM1-M with a slab ocean are not available, but two other approximations of ECS are estimated for the full NorESM1-M. Both methods use simultaneous values of surface air temperature change ($\Delta T(t)$) and TOA radiation imbalance ($\Delta R(t)$) estimated at the time t after the abrupt quadrupling of atmospheric CO₂ concentrations.

Gregory et al. (2004) proposed to use a linear regression between $\Delta R(t)$ and $\Delta T(t)$, assuming negligible contributions from time-varying feedbacks. The slope of the regression line is the overall feedback parameter $\lambda = -d\Delta R/d\Delta T$ (in units of $\text{W m}^{-2} \text{K}^{-1}$), the intercept at $\Delta T = 0$ approximates the instantaneous forcing R_f , while the intercept

**NorESM1-M –
climate response and
scenario projections**

T. Iversen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ΔT_{reg} at $\Delta R = 0$ approximates the ECS. In reality, this estimate of R_f disregards rapid adjustments during the first year of the simulation, and it therefore underestimates the true instantaneous forcing of the quadrupled CO_2 (Andrews et al., 2012).

Murphy (1995) proposed to use the remaining TOA radiative imbalance $\Delta R(t)$ at the time t to approximate ECS. This approximation, termed the effective climate sensitivity and denoted $\Delta T_{\text{eff}}(t)$, is:

$$\Delta T_{\text{eff}}(t) = \frac{\Delta T(t)R_f}{R_f - \Delta R(t)} \quad (1)$$

Assuming the same linear relationship between $\Delta T(t)$ and $\Delta R(t)$, ΔT_{eff} should not depend on time. However, slow feedback processes, for example involving the deep ocean, may cause changes to occur over decades and centuries (Senior and Mitchell, 2000). Furthermore, chaotic fluctuations in the climate response may lead to high-frequency variations in $\Delta R(t)$. Figure 1a shows results for both $\Delta T(t)$ (black dots for years 1–150) and ΔT_{eff} (red dots for years 111–150), where we assume $R_f = 7.0 \text{ W m}^{-2}$ as estimated by Kay et al. (2012).

The two approximations to ECS are $\Delta T_{\text{reg}}(4 \times \text{CO}_2) = 5.74 \text{ K}$ from the regression, with feedback parameter $\lambda \cong 1.101 \text{ W m}^{-2} \text{ K}^{-1}$, and $\Delta T_{\text{eff}}(4 \times \text{CO}_2) = 5.71 \text{ K}$ using Eq. (1) with values averaged over the last 40 of the 150 yr of the abrupt $4 \times \text{CO}_2$ experiment (black cross in Fig. 1). The numbers in Table 1 are these divided by 2 since the effect of CO_2 doubling are more standard in the literature (e.g. Andrews et al., 2012). Notice that the forcing approximated by the regression (see Fig. 1a) is only 6.32 W m^{-2} due to the fast adjustments during the first year of the integration. Furthermore, slow deep-oceanic feedbacks may delay the response and thus render the linear regression inaccurate. For example, a regression for years 1–76 yields a smaller approximation of the ECS: 5.18 K, indicating that there may be slow feedback mechanisms at work. Andrews et al. (2012) indicate that short-wave radiative effects of clouds over oceans may cause non-linearity over the first decades.

NorESM1-M – climate response and scenario projections

T. Iversen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

reduced by 8–10 Sv over the first couple of decades and then remains almost constant in the abrupt experiment, the reduction is slower and almost linear with time in the gradual experiment. The efficiency at which the net downward radiative heat flux at the top of the model is calculated to penetrate downwards in the world ocean is illustrated in Fig. 1e and f, which show the heat flux at different ocean depths averaged over the entire globe.

By the time of CO₂ doubling, AMOC is reduced with about 3–5 Sv in the gradual experiment. The heat fluxes into the deep ocean shown in Fig. 1e and f reduces the fraction of the net heat flux at the top of the model that is available for further increase in surface temperatures, evaporation of water, and melting of ice. An efficient heat transport into deep oceans thus reduces the traditional measures of climate sensitivity. It can be seen from Fig. 1f that a slab ocean model with 200 m thickness of the mixed layer would require almost 50 yr spin-up to reach a quasi-equilibrium state for the 4 × CO₂ climate. The transfer of heat into the deep ocean is a much slower and spatially heterogeneous process.

Despite the stronger AMOC in the experiment with gradual CO₂-increase, the heat transport into the deep ocean may appear more efficient in the abrupt experiment. This is an artefact caused by the exponential increase in atmospheric CO₂ (1 % increase per year) starting from pre-industrial levels. These annual forcing increments add to the TOA imbalance, and the increments penetrate into the deep ocean with a characteristic time which is influenced by the strength of the AMOC. As the AMOC strength decreases gradually, the downward heat transport at high latitudes reduces. The deep ocean heating will therefore continue several decades even without further CO₂ increase after the doubling (when the TRC is estimated), but the efficiency will gradually decrease as AMOC steadily reduces before stabilizing at a smaller value due to the heating and freshening of the upper ocean layers at high latitudes. This slow reduction of the deep ocean heating efficiency is different from the abrupt experiment which establishes a new quasi-stable AMOC already after a few decades. We hypothesize that

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\Delta T_{\text{TRC,eff}}$ at the time of CO₂-doubling underestimates the true equilibrium temperature after CO₂-doubling, due to this multi-decadal non-linear contribution to the feedbacks.

Tables 3 and 4 summarize a few results after applying linear regression between changes of a selection of global variables characterizing the global climate, and the changed surface air temperature for the abrupt 4 × CO₂ experiment. For all the hydroclimatic variables in Table 3, there are positive gross feedback factors, i.e. their change increase with increased temperature change. The factor for global precipitation increase is equivalent to ca 2.7 %K⁻¹, which probably is on the high side (e.g. Trenberth, 2011). The factor is about 6 times larger over the oceans than over continents, but almost all the response over the oceans are due to recycling of oceanic evaporation. The slight surplus of 0.02 × 10³ km³ K⁻¹ for oceanic evaporation over precipitation equals the deficit over the continents. This number results from a small imbalance between terms that are several orders of magnitude larger, and the implied uncertainty is shown as an interval in Table 2. Nevertheless, based on the abrupt 4 × CO₂ experiment, the model predicts a more intense water cycle with a small but uncertain increase in the atmospheric lifetime of water vapour with increased temperatures.

Corresponding factors for change in yearly mean sea-ice volume and area in each of the hemispheres are given in Table 4. The sensitivity parameters are all negative, and the sensitivity is considerably higher in the Arctic than in the Antarctic. In the Arctic, melting of sea-ice is particularly associated with the surface albedo feedback effect, which also involves changes in the snow cover.

3 Time trends of interactive forcing agents

The only prescribed aerosol concentrations in the model are stratospheric sulphate from explosive volcanoes in the historical period. Other aerosol components are calculated from prescribed emission data, or, for sea-salt, from emissions calculated as a function of wind speed and ocean temperature. Kirkevåg et al. (2012) present and evaluate the aerosol module, including estimates of direct and indirect aerosol forcing.

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



It is emphasized that a correct simulation of forcing of anthropogenic aerosols depends on the amount and properties of the natural background aerosols and the associated cloud droplet properties (see e.g. Hoose et al., 2009). It should be noted that there were considerable anthropogenic aerosols already in 1850 due to biomass burning.

In a few places, emissions from forest fires and also from natural secondary organics from areas that were forested, but no longer are, were larger in 1850. The model calculates mass concentrations of sulphate, black carbon (BC), and particulate organic matter (POM), which includes the secondary organics (SOA) in addition to the major natural components sea-salt and mineral dust. The aerosols interact directly with solar radiation, and a prognostic equation for liquid water droplets in stratiform clouds uses CCN activation of the aerosols according to size and composition (Storelvmo et al., 2006).

Figure 2 shows the historical and scenario developments of the average global loadings of particulate sulphate, BC, and POM since 1850 as simulated by NorESM1-M. Both natural and anthropogenic aerosols are included, but the major part of the long-term trends since 1850 are due to anthropogenic activities involving fossil fuel combustion and to some extent biomass burning. POM has a relatively larger fraction of natural aerosols because of biogenic emissions from oceans and from land vegetation. All the RCP scenarios, and RCP2.6 for BC in particular, peak during the first decades of the 21st century before decaying to slightly higher levels than in 1850 towards the end of the century. The globally averaged aerosol optical depth and the absorption component separately show the natural part in the historical period and include the contribution of stratospheric sulphate from known explosive volcanoes since 1850. The scattering effect of the volcanic aerosols is considerable for 1–3 yr in each case, which also demonstrate that the sustained impacts of the anthropogenic aerosols are due to the continuous replenishment from human activity. The decaying loadings and optical depths in the 21st century therefore follow immediately from assumed changes in emissions. For most greenhouse gases this is not the case, due to their long residence time in the Earth system.

**NorESM1-M –
climate response and
scenario projections**

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The curves in Fig. 3 show the calculated TOA long-wave, short-wave and net radiative imbalances in the period 1850 to 2300. While a negative trend is simulated for the long-wave as well as the short-wave radiation with a close to zero net effect up until 1970s. Then the trends become positive and increase in particular for the future RCP scenarios. The net TOA imbalance was ca. 0.6 W m^{-2} by the first decade of the 21st century, but the year-to-year variation is substantial. The effect of this change is seen in the global mean near surface temperature and in the global precipitation rate. Bentsen et al. (2012) discuss the realism of this and other results for the historical period.

4 Historical single forcing simulations

As an element in attributing climate change and variability since 1850 to possible causes, a few selected single forcing simulations are made as a part of the CMIP5 protocol. We have only run single realizations for each of these forcing simulations, which is insufficient to estimate statistical significance with respect to attribution of climate variations. However, they contribute to the multi-model ensemble in CMIP5 for IPCC AR5. Here we discuss three such experiments. In “GHG only”, all but the prescribed greenhouse gas concentrations are kept constant at the 1850-level; in “Aerosol only” all but aerosol emissions are as in 1850; and in “Natural forcing only”, only the natural contributions to the forcing are varied after 1850.

Figure 4 shows results for surface air temperature and precipitation. For temperature it appears that the simulated warming since the 1970s hardly can be reproduced with natural forcing only. Furthermore, the greenhouse gases alone will probably lead to an exaggerated warming estimate, while aerosols significantly dampen the warming exerted by GHG. For global precipitation, the picture is much less clear, and the regional variations in the simulated precipitation changes are crucial. In some continental regions considerably reduced annual precipitation is estimated even if the global trend is positive.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The maps in Fig. 4 also show that regional responses to GHG forcing and aerosol forcing show many similar geographical patterns, but the patterns have opposite sign to each other. Given that the spatial forcing patterns of GHG and aerosols are very different, the similarity in the response pattern demonstrates that internal dynamics (Palmer, 1999; Branstator and Selten, 2009) and geographically determined feedbacks (Boer and Yu, 2003) determine the nature of the climate response, rather than the forcing pattern itself. Kirkevåg et al. (2008b) found similar results with a model coupled to a slab ocean.

There are, however, some important exceptions for precipitation, since the response to GHG tends to systematically have the same sign as the response to aerosols in some areas. Such mutual reinforcement may, however, also occur by chance due to chaotic internal variability, and care must be taken not to exaggerate conclusions based on single realizations of the experiment. In some cases random reinforcements can be checked by a non-linearity test. The two spatial response fields of each of the single forcing experiments are then added, and the sum is compared to the result of a single experiment that employs the sum of the two forcing components. In areas where these two fields differ considerably, chaotic internal variations may dominate over systematic mutual reinforcement. This will not provide a final proof, however, since random patterns also may behave similarly by chance in the two single-forcing experiments. A stronger proof requires multiple ensemble members.

In the two maps in Fig. 5 contributions from GHG only and aerosol only are added (left) and can be compared to the changes in the Historic1 run (right). Although there are other minor forcing agents in the Historic1 run such as land usage, volcanoes, and solar radiation changes, the response patterns which differ considerably between the two maps can be regarded as dominated by chaotic randomness. In the tropics and subtropics, however, there are patterns that are less likely to be entirely random. The impact of GHG on the tropical and subtropical precipitation patterns shown in Fig. 4 (right middle panel) relates to an intensified Hadley circulation with increased precipitation close to the equator and reduced in the adjacent subtropics in both hemispheres.

**NorESM1-M –
climate response and
scenario projections**

T. Iversen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The changes over oceans west of South America and Africa are not statistically significant. There are also signs of an extended dry zone towards the middle latitudes. The impacts of aerosols tend to balance these GHG-driven changes at low latitudes (Fig. 4 lower right panel), but there are important exceptions since the cooling effects of anthropogenic aerosol are larger in the northern than the Southern Hemisphere. This asymmetry leads to a southward displacement of the Hadley circulation and the associated ITCZ, a result that was reported by e.g. Rotstayn and Lohmann (2002), Kristjánsson et al. (2005), and Kirkevåg et al. (2008b) using slab ocean models. This combination of a reduced and displaced Hadley circulation may lead to reinforcement of the considerably more symmetric strengthening caused by the GHG in some areas.

Since aerosols predominantly influence solar radiation, the asymmetric response is particularly pronounced during the NH summer. This can be seen by comparing Fig. 5a and b. The reinforcement of drying the subtropics appears more pronounced in the Northern Hemisphere than in the Southern while the moistening of the tropics is more pronounced in the southern flank of the ITCZ. As discussed in the companion paper by Bentsen et al. (2012), we relate the effects of aerosols to an improved precipitation pattern in NorESM1-M with a reduced split of the ITCZ over low latitudes in the Pacific Ocean compared to CCSM4 (Gent et al., 2011).

5 Climate projections based on RCP-scenarios

The first ensemble member of the historical period is extended with 4 climate projections to year 2100 in accordance with the timing of the prescribed scenarios for concentrations and emissions in the RCPs. RCP4.5 is extended further to year 2300. Results discussed here are given in Tables 5 and 6 and Figs. 6 to 10.

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5.1 Surface temperatures and sea-ice

Figure 6 shows calculated surface air temperatures at reference height relative to the 1850–1899 mean for a global average and for averaged over the polar region north of 65 degrees north. By the end of the 21st century the global mean ranges from ca. +1.2 K for RCP2.6 to ca. +4.0 K for RCP8.5 and the polar region mean from ca. +3.5 K to 9.7 K, with a notable downward trend for RCP2.6. The global numbers in Table 5 are smaller as they represent changes for two 30-yr periods separated by 95 yr (1976–2005 to 2071–2100).

The year-to-year variability is also considerably larger for the NH polar region than globally. The signal to noise ratio is not quantified, but is probably smaller in the polar region due to the large natural variability in the region. Bentsen et al. (2012) showed, in their Fig. 25, that the signal to noise ratio was too small in the area north of 60° N relative to the observed historical data to detect a significant trend.

Figure 7 shows a comparison of NorESM1-M surface air temperature relative to the 1850–1899 average, with statistics from 15 CMIP5 models (one being NorESM1-M, Andrews et al., 2012). The selected results are for the RCP2.6 and RCP8.5 scenarios and are global means (left) and averages over land areas (right). NorESM-values are within one standard deviation away from the multi-model mean but persistently lower than the average. This is in accordance with the relatively small climate sensitivity found in Sect. 2. A full explanation will require a separate study of the properties of all the models. Candidate explanations should for example take into account that NorESM1-M includes negative forcing contributions from the direct as well as the indirect aerosol effect. Furthermore, NorESM's gross cloud radiative feedback is negative, and the model's strong AMOC may transfer heat into the deep oceans more efficiently than many other models. Figure 7 also shows that both the inter-model spread and the size of the warming are considerably larger over land than globally, a feature which is also seen when comparing the changes in SST to those of global temperatures in Table 5. The simulated SST changes account for 55–60 % of the changes in global surface air

GMDD

5, 2933–2998, 2012

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



temperatures. This is well known, e.g. from IPCC reports (e.g. Trenberth et al., 2007). It can be attributed as a manifestation of the cold-ocean warm-land (COWL) pattern (Wallace et al., 1996), caused by the low heat capacity of the continents compared to the oceans where heat is mixed in deep water masses.

5 The geographical distribution of the simulated annual temperature changes for 2071–2100 relative to 1976–2005 for RCP2.6 and RCP8.5 are given in Fig. 8a and b. The temperature increase is considerably larger in RCP8.5 than for RCP2.6. The patterns of COWL and the Arctic amplification are seen for both scenarios. The Arctic sea-ice extent is also projected to decrease by 2100 for all scenarios (see Fig. 9c). The
10 reduction is particularly large for the annual minimum sea-ice extent, where the Arctic September sea-ice has almost disappeared by ca. 2100 for the RCP8.5 scenario. For the other scenario projections, some sea-ice always remains. The response in the Southern Hemisphere is considerably smaller. The simulated response from 1976–2005 to 2071–2100 in the total global and annual mean sea-ice area is given for each
15 RCP scenario in the bottom row in Table 5. In relative numbers the reduction varies from ca 7 % (RCP2.6) to ca 30 % (RCP8.5).

5.2 Precipitation

The climatology of precipitation and dry spells has strong impacts on nature as well as human society. Changes in annual total amounts as well as the intensity of precipitation events are important. As discussed by Bentsen et al. (2012) NorESM1-M over-
20 estimates the global precipitation by the end of the 20th century by ca 0.14 mm d^{-1} ($2.81\text{--}2.67 \text{ mm d}^{-1}$) compared to the GPCP data (Adler et al., 2003), where the over-estimations are particularly pronounced in the tropics. However, Trenberth (2011) mentions that GPCP values may underestimate warm rain in the extratropics and refers to increased estimates of 5 % over the ocean. NorESM1-M also has a slightly too intense hydrological cycle, since oceanic evaporation is about 4 % larger than estimates from synthesized data (Trenberth et al., 2011). Opposed to many other global climate
25 models with too fast recycling of water vapour, NorESM1-M slightly overestimates the

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



atmospheric residence time of oceanic water vapour, and the atmospheric transfer of water vapour from ocean to land is overestimated by about 8 % compared to the estimate of Trenberth et al. (2011).

Figure 8 shows projections of relative change (%) in annual precipitation amounts by 2071–2100 relative to 1976–2005 for the extreme scenarios RCP2.6 and RCP8.5. In general the patterns are the same, but RCP8.5 has larger anomalies. Some increases occur over arid regions giving very large relative changes, e.g. over central parts of Northern Africa. Otherwise, there are strong increases over the tropical Pacific Ocean paired with strong decreases in the subtropics. There is also reduced precipitation in the Mediterranean region and in southern parts of North America. Otherwise there are mainly precipitation increases, including at polar latitudes in both hemispheres. Many of these features are more pronounced in the seasonal maps for RCP8.5. Drying in the northern hemispheric continents is more pronounced in NH summer, whilst in the NH winter the precipitation increase is more ubiquitous.

Figure 9a and 9b show the simulated time development of $E - P$ and E integrated over the global oceans. The oceanic $E - P$ is the net water vapour transported from ocean to land in the atmosphere, while the oceanic E is a measure of the intensity of the hydrological cycle. Both quantities increase in the simulations of the RCP scenarios. The budget changes in Table 5 indicate that all gross quantities but one increase with the size of the forcing by 2071–2100 compared to 1976–2005. The quantity that does not increase is evaporation from land, which thus appears to be insensitive to the size of the forcing. The global annual precipitation amounts increase by 2.3–5.2 %, the oceanic evaporation by 2.3–4.5 %, the net atmospheric transfer from ocean to land by 2–18 %, and the precipitation on land by 2.5–8.2 %, where the low numbers are for RCP2.6 and the high for RCP8.5.

When evaporation from land in practice is preserved while both the net flux of vapour from ocean to land and the precipitation over land increase, the space-time fraction of dry spells over land must increase and thus also the average intensity of precipitation. Such effects were deduced for a warmer climate by Trenberth et al. (2003) and further

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



elaborated by Trenberth (2011). The relevance of dry spells in diagnosing the intensity of the hydrological cycle was thoroughly discussed and analyzed by Giorgi et al. (2011).

Table 6 analyses the possible future situation in Europe for the RCP scenarios. The table clearly shows that the NorESM1-M simulations produce a striking difference between increased precipitation in Northern Europe and more dry conditions in Southern Europe towards 2100. According to the simulations in Historic1, such a development may already have occurred.

5.3 AMOC and ocean temperatures

We have already discussed the possible regulating role of the AMOC for the impacts of radiative forcing on near surface air temperature, SST, surface evaporation, and melting of sea-ice. This 3-dimensional current is regarded as a part of the global ocean conveyor belt. A common view is that the upward closing branch is a large scale balance between upwelling and diapycnal mixing (e.g. Munk and Wunsch, 1998). AMOC is driven by wind stress and by thermohaline forcing. The latter occurs when cold and saline water is produced at high latitudes and becomes negatively buoyant and sinks.

Under anthropogenic climate change, increased precipitation and sea-ice melting of the cryosphere may stabilize the vertical water column at high latitudes and lead to a reduced AMOC (e.g. Hofmann and Rahmstorf, 2009). We have already seen that NorESM1-M simulates increased precipitation in the Northern North Atlantic Ocean and reduced precipitation in the subtropics under RCP scenarios (Fig. 8). Observational studies indicate that surface water has become fresher in areas relevant for deep water formation already during the recent decades (Curry and Mauritzen, 2005). A slower AMOC may be associated with reduced poleward transport of heat in the upper ocean and cause colder climate regionally over the Northern North Atlantic Ocean, the Arctic, and in North-West Europe. Persistent wind stress in the storm-track regions combined with increased subtropical surface salinity may compensate if more saline water is transported northwards by the surface wind driven currents (Bethke et al., 2006).

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 10 (upper panel) shows the time series of maximum AMOC strength at 26.5° N in the NorESM1-M simulations runs piControl, Historic1, 2 and 3, and the 4 RCP scenarios. The piControl time series has a mean value of 30.8 Sv and a small but significant (p -value < 0.01) linear trend of -0.6 Sv over 500 yr (Bentsen et al., 2012). The historical experiments do not clearly agree on a substantial deviation from the long-term evolution from the piControl experiment, which shows considerable amplitudes due to unforced internal variability. However, they all show a decreasing AMOC strength after about 1980, and in the moving averaged time series two of the historical members end at an AMOC strength near the minimum value encountered during the whole piControl integration.

The reduction seen in the RCP experiments is considerably larger than the trend and variability of the piControl. When comparing the mean AMOC strength of the years 2091–2100 in the scenario experiments to the mean strength of the control, the reductions are 3.6 Sv for RCP2.6, 5.1 Sv for RCP4.5, 5.6 Sv for RCP6.0, and 9.9 Sv for RCP8.5. The relative reductions are thus approximately 12 %, 17 %, 18 %, and 32 %, respectively. In the SRES A1B scenario experiment of 16 models participating in CMIP3, Schneider et al. (2007) found a mean reduction of maximum AMOC strength at 30° N of about 4 Sv from year 2000 to 2100. This amounts to an average decline of about 25 %. The SRES A1B scenario is closest to the RCP6.0 scenario in terms of estimated radiative forcing towards year 2100 (Houghton et al., 2001; van Vuuren, 2011). The NorESM1-M simulation based on the RCP4.5 scenario, which was extended to year 2300 with constant aerosol emissions and greenhouse gas concentrations after year 2100, shows a rather stable AMOC strength of 25–26 Sv after year 2100.

The zonal mean temperature change for years 2071–2100 of NorESM RCP8.5 compared to corresponding years of the piControl experiment is shown in the two bottom panels in Fig. 10. In the upper 200 m the warming is in excess of 2 K most places except where sea-ice is present and constrains the temperature at the freezing temperature. The warming is in general reduced with depth, and below 3000 m the warming signal is weak, particularly in the global average shown in the right panel of Fig. 10. At high

**NorESM1-M –
climate response and
scenario projections**

T. Iversen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

latitudes, the warming penetrates deeper. A generally stronger warming signal is seen for the Atlantic Ocean in the left panel of Fig. 10 compared to the global ocean. Also, for the Atlantic Ocean, the upper ocean temperature increase is efficiently communicated to the ocean interior at high latitudes.

5 The overturning circulation of the Atlantic seems to carry a warming signal southward in the Atlantic at 2000–3000 m depth. Given that AMOC strength is reduced by a third by 2100, further penetration of heat to the deep ocean will be considerably reduced, although it will remain strong in this model. A larger fraction of the greenhouse gas heating in the RCP8.5 scenario will nevertheless remain in the atmosphere and contribute to enhance the globally averaged surface warming. However, a considerably reduced heating (possibly cooling) may occur regionally at high latitudes adjacent to where the negatively buoyant water normally is formed in the Atlantic Ocean. In order to study such consequences, the RCP8.5-based simulation should be prolonged. Some caution should also be taken with regard to these results since NorESM1-M probably overestimate the strength of AMOC in the first place.

6 Changes in regional flow patterns

20 The climate of the mid-latitudes is closely linked to large-scale cyclones that develop and propagate in the westerly jet-stream systems. In particular, extreme precipitation and flooding are linked to storminess and transport in the associated warm conveyor belt (Stohl et al., 2008; Gimeno et al., 2011). Extratropical cyclones are also important vehicles for the atmospheric meridional transport of heat, humidity and momentum between the low and high latitudes, as well as the maintenance of the jet-streams themselves (e.g. Bratseth, 2001, 2003). Absence of cyclones associated with persistent blocking events is likewise important for the occurrence of droughts. Furthermore, the changed occurrence of flow regimes or prevalent intrinsic weather modes can be claimed to be a regional manifestation of global climate change (Corti et al., 25 1999; Branstator and Selten, 2009). In this section we address these aspects of

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the NorESM1-M simulation results, emphasizing the Northern Hemisphere (NH) and ENSO.

6.1 NH storminess

The climatological storminess in the Northern Hemisphere (NH) extratropics simulated with NorESM1-M is diagnosed using the standard deviation of 2.5–6 days band-pass filtered 500 hPa geopotential height. The band-pass filter is the same as used by Blackmon (1976), which has been shown to retain baroclinic waves consistent with theoretical and modelling studies (e.g. Chang, 2002; and references therein). Although this field represents baroclinic wave activity we will refer to it here as a measure of storm track activity. Figure 11 shows the annual mean storm tracks for the NorESM1-M historical simulation compared to the ERA-40 reanalysis (Uppala et al., 2005). Many of the main characteristics of the NH storm tracks are well simulated, although the amplitude of the band-pass filtered variability overall are slightly too weak. Note that this amplitude bias is significantly reduced for the AMIP run of the NorESM1-M model with prescribed SSTs based on observations. Parts of the bias in the fully coupled NorESM1-M can thus be attributed to systematic errors in the simulated SST field. Another notable bias is found over the North-Atlantic Ocean where the storm track is too zonal and lacks the characteristic poleward tilt in the ERA-40 data. This bias is very similar to that reported for the CAM3 model (Hurrell et al., 2006), and the bias appears to be intrinsic to the atmospheric model component, since it is virtually unchanged in the AMIP run of the NorESM1-M.

Figure 12 shows the projected changes for 2071–2100 compared to 1976–2005 for the RCP8.5 scenario. The annual mean change is dominated by a slight general weakening of the band-pass filtered variability except over the Northern North-Atlantic Ocean where a poleward shift in the storminess is apparent. This poleward shift is prominent mainly during summer and autumn. Although the shift is statistically significant at the 5% level relative to the variational spread in the time series, Fig. 12 also includes the spread between the 3 historical NorESM1-M simulations for the period

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1976–2005. This sample is of course too small to adequately represent the model's internal variability as an ensemble, but the amplitude of the poleward shift as well as the more general weakening can, at least, be seen to exceed this ensemble spread. A poleward shift of mid-latitude storm tracks has also been diagnosed in many other climate model simulations as a response to anthropogenic greenhouse-gas forcing (e.g. Yin, 2005).

6.2 NH blocking

Whilst storminess is associated with frequent occurrence of precipitation and possibly flooding, the blocking phenomenon is closely connected with persistent anticyclones, which tend to suppress precipitation at mid-latitudes for periods of up to several weeks. Incidents of extensive droughts can be associated with blocking, and the ability of climate models to simulate and project the climatic occurrence of droughts at mid-latitudes will be influenced by their ability to simulate blocking. Many blockings will also include a cold cyclone with low static stability and heavy convective storms.

To diagnose atmospheric blocking we use the index originally proposed by Lejanäs and Økland (1983) and later modified by Tibaldi and Molteni (1990) (TM). The TM index uses a persistent reversal of the gradient of the 500 hPa geopotential height around a central latitude (50° N) as an indicator of blocked flow. The central latitude is allowed to vary by approximately 3.8 degrees (2 grid points in the NorESM) to include small latitudinal shifts in the block. The blocks were further required to last for at least 5 days and be present at 7.5 degrees consecutive longitudes.

Pelly and Hoskins (2003) have shown that a fixed central blocking latitude suitable for detecting blocks over the North Atlantic leads to spurious detection over the North Pacific. We therefore also calculate a “vTM” index where the central latitude varies with the longitude of the climatological storm track. The central latitude is defined as the maximum of the standard deviation of the 2.5–6 days band-pass filtered geopotential height anomalies at 500 hPa. To account for the seasonal cycle of the storminess, the

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



central latitude for a given month is calculated as the climatological 3-month average centred on that month.

Figure 13 shows the seasonal blocking frequency for the NorESM1-M Historic1 simulation for 1976–2005 compared to the ERA-40 reanalysis for 1979–2002. The variable latitude “vTM” index is shown. For all seasons the model largely fails to adequately simulate blocking over the North-Atlantic Ocean and Western Europe in NH winter and spring. This is consistent with the too zonal propagation of storms in this sector (Fig. 11). This common deficiency amongst climate models (e.g. D’Andrea et al., 1998) may be partly attributed to the coarse resolution, as the investigation of Jung et al. (2012) suggests that around 40 km resolution is needed. Matsueda et al. (2009) even found that a horizontal resolution of 20 km was required to accurately simulate the frequency of Euro-Atlantic blocking, and that higher resolution generally improves the representation in this sector. Further to the east, over the Eurasian continent, blocking is better simulated but exaggerated. Similar results are seen when blocking is defined at 50° N fixed latitude, although the bias is smaller in spring (not shown). For blocking in the NH summer and fall, the Euro-Atlantic blocking is better reproduced, but still underestimated.

Over the Pacific Ocean the simulated blocking frequency is closer to the observed, which is consistent with the conclusion by Matsueda et al. (2009) that the required horizontal resolution is coarser in the Pacific sector than in the Atlantic. It should be noted that there is some evidence that blocking deficits also can be reduced, even with relatively coarse resolution, by improving the SST field and reducing the time mean bias in the westerlies (Scaife et al., 2011).

The projected changes in blocking frequency during 2071–2100 for the RCP8.5 scenario compared to 1976–2005 are shown in Fig. 13. Using again the “vTM” index an increased blocking frequency is apparent for the sector 0–100° E for the summer months as well as in the sector 0–40° E during spring. We emphasize, however, that since there are large systematic biases in blocking frequency for the same sectors, these projections must be interpreted with great caution. The projected changes diagnosed using

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the index with fixed 50° N latitude (not shown), show considerably smaller changes than “vTM”, which may be due to the simultaneous poleward shift in the position of the westerlies. This contradicts the use of fixed latitudes for the blocking index. Such sensitivity to choice of index has also been reported by Barnes et al. (2012), and adds to the uncertainties connected with blocking simulation in climate models. The projected increase in blocking frequency can be linked to the reduced precipitation in the region except for Northern Europe.

6.3 NH EOF-analysis

In order to describe the low frequency variability in the NorESM1-M, an EOF (empirical orthogonal function) analysis has been applied to monthly mean 500 hPa geopotential height anomalies during extended winter seasons (DJFM) from 1976 to 2005, where the JFM-months define the year associated with each season. The three historical ensemble members for the mentioned period define the climatology about which the anomalies are calculated.

To calculate the EOFs (see e.g. Bjørnson, 1997; Hannachi, 2007; Monahan, 2009) the anomalies for each historical ensemble member are detrended by calculating deviations from a 5 yr moving average, and a common seasonal cycle for the simulated 1976–2005 period estimated by subtracting separate averages for each month (3 × 30 values are averaged per month). The EOFs thus represent spatial structures of the 500 hPa geopotential height fields associated with non-seasonal variations up to a few years, similar to the analysis of Corti et al. (1999) which was further extended by Molteni et al. (2006), based on NCEP re-analysis data (Kalnay et al., 1996).

The detrended, non-seasonal monthly anomalies are formally organized into a matrix **A**, where the rows are the 3 × 30 × 4 spatial anomaly patterns and the columns are the monthly values of the anomaly in each point in space. The EOFs are the eigenvectors (wrt. the standard Euclidian inner product) of the covariance matrix, $\mathbf{C} = \mathbf{A}^T \mathbf{A}$, i.e. $\mathbf{C}\mathbf{R} = \mathbf{R}^T \mathbf{A}$ where **A** is the diagonal matrix containing the eigenvalues, λ_i , of **C**. The column vectors in **R** are the mutually orthogonal eigenvectors *eof_i* of **C**. Each eigenvalue

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



measures the fraction of the total variance that the corresponding EOF pattern accounts for.

The associated principal components (PC) are time-series of the projections of \mathbf{A} onto the EOFs: $PC_i = \mathbf{A}^T \cdot \mathbf{eof}_i$. In addition to calculating the PCs for the three historical simulations for the DJFM-winters 1976–2005, the time-series of the projections of the non-detrended anomaly data on each EOF have been estimated for the GHG only (GHG), aerosol only (AER), and natural forcing only (NAT), as well as the projected scenario RCP8.5. Note that all anomalies are calculated w.r.t. the 1976–2005 climatology for the three historical ensemble members before projection onto the EOFs.

One reason for using EOF-based flow regime analysis in the context of climate change, is that internal dynamics in the climate system may determine the patterns of climate response to external forcing rather than the structure of the forcing itself (Palmer, 1999; Branstator and Selten, 2008; Iversen et al., 2008), although this view needs to be extended to account for local internal feedbacks in the climate system, which can be particularly strong in connection with Arctic snow and sea-ice (Boer and Yu, 2003).

The obtained projections onto selected EOFs for scenario simulations can therefore be directly compared with the PCs of the three historical ensemble members. If systematic differences are found, these can potentially be attributed to the differences in external conditions, whilst differences between the historical ensemble members' PCs will reflect internal variability. At present we have not estimated any probability density functions for projection coefficients; a method whose validity is under some discussion (Stephenson et al., 2004; Molteni et al., 2006).

The validity of potential results described above will not be convincing if the EOFs for the modelled fields differ considerably from “observed” EOFs from re-analysis data (e.g. Corti et al., 2003). One additional purpose of an analysis like this is therefore to investigate to what extent selected atmospheric flow regime patterns are reproduced in NorESM1-M. Hence we have also calculated EOFs to the NCEP reanalysis data for

GMDD

5, 2933–2998, 2012

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the same extended winter seasons of 1976–2005, in the same way as was done by Corti et al. (1999) and Molteni et al. (2006), but for a shorter period.

Figure 14 shows the 500 hPa geopotential height pattern for the 4 leading EOFs from the detrended monthly data for the three ensemble members of the historical simulations with NorESM1-M. Each EOF is scaled to represent one standard deviation of its principal component. They are then compared to the two leading EOFs calculated in the same way for the reanalysis data from NCEP for the same 30 winters (Kalnay et al., 1996). The first EOF for both the reanalysis and the model represents the Arctic Oscillation (AO), or alternatively the Northern Annular Mode (NAM). The shapes are slightly different in NorESM1-M with a maximum centred in the Central Arctic, while the NCEP data has a centre displaced over to the Atlantic sector, but both have secondary maxima of the opposite sign in Southern Europe, North-Eastern North America, and the Northern Pacific Ocean. These differences for EOF1 may be associated with systematic errors in the storminess patterns and the blocking occurrence over the North Atlantic Ocean and Europe mentioned in Sects. 6.1 and 6.2.

The second EOF of NorESM1-M differs significantly from that of the NCEP data. Both EOFs are combinations of patterns reminiscent of the Pacific North American (PNA) (Wallace and Gutzler, 1981) and the cold-ocean-warm-land (COWL) (Wallace et al., 1996) in combination with the North Atlantic Oscillation (NAO). However, while the NCEP EOF2 is dominated by the latter, the NorESM1-M EOF2 is dominated by the PNA. Furthermore, the NAO resemblance in NorESM1-M's EOF2 is weak and slightly displaced eastwards, while it turns up considerably more clearly in the fourth EOF. The third EOF of NorESM1-M resembles the EOF4 of NCEP (not shown).

Another difference between NorESM1-M and NCEP is revealed by the eigenvalues associated with the EOFs, i.e. the “explained variance” of each EOF as shown by the percentages above each EOF in Fig. 14. The low-order EOFs from the NorESM1-M data “explain” a larger portion than the corresponding low-order NCEP EOFs.

Figure 15 shows the time series of the decomposition of monthly 500 hPa geopotential height anomalies from NorESM1-M on the respective EOF1, EOF2 and EOF4

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with the sign shown in Fig. 14. The data are from the three historical ensemble members and the RCP8.5 scenario for the years 2071–2100. These data include the inter-annual (and longer) trends and the systematic differences between the historical and the scenario, but not the seasonal variations. Since all the anomalies are calculated with respect to the same climatology, defined by the three historical ensemble members for 1976–2005, a given value of the component identifies an exact monthly state, irrespective of the data source. The main curves in the diagrams are the 10-yr moving averages of the components. If a coloured curve lies outside of the range of the three historical ensemble members, this may indicate that the scenario assumption (RC8.5) leads to systematic differences from internal natural variability.

For PC1, this is clearly not the case, and neither for PC3 (not shown). For PC2 and PC4, however, there are indications of systematic differences, although less clear for the latter. One possible interpretation is that in the RCP8.5 climate towards the end of the 21st century, positive phase PNA may occur less frequently or the negative phase PNA may occur more often. Furthermore, but with less confidence, positive phase NAO may occur more often or negative phase NAO less frequently. More investigations of the significance of this and on probability density functions for the different combinations of PCs are ongoing.

Finally, Fig. 15 also includes the components of anomalies associated with the historical single forcing experiments. Even though there are some signs of opposite results for the GHG and the aerosol experiments, none of these are outside the ranges of natural variability defined by the three ensemble members with all forcing components included.

6.4 ENSO

The “El Niño Southern Oscillation” phenomenon is a dominant interannual regional weather pattern in the coupled atmosphere–ocean system (e.g. Trenberth, 1997). The amplitude (Trenberth and Shea, 1987; Wang, 1995), the frequency of occurrence, and the pattern structure (An and Wang, 2000) are modulated on multi-decadal timescales.

Nevertheless, it is a well recognized weather pattern with large impacts over the western equatorial South America. It also has considerable remote impacts (Trenberth et al., 1998) showing up as a pattern in the NH extratropical troposphere reminiscent of the PNA internal mode of variability although the patterns of the ENSO-response and the PNA are different (Straus and Shukla, 2002). The annual global mean surface air temperature is influenced by the ENSO phase.

Bentsen et al. (2012) show that for the NINO3.4 index NorESM1-M simulates variability on shorter time-scales (2–4 yr) than the HadISST observations (3–7 yr, Rayner et al., 2003). It has not been investigated to what extent this is dominated by model errors or if it can be related to interdecadal modes of variability (An and Wang, 2000). However, the recent analysis by Kim and Yu (2012) indicates that both modes of the ENSO variability are represented in the NorESM1-M simulations, as one of the 9 out of 20 CMIP5 models.

Figure 16 shows the time series of the NINO3.4 index for HadISST data, and from NorESM1-M, the piControl, the Historic1, and the RCP8.5. The more frequent recurrence in the NorESM1-M results for piControl and Historic1 is possible to recognize from the time series. A difference between the time series for RCP8.5 and either Hirstoric1 or piControl is less evident even though both amplitudes and return periods appear slightly reduced in the scenario. To the extent that it is correct to associate warm-phase ENSO with a positive PNA pattern, this result is consistent with the EOF-analysis in Sect. 6.3.

The spectra in Fig. 16b also indicate such changes. There are two peaks in the piControl, a primary peak around 3 yr and a secondary around 6–7 yr. Except for RCP4.5, the two peaks are less distinct in the scenarios. For RCP4.5 the two peaks appear distinct with a smaller difference between them, but both peaks occur at shorter periods than in piControl. There are also signs of less energy on periods longer than a decade for all RCPs except RCP6.0.

Further investigations with a larger number of ensemble members are required to establish the significance of these changes. The significant biases in the model

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



simulations also reduce the confidence in the changes, even though they are internally consistent and the preliminary study by Guilyardi et al. (2012) confirms that NorESM1-M is one of the two out of 14 CMIP5 models which simulate significantly reduced ENSO variability (in that case, the NINO3 index) for the abrupt $4 \times \text{CO}_2$ experiment, and close to significantly reduced for the gradual experiment.

7 Summary and conclusions

This paper presents a wide range of results of simulations with the new global climate model NorESM1-M. While the basics of the model predominantly is presented together with validation studies in the companion paper by Bentsen et al. (2012), we have presented and discussed different aspects of the model's properties concerning the climate sensitivity and response to prescribed changes that lead to radiative forcing. We believe that the results from CMIP5 experiments with NorESM1-M, which are only discussed to some extent in this paper, are valuable contributions to the development of climate system science, as well as to the total evaluation of possible human induced climate change. The data are open for anyone to download and analyze from <http://cmip-pcmdi.llnl.gov/cmip5/index.html>.

The clouds in the NorESM1-M tend to dampen the response to GHG forcing ($-0.09 \text{ W m}^{-2} \text{ K}^{-1}$), as the long-wave cloud response is considerably smaller than the negative short-wave response. The clear-sky response is negative ($-1.02 \text{ W m}^{-2} \text{ K}^{-1}$), in close agreement with other models (Andrews et al., 2012). The model has a strong Atlantic meridional overturning circulation (AMOC) of 30.8 Sv averaged over the piControl simulation period. This contributes to an efficient transport of heat into the deep oceans and reduces the heat available for increasing the surface temperature and to melt ice and snow. Exceptions may include regions adjacent to where major deep water formation takes place, where strong convergence of warm water in the upper ocean levels may occur, such as in the Atlantic sector of the Arctic and in Northwest Europe. The two estimates we have made of the equilibrium climate sensitivity are both

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



slightly lower than 2.9 K for a long-term adjustment to an abrupt doubling of CO₂. The transient climate sensitivity is estimated at slightly less than 1.4 K for gradual CO₂-increase until doubling. We argue that the latter may be an underestimate since very slow feedback processes may occur in response to a reduction of AMOC, which will gradually decrease the efficiency of deep ocean heat uptake. Nevertheless, NorESM1-M is amongst the less sensitive global climate models (Andrews et al., 2012).

We have also seen that the projections of global temperature increase based on RCP scenarios also are substantially smaller than most other global climate models contributing to the CMIP5 experiment, although inside one standard deviation below the ensemble mean. In addition to the low climate sensitivity, NorESM1-M also includes the predominant cooling of aerosols, both the direct effect and the indirect effects of pure water clouds, although their magnitude in the present model version is quite moderate (Kirkevåg et al., 2012). The geographical distribution of the heating shows the well-known response patterns of stronger warming over continents than oceans and a considerable amplification in the Arctic. The response in sea-ice is projected to be considerably larger in the Arctic than the Antarctic, and the extent in September is reduced to less than half by 2100 for three of the four RCP scenarios. For RCP8.5 this annual minimum is reduced to zero. Even though the winter maximum in March is relatively much less reduced, this will mean that the major extent of the winter Arctic sea-ice is generated during the same season.

For precipitation the largest response in the RCP scenarios towards the end of the 21st century is simulated at low latitudes, with an increase in the tropics and a decrease in the subtropics. In the extratropics and the high latitudes, precipitation is projected to increase, but in the NH summer the subtropical drying is extended northwards to mid-latitudes, including southern parts of North America and Europe. Projections of gross budget numbers in the atmospheric water cycle indicate intensification of all components except the evaporation from land. This reflects that the occurrence of both dry spells and high precipitation intensities increase over land, while widespread medium precipitation intensities probably occur less frequently.

**NorESM1-M –
climate response and
scenario projections**

T. Iversen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

NorESM1-M – climate response and scenario projections

T. Iversen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



The increased precipitation and sea-ice melting in the Arctic will influence the thermohaline forcing of the deep water formation, and thus the strength of the AMOC. All the RCP-scenarios are projected to lead to significantly reduced AMOC. RCP8.5 reduces AMOC by ca. one third by 2100, and in RCP4.5, which was run to 2300 with a stabilization scenario after 2100, the AMOC levels off at about 15–17 % lower intensity. Since NorESM1-M probably overestimates the AMOC speed, there are doubts associated with the quality of these results.

Precipitation and droughts in the extratropics is associated with storminess and blocking occurrence. The model is found to generally underestimate the variability in the zones of extratropical storminess, and a considerable part of this bias is linked to errors in the SST-simulations. The patterns of storminess are also too zonal over the North Atlantic Ocean. This is consistent with the too rare occurrence of blocking over the North Atlantic Ocean and Western Europe. Other recent investigations indicate that this can be related to too coarse horizontal resolution (Jung et al., 2012). Under RCP scenarios, NorESM1-M generally projects a northward displacement of the storminess. For blocking, an increased occurrence in the European-Atlantic sector in spring and further extended over Eurasia during summer is projected for the most extreme scenario (RCP8.5).

The mid-tropospheric pattern of the leading EOF during winter, representing the northern annular mode (NAM), has some pattern errors that can be associated with the systematic errors of the storminess. Furthermore, while the second EOF is dominated with a pattern reminiscent of the Pacific North American (PNA) pattern, the main influence of the North Atlantic Oscillation (NAO) turns up in the fourth EOF. From NCEP re-analysis data, the second EOF contains the main pattern of the NAO in addition to PNA. Under the RCP8.5 scenario towards 2100, there are indications that the average NAO index will increase (more positive and less negative events) and the average PNA index will decrease (more negative and less positive). The first result is consistent with the changes in storminess, while changes in blocking are insignificant in winter. The reduced PNA can be associated with the reduced amplitude of the projected NINO3.4

index, even though both the change in ENSO and the relation between ENSO and PNA are associated with low confidence.

Two of the single forcing experiments for the historical period have been addressed: the GHG only and the aerosol only. The response patterns by 1976–2005 compared to the pre-industrial control run are to a large extent similar but with opposite sign. This is also true for precipitation, but with important exceptions at low latitudes. The model simulates a southward displacement of ITCZ due to aerosol forcing, and in particular during NH summer. This change may partly cancel and partly reinforce changes caused by GHG alone. Reinforcements are simulated in the northern hemispheric subtropics with reduced precipitation and increased droughts, and at the southern flank of the ITCZ in the Pacific Ocean with increased precipitation. In consequence, the double ITCZ systematic error seen in many models (also CCSM4, Gent et al., 2011) is reduced in NorESM1-M partly due to the impact of aerosols (see Bentsen et al., 2012).

Acknowledgements. This work has been supported by the Research Council of Norway through the EarthClim (207711/E10) and NOTUR/NorStore projects, the Center for Climate Dynamics at the Bjerknes Centre, and through the European Commission FP7 projects PEGASOS (FP7-ENV-2010–265148) and ACCESS (FP7-ENV-2010-265863). The development of NorESM has been possible because of the granted early access to the later public versions of the CCSM4 and CESM1. We are particularly grateful to P. J. Rasch, A. Gettelman, J. F. Lamarque, S. Ghan, M. Vertenstein, B. Eaton, M. Flanner, and others, for invaluable advice on numerous scientific and technical issues, and the support by the CESM program directors during the development period, P. Gent and J. Hurrell. We are grateful for contributions during different versions of NorESM by K. Alterskjær, A. Ekman, C. Heinze, C. Hoose, Ø. Hov, M. Ø. Køltzow, D. Nilsson, O. H. Otterå, D. Olivié, C. Roelandt, M. Simonsen, H. Struthers, T. Storelvmo, J. Berger, J. Su, and J. Tjiputra.

Section 6.3 in this paper is intended to be part of a PhD thesis at the University of Oslo (M. Sand) in agreement with all the co-authors.

GMDD

5, 2933–2998, 2012

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Adler, R. F., Huffman, G. J., Chang, A., Ferraro, R., Xie, P., Janowiak, J., Rudolf, B., Schneider, U., Curtis, S., Bolvin, D., Gruber, A., Susskind, J., and Arkin, P.: The Version 2 Global Precipitation Climatology Project (GPCP) monthly precipitation analysis (1979–present), *J. Hydrometeorol.*, 4, 1147–1167, 2003.
- Ammann, C. M., Meehl, G. A., Washington, W. M., and Zender, C.: A monthly and latitudinally varying volcanic forcing dataset in simulations of 20th century climate, *Geophys. Res. Lett.*, 30, 1657, doi:10.1029/2003GL016875, 2003.
- An, S.-I. and Wang, B.: Interdecadal change of the structure of the ENSO mode and its impact on the ENSO frequency, *J. Climate*, 13, 2044–2055, 2000.
- Andrews, T., Gregory, J. M., Webb, M. J., and Taylor, K. E.: Forcing, feedbacks and climate sensitivity in CMIP5 coupled atmosphere–ocean climate models, *Geophys. Res. Lett.*, 39, L09712, doi:10.1029/2012gl051607, 2012.
- Assmann, K. M., Bentsen, M., Segschneider, J., and Heinze, C.: An isopycnic ocean carbon cycle model, *Geosci. Model Dev.*, 3, 143–167, doi:10.5194/gmd-3-143-2010, 2010.
- Barnes, E. A., Slingo, J., and Woollings, T. J.: A methodology for the comparison of blocking climatologies across indices, models and climate scenarios, *Clim. Dynam.*, 38, 2467–2481, doi:10.1007/s00382-011-1243-6, 2012.
- Bentsen, M., Bethke, I., Debernard, J. B., Iversen, T., Kirkevåg, A., Seland, Ø., Drange, H., Roe-landt, C., Seierstad, I. A., Hoose, C., and Kristjánsson, J. E.: The Norwegian Earth System Model, NorESM1-M – Part 1: Description and basic evaluation, *Geosci. Model Dev. Discuss.*, 5, 2843–2931, doi:10.5194/gmdd-5-2843-2012, 2012.
- Bethke, I., Furevik, T., and Drange, H.: Towards a more saline North Atlantic and a fresher Arctic under global warming, *Geophys. Res. Lett.*, 33, L21712, doi:10.1029/2006GL027264, 2006.
- Bitz, C. M., Shell, K. M., Gent, P. R., Bailey, D. A., Danabasoglu, G., Armour, K. C., Holland, M. M., and Kiehl, J. T.: Climate Sensitivity of the Community Climate System Model, version 4, *J. Climate*, 25, 3053–3070, doi:10.1175/JCLI-D-11-00290.1, 2012.
- Björnsson, H. and Venegas, S. A.: A manual for EOF and SVD analyses of climate data, McGill University, CCGCR Report No. 97-1, Montréal, Québec, 52 pp., 1997.
- Blackmon, M. L.: A climatological spectral study of the 500 mb geopotential height of the Northern Hemisphere, *J. Atmos. Sci.*, 33, 1607–1623, 1976.

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Boer, G. J. and Yu, B.: Climate sensitivity and response, *Clim. Dynam.*, 20, 415–429, doi:10.1007/s00382-002-0283-3, 2003.
- Branstator, G. and Selten, F.: “Modes of variability” and climate change, *J. Climate*, 22, 2639–2658, doi:10.1175/2008JCLI2517.1, 2009.
- 5 Bratseth, A. M.: Eddy acceleration of the mean flow in the atmosphere, *J. Atmos. Sci.*, 58, 3328–3339, 2001.
- Bratseth, A. M.: Zonal-mean transport characteristics of ECMWF re-analysis data, *Q. J. Roy. Meteorol. Soc.*, 129, 2331–2346. doi:10.1256/qj.02.90, 2003.
- 10 Chang, E. K. M., Lee, S., and Swanson, K. L.: Storm track dynamics, *J. Climate*, 15, 2163–2183, 2002.
- Corti, S., Molteni, F., and Palmer, T. N.: Signature of recent climate change in frequencies of natural atmospheric circulation regimes, *Nature*, 398, 799–802, 1999.
- Corti, S., Gualdi, S., and Navarra, A.: Analysis of the mid-latitude weather regimes in the 200-year control integration of the SINTEX model, *Ann. Geophys.*, 46, 27–37, 2003, <http://www.ann-geophys.net/46/27/2003/>.
- 15 Curry, R. and Mauritzen, C.: Dilution of the Northern North Atlantic Ocean in recent decades, *Science*, 308, 1772–1774, doi:10.1126/science.1109477, 2005.
- D’Andrea, F., Tibaldi, S., Blackburn, M., Boer, G., Deque, M., Dix, M. R., Dugas, B., Ferranti, L., Iwasaki, T., Kitoh, A., Pope, V., Randall, D., Roeckner, E., Straus, D., Stern, W., Van den Dool, H., and Williamson, D.: Northern Hemisphere atmospheric blocking as simulated by 15 atmospheric general circulation models in the period 1979–1988, *Clim. Dynam.*, 14, 385–407. 1998.
- 20 Fetterer, F., Knowles, K., Meier, W., and Savoie, M.: Sea Ice Index, National Snow and Ice Data Center, Boulder, CO, digital media, 2009.
- Flanner, M. G. and Zender, C. S.: Linking snowpack microphysics and albedo evolution, *J. Geophys. Res.*, 111, D12208, doi:10.1029/2005JD006834, 2006.
- Furevik, T., Bentsen, M., Drange, H., Kindem, I. K. T., Kvamstø, N. G., and Sorteberg, A.: Description and evaluation of the bergen climate model: ARPEGE coupled with MICOM, *Clim. Dynam.*, 21, 27–51, doi:10.1007/s00382-003-0317-5, 2003.
- 30 Gent, P. R., Danabasoglu, G., Donner, L. J., Holland, M. M., Hunke, E. C., Jayne, S. R., Lawrence, D. M., Neale, R. B., Rasch, P. J., Vertenstein, M., Worley, P. H., Yang, Z.-L., and Zhang, M.: The Community Climate System Model version 4, *J. Climate*, 24, 4973–4991, doi:10.1175/2011JCLI4083.1, 2011.

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Gettelman, A., Kay, J. E., and Shell, K. M.: The evolution of climate sensitivity and climate feedbacks in the Community Atmosphere Model, *J. Climate*, 25, 1453–1469, doi:10.1175/JCLI-D-11-00197.1, 2012.

Ghil, M., Allen, R. M., Dettinger, M. D., Ide, K., Kondrashov, D., Mann, M. E., Robertson, A., Saunders, A., Tian, Y., Varadi, F., and Yiou, P.: Advanced spectral methods for climatic time series, *Rev. Geophys.*, 40, 1–41, doi:10.1029/2000RG000092, 2002.

Gimeno, L., Nieto, R., Drumond, A., Durán-Quesada, A. M., Stohl, A., Sodemann, H., and Trigo, R. M.: A close look at oceanic sources of continental precipitation, *Eos Trans. AGU*, 92, 193–195, 2011.

Giorgi, F., Im, E.-S., Coppola, E., Diffenbaugh, N. S., Gao, X. J., Mariotti, L., and Shi, Y.: Higher hydroclimatic intensity with global warming, *J. Climate*, 24, 5309–5324, doi:10.1175/2011JCLI3979.1, 2011.

Gregory, J. M., Ingram, W. J., Palmer, M. A., Jones, G. S., Stott, P. A., Thorpe, R. B., Lowe, J. A., Johns, T. C., and Williams, K. D.: A new method for diagnosing radiative forcing and climate sensitivity, *Geophys. Res. Lett.*, 31, L03205, doi:10.1029/2003GL018747, 2004.

Guilyardi, E., Bellenger, H., Collins, M., Ferrett, S., Cai, W., and Wittenberg, A.: A first look at ENSO in CMIP5, CLIVAR Exchanges no. 58, 17, 29–32, available at: http://www.gfdl.noaa.gov/~atw/yr/2012/guilyardi.etal_2012_clivex.pdf (last access: February 2012), 2012

Hannachi, A., Jolliffe, I. T., and Stephenson, D. B.: Empirical orthogonal functions and related techniques in atmospheric science: a review, *Int. J. Climate*, 27, 1119–1152, 2007.

Hofmann, M. and Rahmstorf, S.: On the stability of the Atlantic meridional overturning circulation, *Proc. Natl. Acad. Sci.*, 106, 20584–20589, 2009.

Hoose, C., Kristjánsson, J. E., Iversen, T., Kirkevåg, A., Seland, Ø., and Gettelman, A.: Constraining cloud droplet number concentration in GCMs suppresses the aerosol indirect effect, *Geophys. Res. Lett.*, 36, L12807, doi:10.1029/2009GL038568, 2009.

Houghton, J. T., Ding, Y., Griggs, D. J., Noguera, M., van der Linden, P. J., Dai, X., Maskell, K., and Johnson, C. A.: *Climate Change 2001: The Scientific Basis*, Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, 881 pp., 2001.

Hurrell, J. W., Hack, J. J., Phillips, A. S., Caron, J., and Yin, J.: The dynamical simulation of the Community Atmosphere Model version 3 (CAM3), *J. Climate*, 19, 2162–2183, 2006.

NorESM1-M – climate response and scenario projections

T. Iversen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Iversen, T. and Seland, Ø.: A scheme for process-tagged SO₄ and BC aerosols in NCAR-CCM3, validation and sensitivity to cloud processes, *J. Geophys. Res.*, 107, D244751, doi:10.1029/2001JD000885, 2002.

Iversen, T. and Seland, Ø.: Correction to “A scheme for process-tagged SO₄ and BC aerosols in NCAR-CCM3, validation and sensitivity to cloud processes”, *J. Geophys. Res.*, 108, D164502, doi:10.1029/2003JD003840, 2003.

Iversen, T., Kristiansen, J., Jung, T., and Barkmeijer, J.: Optimal atmospheric forcing perturbations for the cold-ocean warm-land pattern, *Tellus A*, 60, 528–546, doi:10.1111/j.1600-0870.2008.00310.x, 2008.

Jiang, J.H., Su, H., Zhai, C., Perun, V., Del Genio, A. D., Nazarenko, L. S., Donner, L. J., Horowitz, L. W., Seman, C. J., Cole, J., Gettelman, A., Ringer, M. A., Rotstayn, L.D., Jeffrey, S. J., Wu, T., Brient, F., Dufresne, J.-L., Kawai, H., Koshiro, T., Masahiro, W., L'Écuyer, T. S., Volodin, E. M., Iversen, T., Drange, H., dos Santos Mesquita, M., Read, W. G., Waters, J. W., Tian, B., Teixeira, J., and Stephens, G. L.: Evaluation of cloud and water vapor simulations in CMIP5 climate models using NASA “A-Train” satellite observations, *J. Geophys. Res.*, 117, D14105, doi:10.1029/2011JD017237, 2012.

Jung, T., Miller, M. J., Palmer, T. N., Towers, P., Wedi, N., Achuthavari, D., Adams, J. M., Alshuler, E. L., Cash, B. A., Kinter III, J. L., Marx, L., Stan, C., and Hodges, K. I.: High-resolution global climate simulations with the ECMWF model in project Athena: experimental design, model climate, and seasonal forecast skill, *J. Climate*, 25, 3155–3172. doi:10.1175/JCLI-D-11-00265.1, 2012.

Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–470, 1996.

Kay, J. E., Holland, M. M., Bitz, C. M., Blanchard-Wrigglesworth, E., Gettelman, A., Conley, A., and Bailey, D.: The influence of local feedbacks and northward heat transport on the equilibrium Arctic climate response to increased greenhouse gas forcing, *J. Climate*, 25, 5433–5450, doi:10.1175/JCLI-D-11-00622.1, 2012.

Kim, S. T. and Yu, J.-Y.: The two types of ENSO in CMIP5 models, *Geophys. Res. Lett.*, 39, L11704, doi:10.1029/2012GL052006, 2012.

Kirkevåg, A. and Iversen, T.: Global direct radiative forcing by process-parameterized aerosol optical properties, *J. Geophys. Res.*, 107, D204433, doi:10.1029/2001JD000886, 2002.

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kirkevåg, A., Iversen, T., Seland, Ø., Debernard, J. B., Storelvmo, T., and Kristjánsson, J. E.: Aerosol-cloud-climate interactions in the climate model CAM-Oslo, *Tellus A*, 60, 492–512, doi:10.1111/j.1600-0870.2008.00313.x, 2008a.
- 5 Kirkevåg, A., Iversen, T., Kristjánsson, J. E., Seland, Ø., and Debernard, J. B.: On the additivity of climate response to anthropogenic aerosols and CO₂, and the enhancement of future global warming by carbonaceous aerosols, *Tellus A*, 60, 513–527, doi:10.1111/j.1600-0870.2008.00308.x, 2008b.
- Kirkevåg, A., Iversen, T., Seland, Ø., Hoose, C., Kristjánsson, J. E., Struthers, H., Ekman, A. M. L., Ghan, S., Griesfeller, J., Nilsson, E. D., and Schulz, M.: Aerosol-climate interactions in the Norwegian Earth System Model – NorESM, *Geosci. Model Dev. Discuss.*, 5, 2599–2685, doi:10.5194/gmdd-5-2599-2012, 2012.
- 10 Kristjánsson, J. E.: Studies of the aerosol indirect effect from sulphate and black carbon aerosols, *J. Geophys. Res.*, 107, D154246, doi:10.1029/2001JD000887, 2002.
- Kristjánsson, J. E., Iversen, T., Kirkevåg, A., Seland, Ø., and Debernard, J. B.: Response of the climate system to aerosol direct and indirect forcing: role of cloud feedbacks, *J. Geophys. Res.*, 110, D24206, doi:10.1029/2005JD006299, 2005.
- 15 Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.
- Lean, J., Rottman, G., Harder, J., and Kopp, G.: *SORCE* contributions to new understanding of global change and solar variability, *Sol. Phys.*, 230, 27–53, 2005.
- 25 Lejanäs, H. and Økland, H.: Characteristics of Northern Hemisphere blocking as determined from a long series of observational data, *Tellus A*, 35, 350–362, 1983.
- Matsueda, M., Mizuta, R., and Kusunoki, S.: Future change in wintertime atmospheric blocking simulated using a 20-km-mesh atmospheric global circulation model, *J. Geophys. Res.*, 114, D12114, doi:10.1029/2009JD011919, 2009.
- 30 Medhaug, I. and Furevik, T.: North Atlantic 20th century multidecadal variability in coupled climate models: sea surface temperature and ocean overturning circulation, *Ocean Sci.*, 7, 389–404, doi:10.5194/os-7-389-2011, 2011.

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Meehl, G. A., Covey, C., McAvaney, B., Latif, M., and Stouffer, R. J.: Overview of the coupled model intercomparison project, *B. Am. Meteorol. Soc.*, 86, 89–93, doi:10.1175/BAMS-86-1-89, 2005.
- Meehl, G. A., Washington, W. M., Arblaster, J. M., Hu, A., Teng, H., Tebaldi, C., Sanderson, B., Lamarque, J.-F., Conley, A., Strand, W. G., and White III, J. B.: Climate system response to external forcings and climate change projections in CCSM4, *J. Climate*, 25, 3661–3683, doi:10.1175/JCLI-D-11-00240.1, 2012.
- Molteni, F., Kucharski, F., and Corti, S.: On the predictability of flow-regime properties on interannual to interdecadal timescales, in: *Predictability of Weather and Climate*, edited by: Palmer, T. and Hagedorn, R., Cambridge University Press, 2006, 365–389, 2006.
- Monahan, A. H., Fyfe, J. C., Ambaum, M. H. P., Stephenson, D. B., and North, G. R.: Empirical orthogonal functions: the medium is the message, *J. Climate*, 22, 6501–6514, 2009.
- Munk, W. and Wunsch, C.: Abyssal recipes II: energetics of tidal and wind mixing, *Deep-Sea Res. Pt. I*, 45, 1977–2010, 1998.
- Murphy, D. M.: Transient response of the Hadley Centre coupled ocean-atmosphere model to increasing carbon dioxide: part III. analysis of global-mean response using simple models, *J. Climate*, 8, 495–514, doi:10.1175/1520-0442(1995)008<0496:TROTHC>2.0.CO;2, 1995.
- Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N., Chin, M., Diehl, T., Easter, R. C., Feichter, J., Ghan, S. J., Hauglustaine, D., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Luo, G., Ma, X., Penner, J. E., Rasch, P. J., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., Wang, Z., Xu, L., Yu, H., Yu, F., Yoon, J.-H., Zhang, K., Zhang, H., and Zhou, C.: Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations, *Atmos. Chem. Phys. Discuss.*, 12, 22355–22413, doi:10.5194/acpd-12-22355-2012, 2012.
- Neale, R. B., Richter, J. H., and Jochum, M.: The impact of convection on ENSO: from a delayed oscillator to a series of events, *J. Climate*, 21, 5904–5924, 2008.
- Neale, R. B., Richter, J. H., Conley, A. J., Park, S., Lauritzen, P. H., Gettelman, A., Williamson, D. L., Rasch, P. J., Vavrus, S. J., Taylor, M. A., Collins, W. D., Zhang, M., and Lin, S.-J.: Description of the NCAR Community Atmosphere Model (CAM 4.0), NCAR technical note, April, 2010.
- Neale, R. B., Richter, J. H., Park, S., Lauritzen, P. H., Vavrus, S. J., Rasch, P. J., and Zhang, M.: The mean climate of the Community Atmosphere Model (CAM4) in forced SST and fully coupled experiments, *J. Climate*, submitted, 2012.

NorESM1-M – climate response and scenario projections

T. Iversen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Otterå, O. H., Bentsen, M., Bethke, I., and Kvamstø, N. G.: Simulated pre-industrial climate in Bergen Climate Model (version 2): model description and large-scale circulation features, *Geosci. Model Dev.*, 2, 197–212, doi:10.5194/gmd-2-197-2009, 2009.
- Otterå, O. H., Bentsen, M., Drange, H., and Suo, L.: External forcing as a metronome for Atlantic multidecadal variability, *Nat. Geosci.*, 3, 688–694, doi:10.1038/NGEO955, 2010.
- Palmer, T. N.: A nonlinear dynamical perspective on climate prediction, *J. Climate*, 12, 575–591, 1998.
- Pelly, J. L. and Hoskins, B. J.: A new perspective on blocking, *J. Atmos. Sci.*, 60, 743–755, 2003.
- Rasch, P. J. and Kristjánsson, J. E.: A comparison of the CCM3 model climate using diagnosed and predicted condensate parameterizations, *J. Climate*, 11, 1587–1614, 1998.
- Rasch, P. J., Coleman, D. B., Mahowald, N., Williamson, D. L., Lin, S.-J., Boville, B. A., and Hess, P.: Characteristics of atmospheric transport using three numerical formulations for atmospheric dynamics in a single GCM framework, *J. Climate*, 19, 2243–2266, 2006.
- Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, *J. Geophys. Res.*, 108, 4407, doi:10.1029/2002JD002670, 2003.
- Richter, J. H. and Rasch, P. J.: Effects of convective momentum transport on the atmospheric circulation in the Community Atmosphere Model, version 3, *J. Climate*, 21, 1487–1499, 2008.
- Rotstayn, L. D. and Lohmann, U.: Tropical rainfall trends and the indirect aerosol effect, *J. Climate*, 15, 2103–2116, 2002.
- Scaife, A. A., Copsey, D., Gordon, C., Harris, C., Hinton, T., Keeley, S., O'Neill, A., Roberts, M., and Williams, K.: Improved Atlantic winter blocking in a climate model, *Geophys. Res. Lett.*, 38, L23703, doi:10.1029/2011GL049573, 2011.
- Schneider, B., Latif, M., and Schmittner, A.: Evaluation of different methods to assess model projections of the future evolution of the Atlantic meridional overturning circulation, *J. Climate*, 20, 2121–2132, 2007.
- Seland, Ø., Iversen, T., Kirkevåg, A., and Storelvmo, T.: Aerosol-climate interactions in the CAM-Oslo atmospheric GCM and investigations of associated shortcomings, *Tellus A*, 60, 459–491, 2008.

NorESM1-M – climate response and scenario projections

T. Iversen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ciated with tropical sea surface temperatures, *J. Geophys. Res.*, 103, 14291–14324, doi:10.1029/97JC01444, 1998.

Trenberth, K. E., Dai, A., Rasmussen, R. M., and Parsons, D. B.: The changing character of precipitation, *B. Am. Meteorol. Soc.*, 84, 1205–1217, 2003.

5 Trenberth, K.E., Jones, P. D., Ambenje, P., Bojariu, R., Easterling, D., Klein Tank, A., Parker, D., Rahimzadeh, F., Renwick, J. A. Rusticucci, M., Soden, B., and Zhai, P.: Observations: surface and atmospheric climate change, in: *Climate Change 2007: The Physical Science Basis*, Intergovernmental Panel on Climate Change 4th Assessment Report, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z. Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, 235–336, 2007.

10 Trenberth, K. E., Fasullo, J. T., and Mackaro, J.: Atmospheric moisture transports from ocean to land and global energy flows in reanalyses, *J. Climate*, 24, 4907–4924, doi:10.1175/2011JCLI4171.1, 2011.

15 Uppala, S. M., Kallberg, P. W., Simmons, A. J., Andrae, U., Bechtold, V. D., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., Van De Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Holm, E., Hoskins, B. J., Isaksen, L., Janssen, P. A. E. M., Jenne, R., McNally, A. P., Mahfouf, J. F., Morcrette, J. J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J. T.: The ERA-40 re-analysis, *Q. J. Roy. Meteorol. Soc.*, 131, 2961–3012, 2005.

20 Yin, J. H.: A consistent poleward shift of the storm tracks in simulations of 21st century climate, *Geophys. Res. Lett.*, 32, L18701, doi:10.1029/2005GL023684, 2005.

25 Van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., and Rose, S. K.: The representative concentration pathways: an overview, *Climatic Change*, 109, 5–31, doi:10.1007/s10584-011-0148-z, 2011.

Wallace, J. M. and Gutzler, D. S.: Teleconnections in the geopotential height field during the Northern Hemisphere winter, *Mon. Weather Rev.*, 109, 784–812, 1981.

30 Wallace, J. M., Zhang, Y., and Bajuk, L.: Interpretation of interdecadal trends in Northern Hemispheric surface air temperature, *J. Climate*, 9, 249–259, 1996.

Wang, B.: Interdecadal changes in El Niño onset in the last four decades, *J. Climate*, 8, 267–285, 1995.

Wang, Y.-M., Lean, J. L., and Sheeley Jr., N. R.: Modeling the Sun's magnetic field and irradiance since 1713, *Astrophys. J.*, 625, 522–538, 2005.

Zhang, G. J. and McFarlane, N. A.: Sensitivity of climate simulations to the parameterization of cumulus convection in the Canadian Climate Centre general circulation model, *Atmos. Ocean*, 33, 407–446, 1995.

5

GMDD

5, 2933–2998, 2012

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NorESM1-M – climate response and scenario projections

T. Iversen et al.

Table 2. Global gross feedback response (λ_X) in TOA radiation parameters (X) as determined by linear regression of model simulated annual change (ΔX) with respect to the corresponding annual surface air temperature change (ΔT) after abrupt $4 \times \text{CO}_2$. The quantity $\lambda_X = d(\Delta X)/d(\Delta T)$, and X is long-wave (LW) and short-wave (SW) all-sky and clear-sky TOA outgoing radiation, long-wave (LWCF) and short-wave (SWCF) cloud forcing, or net cloud radiative effect (CRE).

	$\lambda_{\text{LWAllsky}}$ $\text{Wm}^{-2}\text{K}^{-1}$	$\lambda_{\text{SWAllsky}}$ $\text{Wm}^{-2}\text{K}^{-1}$	$\lambda_{\text{LWclearsky}}$ $\text{Wm}^{-2}\text{K}^{-1}$	$\lambda_{\text{SWclearsky}}$ $\text{Wm}^{-2}\text{K}^{-1}$	λ_{LWCF} $\text{Wm}^{-2}\text{K}^{-1}$	λ_{SWCF} $\text{Wm}^{-2}\text{K}^{-1}$	λ_{CRE} $\text{Wm}^{-2}\text{K}^{-1}$
NorESM1-M, 2 deg.	-1.80	+0.70	-1.86	+0.84	+0.06	-0.15	-0.09

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

NorESM1-M – climate response and scenario projections

T. Iversen et al.

Table 3. Global gross feedback response (λ_X) in parameters (X) characterizing the hydroclimate, as determined by linear regression of model simulated annual change (ΔX) with respect to the corresponding annual surface air temperature change (ΔT) after abrupt $4 \times \text{CO}_2$. X is annual amounts of evaporation (E), precipitation (P), or the difference ($E - P$) accumulated globally, from oceans, or from land ($10^3 \text{ km}^3 \text{ yr}^{-1}$).

	$\lambda_{P\text{-GLOB}}$ $10^3 \text{ km}^3 \text{ K}^{-1}$	$\lambda_{E\text{-OCEAN}}$ $10^3 \text{ km}^3 \text{ K}^{-1}$	$\lambda_{P\text{-OCEAN}}$ $10^3 \text{ km}^3 \text{ K}^{-1}$	$\lambda_{(E-P)\text{-OCEAN}}$ $10^3 \text{ km}^3 \text{ K}^{-1}$	$\lambda_{E\text{-LAND}}$ $10^3 \text{ km}^3 \text{ K}^{-1}$	$\lambda_{P\text{-LAND}}$ $10^3 \text{ km}^3 \text{ K}^{-1}$
NorESM1-M, 2 deg.	14.58	12.42	12.40	+0.02 (−0.29+0.32)	2.16	2.18

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


NorESM1-M – climate response and scenario projections

T. Iversen et al.

Table 4. Global gross feedback response (λ_X) in ($X =$) yearly averaged sea-ice area (AREA, $10^6 \text{ km}^2 \text{ yr}^{-1}$), and volume (VOL, $10^3 \text{ km}^3 \text{ yr}^{-1}$) in the northern (NH) or southern (SH) hemispheres, as determined by linear regression of model simulated annual change (ΔX) with respect to the corresponding annual surface air temperature change (ΔT) after abrupt $4 \times \text{CO}_2$.

	$\lambda_{\text{AREA-NH}}$ $10^6 \text{ km}^2 \text{ K}^{-1} \text{ yr}^{-1}$	$\lambda_{\text{VOL-NH}}$ $10^3 \text{ km}^3 \text{ K}^{-1} \text{ yr}^{-1}$	$\lambda_{\text{AREA-SH}}$ $10^6 \text{ km}^2 \text{ K}^{-1} \text{ yr}^{-1}$	$\lambda_{\text{VOL-NH}}$ $10^3 \text{ km}^3 \text{ K}^{-1} \text{ yr}^{-1}$
NorESM1-M, 2 deg.	-2.39	-10.55	-0.86	-2.52

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


NorESM1-M – climate response and scenario projections

T. Iversen et al.

Table 5. Simulated changes in selected global annual data with NorESM1-M from the period 1976–2005 (Historic1) to 2071–2100 based on the four projected representative concentration pathways (RCP) scenarios. The hydro-climatic quantities marked with a star (*) are estimated using the fact that $E_{\text{GLOBAL}} = P_{\text{GLOBAL}}$ in the model. The rightmost columns contain total values simulated for 1976–2005 (Historic1), and the difference between this and piControl.

	RCP8.5 – Historic1	RCP6.0 – Historic1	RCP4.5 – Historic1	RCP2.6 – Historic1	Historic1 1976–2005	Historic1 – piControl
T_{2m} K	+3.07	+1.86	+1.65	+0.94	286.78	+0.50
SST K	+1.76	+1.06	+0.95	+0.59	282.92	+0.34
AREA _{SeaIce} 10 ⁶ km ²	–6.24	–3.48	–2.97	–1.43	20.76	–1.14
P_{GLOBAL} 1000 km ³ yr ^{–1}	+27	+17	+17	+12	521	0
E_{OCEANS} 1000 km ³ yr ^{–1}	+25	+15	+14	+10	442	+1
$(E - P)_{\text{OCEANS}}$ 1000 km ³ yr ^{–1}	+8	+4	+2	+1	43	+1
P_{OCEANS}^* 1000 km ³ yr ^{–1}	+17	+11	+12	+9	399	0
P_{LAND}^* 1000 km ³ yr ^{–1}	+10	+6	+5	+3	122	0
E_{LAND}^* 1000 km ³ yr ^{–1}	+2	+2	+3	+2	79	–1

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


NorESM1-M – climate response and scenario projections

T. Iversen et al.

Table 6. Simulated changes in annual total precipitation (P , mm yr^{-1}) and annual total runoff (R , mm yr^{-1}) with NorESM1-M in Europe, Northern Europe, and the Mediterranean region, from the period 1976–2005 (Historic1) to 2071–2100 based on the four projected representative concentration pathways (RCP) scenarios. The rightmost columns contain total values simulated for 1976–2005 (Historic1), and the difference between this and piControl.

		RCP8.5 – Historic1	RCP6.0 – Historic1	RCP4.5 – Historic1	RCP2.6 – Historic1	Historic1 1976–2005	Historic1 – piControl
Europe	$P_{\text{EUR}}/$ mm yr^{-1}	+32.3	+12.0	+25.5	+42.1	862.1	–19.8
	$R_{\text{EUR}}/$ mm yr^{-1}	–11.8	–16.5	–10.1	+5.7	316.4	–2.0
North. Eur.	$P_{\text{N-EUR}}/$ mm yr^{-1}	+81.9	+61.2	+53.5	+18.9	723.1	+43.3
	$R_{\text{N-EUR}}/$ mm yr^{-1}	+24.1	+21.8	+11.6	–10.3	468.4	+40.6
Mediterr.	$P_{\text{MED}}/$ mm yr^{-1}	–79.7	–36.3	–5.8	+20.6	611.5	–41.1
	$R_{\text{MED}}/$ mm yr^{-1}	–25.3	–12.2	–3.6	+3.9	90.3	–9.77

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NorESM1-M –
climate response and
scenario projections

T. Iversen et al.




Fig. 1. Aspects of climate sensitivity and gross feedback effects in NorESM1-M based on the experiments “abrupt $4 \times CO_2$ ” and “gradual $4 \times CO_2$ ”. **(a)** Model simulated change in yearly TOA net downward radiation (Wm^{-2}) as a function of changed global surface air temperature (K) (black dots) and effective temperature response, ΔT_{eff} (n) (red dots, $n = 111, \dots, 150$), for the abrupt $4 \times CO_2$ experiment. The black line is the linear regression with a slope $\lambda = 1.101 W m^{-2} K^{-1}$ and intercept $\Delta T_{reg}(4 \times CO_2) = 5.74 K$ with the x-axis. The black cross is the average for the red dots with $\Delta T_{eff}(4 \times CO_2) = 5.71 K$. **(b)** Calculated changes in TOA long-wave (blue), short wave (red) and net (black) downward cloud radiation (Wm^{-2}), as a function of changed global surface air temperature for the abrupt $4 \times CO_2$ experiment. The slopes of the linear regression lines are given in Table 2. **(c)** The maximum AMOC (Sv) at $26.5^\circ N$ as a function of time for piControl (grey), abrupt $4 \times CO_2$ (blue) and gradual $4 \times CO_2$ (red). **(d)** Changed temperature zonally averaged for global oceans for the gradual $4 \times CO_2$ experiment at the time of CO_2 doubling. **(e)** The global TOA radiation heat flux as a function of time in the gradual $4 \times CO_2$ experiment along with the globally averaged downward flux of heat through depth levels in the world oceans. **(f)** Same as **(e)** but for the abrupt $4 \times CO_2$ experiment.




Fig. 2. Globally and annually averaged aerosol column burdens for particulate organic matter (POM), particulate sulphate as S ($\text{SO}_4\text{-S}$), and black carbon (BC) (upper panel), aerosol optical depth (AOD) (middle panel), and aerosol optical depth for absorption (AAOD) (lower panel) from 1850 onwards, calculated online in NorESM1-M. Curves for Historic1 from 1850 to 2005 are black. For scenario projections, green are RCP2.6 for 2005–2100, blue are RCP4.5 for 2005–2300 (negligible variations after 2150), orange are RCP6.0 for 2005–2100, and red are RCP8.5 for 2005–2100. The brown curves are contributions to AOD and AAOD in Historic1 by natural aerosols only, including prescribed stratospheric sulphate from explosive volcanoes.

**NorESM1-M –
climate response and
scenario projections**

T. Iversen et al.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



NorESM1-M –
climate response and
scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. From the top panel and downwards, the figure shows the net global long-wave (positive upwards), short-wave (positive downwards), and total (positive downwards) radiative flux at the top of the atmosphere during the NorESM1-M simulations for 1850 to 2300. The next two panels show diagrams for the global surface air temperature and average daily precipitation. Black: Historic1; green: RCP2.6; blue: RCP4.5; orange: RCP 6.0; red: RCP8.5.




Fig. 4. NorESM1-M single forcing simulations of the historical period 1850–2005 and for the period 1976–2005 compared to piControl. Response in annual mean surface air temperature (left panels) and average daily precipitation amounts (right panels). Upper panels shows global annual from 1850 to 2005 for Historic1, 2, and 3 with full forcing (black), with natural forcing only (green), GHG only (red), and aerosols only (blue). The panels in the middle show changes between piControl and 1976–2005 for GHG only, while the lower panels show corresponding results for aerosols only.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NorESM1-M –
climate response and
scenario projections

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 5. Model simulated change in average daily precipitation during 1976–2005 relative to piControl for the historical forcing experiments “GHG only” and “aerosol only”. Upper panels: zonally averaged changes in daily precipitation amounts for December-January-February **(a)** and June-July-August **(b)**. Red: GHG only; black: aerosol only. Lower panels: the sum of the changes in the GHG only and the aerosol only experiments **(c)**; and the changes in the Historic1 experiment with all natural and anthropogenic forcing agents included **(d)**. White patches indicate that changes are insignificant at the 95 % confidence level compared to internal variability in piControl. Units are mm d^{-1} in all panels.

**NorESM1-M –
climate response and
scenario projections**T. Iversen et al.




Fig. 6. Model simulated development from 1850 to 2100 in surface air temperature relative to the 1850–1899 average for Historic1 until 2005 followed by a range defined by the four RCP scenario projections. Blue: global data; red: the NH polar area north of 65° N. The diagram can be compared with Fig. 25 in Bentsen et al. (2012).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**NorESM1-M –
climate response and
scenario projections**

T. Iversen et al.




Fig. 7. Model calculated annual surface air temperature anomalies relative to the 1850–1899 average for RCP2.6 (upper panels) and RCP8.5 (lower panels), averaged globally (left panels) and over land areas (right panels). Black lines: NorESM1-M; blue and red lines: ensemble mean over 15 other models contributing to CMIP5, gray shading: one standard deviation on each side of the ensemble mean; blue and red shading: range defined by max and min values amongst the 15 models.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 8. Model simulated change in mean surface air temperature (K) (**a**, **b**) and percentage change in precipitation (**c**–**f**) from 1976–2005 to 2071–2100. Annual averages for RCP2.6 (**a**, **c**) and for RCP8.5 (**b**, **d**); RCP8.5 precipitation for (**e**) December-January-February and (**f**) June-July-August.

**NorESM1-M –
climate response and
scenario projections**

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NorESM1-M – climate response and scenario projections

T. Iversen et al.




Fig. 9. Decadal moving average annual evaporation minus precipitation **(a)** and evaporation **(b)** from the oceans, and the northern **(c)** and southern hemispheric **(d)** March and September sea ice extent during the NorESM1-M simulations for 1850 to 2300. Black in **(a)** and **(b)** and blue in **(c)** and **(d)**: the piControl, red: 1850–2005 Historic1, 2 and 3; dark green: RCP2.6 2005–2100; light green: RCP4.5 2005–2300; magenta: RCP 6.0 2005–2100; cyan: RCP8.5 2005–2100. Black curves in **(c)** and **(d)** are sea-ice extents estimated from observations (NSIDC, Fetterer et al., 2009).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NorESM1-M – climate response and scenario projections

T. Iversen et al.




Fig. 10. The top panel shows decadal moving averages of the annual max AMOC at 26.5° N, where grey is piControl, blue are Historic1, 2 and 3, green is RCP2.6, turquoise RCP4.5, violet RCP6.0, and red is RCP8.5. The bottom two panels show annual and zonal mean NorESM1-M simulated ocean temperature change for years 2071–2100 with RCP8.5 compared to piControl. Left: the Atlantic Ocean; right: global oceans.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NorESM1-M – climate response and scenario projections

T. Iversen et al.




Fig. 11. Diagnosis of NH extratropical storminess in NorESM1-M simulations of 1976–2005 (Historic1) by applying a band-pass frequency filter, emphasizing periods from 2.5 to 6 days, to the 500 hPa geopotential height (left). The middle panel shows bias error when compared to the ERA40 reanalysis data for 1976–2002 (Uppala et al., 2005), and the right panel shows the corresponding bias for the period 1979–2005 of the AMIP simulations with NorESM1-M run without coupling to the ocean model but with SST-fields prescribed from observations.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NorESM1-M – climate response and scenario projections

T. Iversen et al.




Fig. 12. NorESM1-M simulated change from 1976–2002 to 2071–2100 in NH extratropical storminess, diagnosed as described in Fig. 11, using the RCP8.5 projection scenario (left). Colours indicate significant changes on the 95 % confidence level. The right panel shows the standard deviation in the storminess amongst the three ensemble-members Historic1–3 for 1976–2005, revealing that the significant changes diagnosed in the left panel are considerably larger for the main maxima.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

NorESM1-M – climate response and scenario projections

T. Iversen et al.




Fig. 13. Northern hemispheric seasonal blocking statistics for the Tibaldi and Molteni (1990) (vTM) index diagnosed relative to the latitudes of the seasonally averaged position of the westlies (Pelly and Hoskins, 2002). Left panels are for the simulated NorESM1-M Historic1 for 1976–2005 compared to ERA40 statistics for 1976–2002, and right panels are projections for 2071–2100 with the RCP8.5 scenario compared to the period 1976–2005 (Historic1) with NorESM1-M. Dots signify longitudes where differences are significant at the 95 % confidence level.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 14. First and second rows show the 500-hPa-geopotential height associated with the 4 leading eof-vectors for detrended, monthly average anomalies of the 500 hPa geopotential height for December-March over the years 1976–2005, based on the three ensemble members (Historic1, 2, 3) simulated with NorESM1-M. Seasonal variations are removed by calculating anomalies relative to the 30-yr average for each month, while trends are removed by subtracting the 5-yr moving average. The third row shows the corresponding maps of the 2 leading eof-vectors calculated in the same way for the same 30 yr using the NCEP re-analysis data (Kalnay et al., 1996).

**NorESM1-M –
climate response and
scenario projections**

T. Iversen et al.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Fig. 15. The left column shows components of the non-detrended geopotential height anomalies for each 4-month season (DJFM) of the 3 historical ensemble members for 1976–2005 (black dashed lines and rings), and for the simulated climate projection for each 4-month season of the years 2071–2100 (denoted on the x-axes) using the RCP8.5 scenario with NorESM1-M (magenta dashed lines and rings). Thick lines are 10-yr forward moving averages. Right column shows similar components for the experiments with GHG-only (red), aerosol-only (blue), and natural forcing only (green) for 1976–2005.

**NorESM1-M –
climate response and
scenario projections**

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 16. Panel (a) shows time series of detrended monthly SST anomalies of the NINO3.4 region (5°S – 5°N ; 170°W – 120°W). The anomalies are found by subtracting the monthly means for the whole time series. Red (blue) colours indicate that anomalies are larger (smaller) than $+0.4\text{ K}$ (-0.4 K), see Trenberth (1997) for recommendations. Upper time series shows Hadley Centre Sea Ice and Sea Surface Temperature data set (HadISST; Rayner et al., 2003) for years 1900–2005; middle time series consist of NorESM1-M Historic1 for years 1900–2005 continued with NorESM1-M RCP8.5 for years 2006–2100; lower time series displays NorESM1-M piControl for years 750–950. (b) shows power spectra of the NINO3.4 index (the SST anomalies normalized with its standard deviation) using the multi-taper method of Ghil et al. (2002) with resolution $p = 4$ and number of tapers $t = 7$. Data sources are NorESM1-M piControl years 856–950 (black), and NorESM1-M RCP2.6 (blue), RCP4.5 (cyan), RCP6.0 (red), and RCP8.5 (magenta) for years 2006–2100.

**NorESM1-M –
climate response and
scenario projections**

T. Iversen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

