



This discussion paper is/has been under review for the journal Geoscientific Model Development (GMD). Please refer to the corresponding final paper in GMD if available.

GO5.0: The joint NERC-Met Office NEMO global ocean model for use in coupled and forced applications

A. Megann¹, D. Storkey², Y. Aksenov¹, S. Alderson¹, D. Calvert², T. Graham², P. Hyder², J. Siddorn², and B. Sinha¹

¹Marine Systems Modelling, National Oceanography Centre, Southampton, SO14 3ZH, UK

²Met Office, Hadley Centre, Exeter, Devon, EX1 3PB, UK

Received: 25 October 2013 – Accepted: 8 November 2013 – Published: 26 November 2013

Correspondence to: A. Megann (apm@noc.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We describe a new Global Ocean standard configuration (GO5.0) at eddy-permitting resolution, developed jointly between the National Oceanography Centre and the Met Office as part of the Joint Ocean Modelling Programme (JOMP). This programme is a working group of the UK's National Centre for Ocean Forecasting (NCOF) and part of the Joint Weather and Climate Research Programme (JWCRP). The configuration has been developed with the seamless approach to modelling in mind for ocean modelling across timescales and for a range of applications, from short-range ocean forecasting through seasonal forecasting to climate predictions as well as research use. The GO5.0 configuration has been coupled with sea-ice (GSI5.0), atmosphere (GA5.0) and land-surface (GL5.0) configurations to form a standard coupled global model (GC1). The GO5.0 model will become the basis for the ocean model component of the Forecasting Ocean Assimilation Model, which provides forced short-range forecasting services. The global coupled model (GC1) or future releases of it will be used in coupled short-range ocean forecasting, seasonal forecasting, decadal prediction and for climate prediction as part of the UK Earth System Model.

A 30 yr integration of GO5.0, run with CORE2 surface forcing from 1976 to 2005, is described, and the performance of the model in the final ten years of the integration is evaluated against observations and against a comparable integration of an earlier configuration, GO1. An additional set of 10 yr sensitivity studies, carried out to attribute changes in the model performance to individual changes in the model physics, is also analysed. GO5.0 is found to have substantially reduced subsurface drift above the depth of the thermocline relative to GO1, and also shows a significant improvement in the representation of the annual cycle of surface temperature and mixed-layer depth.

GMDD

6, 5747–5799, 2013

GO5.0

A. Megann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Coupled climate models developed at the UK Met Office have been at the forefront of international climate research and projections for the past fifteen years. HadCM3 (Gordon et al., 2000) was used in the Third and Fourth Assessment Reports (Houghton et al., 2001; Solomon et al., 2007) and is still widely used as a standard tool in climate research, while HadGEM1 (Johns et al., 2006), HadGEM2 (Collins et al., 2008) and HadGEM3 (Hewitt et al., 2011) have offered improvements in resolution, numerics and physics. All these models have an ocean on a horizontal grid of around 1° , although the HadGEM models have a refinement of the north-south grid scale close to the Equator down to $1/3^\circ$. In this paper we will refer to the model described by Hewitt et al. (2011) as HadGEM3; however, newer versions currently in development, with a higher resolution ocean, are also commonly referred to as HadGEM3.

Global ocean models are also used at the Met Office as part of seasonal and decadal forecasting systems (Arribas et al., 2011; Smith et al., 2007) and for ocean analysis and short-range forecasting (Storkey et al., 2010). At the Met Office and elsewhere there is increasing interest in using a seamless modelling system for use at all timescales from short range forecasting to climate prediction (Brown et al., 2012).

Increased horizontal resolution in the ocean has been shown to have several benefits for modelling climate. In the north Atlantic the improved path of the Gulf Stream and North Atlantic Drift reduces the magnitude of a large cold bias off Grand Banks seen in many low-resolution climate models (e.g. Gnanadesikan et al., 2007; Danabasoglu et al., 2010). Reducing this bias has been shown to improve the frequency of blocking in a climate model (Scaife et al., 2011). In the tropical Pacific ocean, eddy permitting resolution in HiGEM has been shown to help reduce the equatorial cold tongue bias (Shaffrey et al., 2009; Roberts et al., 2009) and the double intertropical convergence zone (ITCZ) bias, and also to more realistically simulate the westward extent of El Niño. Furthermore, teleconnections to the North Pacific Ocean associated with ENSO were also improved as a result of increased ocean resolution (Dawson et al.,

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2012). Eddy permitting models have an order of magnitude more eddy kinetic energy (EKE) than low-resolution models (Delworth, 2012) and the NEMO model on the 1/4° ORCA025 grid has been shown to simulate 81 % of observed sea level variability on interannual timescales (Penduff et al., 2010). The high resolution also allows significant
 5 improvement in simulating oceanic circulation in the Arctic Ocean as well as exchanges between the Arctic and the North Atlantic (e.g., Aksenov et al., 2011).

Ocean models run on horizontal grids fine enough to resolve eddies in the Southern Ocean show “eddy saturation”, where increased meridional transport of momentum away from the Antarctic Circumpolar Current (ACC) by the eddy field in response
 10 to increases in wind stress mean the circumpolar transport is relatively insensitive to changes in the wind forcing (Tansley and Marshall, 2001; Hallberg and Gnanadesikan, 2006). This is not observed in lower resolution models where the eddy transports are parameterised by diffusive schemes. A similar insensitivity of the global overturning circulation to the Southern Ocean wind forcing (“eddy compensation”) is also seen in
 15 eddy-resolving models (Viebahn and Eden, 2010, Farneti et al., 2010). This implies that important dynamical adjustment processes are not present in models that are unable to represent the Southern Ocean eddy field.

Here we describe a new Global Ocean standard configuration (GO5.0) at eddy-permitting resolution, developed jointly between the National Oceanography Centre and the Met Office as part of the Joint Ocean Modelling Programme (JOMP), a working
 20 group of the UK’s National Centre for Ocean Forecasting (NCOF) and part of the Joint Weather and Climate Research Programme (JWCRP). The configuration has been developed with the seamless approach to modelling in mind and is therefore intended to be used as the basis for ocean modelling across timescales and for a range of applications, from short-range ocean forecasting, through seasonal forecasting, to climate
 25 predictions as well as research use. The configuration has been developed for use throughout the UK academic and operational modelling communities. It has been coupled with the sea-ice (GSI5.0), the atmosphere (GA5.0) and the land-surface (GL5.0) configurations to form a standard coupled global model (GC1). The GO5.0 model will

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

become the basis for the ocean model component of the Forecasting Ocean Assimilation Model (FOAM, Storkey et al., 2010), which presently provides forced short-range forecasting services to MyOcean (<http://myocean.eu>) and other users. The global coupled model (GC1) or future releases of it will be used in coupled short-range ocean forecasting (as future versions of FOAM evolve into coupled systems), for seasonal forecasting as part of the GloSea4 system (Arribas et al., 2011), for decadal prediction as part of the DePreSys system (Smith et al., 2007) and for climate prediction as part of the UK Earth System Model.

We use the term “standard configuration” to denote all the items required to run the model, i.e. model code, input parameters and datasets, and compilation keys, and these are summarised in the Appendices.

The main aim of this paper is to introduce the ocean model constituting GO5.0, and to evaluate its performance in ocean-only configuration, according to a set of first-order metrics. We will also compare the performance of GO5.0 with the previous global ocean configuration, which we denote GO1, and attribute the salient differences between the two model implementations to specific changes in model physics and parameter sets. Section 2 describes the ocean and ice models and the surface forcing fields. Section 3 summarises the main physics choices, and in Sect. 4 the experimental design is described. In Sect. 5 we present the results of the analysis: firstly the GO5.0 configuration is validated against observations; then GO5.0 is compared with the previous global model GO1; and the main improvements identified in GO5.0 are attributed to specific physics choices. Finally in Sect. 6 we summarise the results and discuss upgrades to the model currently under development.

2 Model description

GO5.0 is based on version 3.4 (v3.4) of NEMO (Nucleus for European Models of the Ocean) (Madec, 2008), and is closely related to the global DRAKKAR ORCA025 configuration (Barnier et al., 2006) sharing many of the same dynamics and physics

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

choices. The horizontal grid, known as ORCA025, has $1/4^\circ$ resolution (1442×1021 grid points) at global scale decreasing poleward (an isotropic Mercator grid in the Southern Hemisphere, matched to a quasi-isotropic bipolar grid in the Northern Hemisphere with poles at 107° W and 73° E). The effective resolution is approximately 27.75 km at the equator, but increases with latitude to be for example 13.8 km at 60° S or 60° N. The model has 75 vertical levels where the level thickness is a double tanh function of depth such that the grid spacing increases from 1 m near the surface to 200 m at 6000 m (I. Culverwell, personal communication, 2009). This level set was chosen to provide high resolution near the surface for short to mid range forecasting purposes while retaining reasonable resolution at mid-depths for long term climate studies.

The model bathymetry is based on the ETOPO1 dataset (Amante and Eakins, 2009) with additional data in coastal regions from GEBCO (IOC, 2008). This is a change from the GO1 configuration, which used the DRAKKAR G70 bathymetry based on ETOPO2 with corrections from satellite-based bathymetry and other sources (Remy et al., 2003). Bottom topography is represented as partial steps (Barnier et al., 2006).

The model uses a linear free surface and an energy and enstrophy conserving momentum advection scheme. The horizontal viscosity is bilaplacian with a value of $1.5 \times 10^{11} \text{ m}^2 \text{ s}^{-1}$ at the equator, reducing polewards. Tracer advection uses a Total Variance Dissipation (TVD) scheme (Zalesak, 1979). Lateral tracer mixing is along isoneutral surfaces with a coefficient of $300 \text{ m}^2 \text{ s}^{-1}$. No parameterisation of eddy mixing is used.

With regard to diapycnal mixing processes, the vertical mixing of tracers and momentum is parameterised using a modified version of the Gaspar et al. (1990) Turbulent Kinetic Energy (TKE) scheme (Madec et al., 1998). Unresolved vertical mixing processes are represented by a background vertical eddy diffusivity of $1.2 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$, which decreases linearly from $\pm 15^\circ$ latitude to a value of $1.2 \times 10^{-6} \text{ m}^2 \text{ s}^{-1}$ at $\pm 5^\circ$ latitude (Gregg et al., 2003) and a globally constant background viscosity of $1.2 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$. A parameterisation of double diffusive mixing is included at GO5.0 (Merryfield et al., 1999).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Bottom friction is quadratic with an increased coefficient in the Indonesian Throughflow, Denmark Strait and Bab al Mandab regions. An advective and diffusive bottom boundary layer scheme is included (Beckmann and Doscher, 1997). The tidal mixing parameterisation of Simmons et al. (2004) is included with a special formulation for the Indonesian Throughflow (Koch-Larrouy et al., 2008). At GO5.0 a climatological geothermal heat flux is added as a bottom boundary condition; this was not used at GO1.

GO5.0 is driven over the period 1976–2005 by the CORE2 surface forcing data set (Large and Yeager, 2004). CORE2 supplies monthly precipitation and daily downward shortwave and longwave radiation which are used to force the model directly, and 6 hourly 10 m wind, 2 m air humidity and 2 m air temperature which are used to compute turbulent air/sea and air/sea ice fluxes during model integration using the bulk formulae proposed by Large and Yeager (2004). The source data for precipitation and radiative fluxes are only available from 1979 and 1984 onward, respectively. Prior to these dates the respective climatologies are used. Climatological monthly runoffs derived from the Dai and Trenberth, 2002 climatology are applied along the land mask (Bourdalle-Badie and Treguier, 2006). No diurnal cycle is imposed in the radiative forcing.

Initial conditions for temperature and salinity are obtained from an average of years 2004–2008 of the EN3 monthly objective analysis (Ingleby and Huddleston, 2007) and the model is started from a state of rest. To avoid unacceptable drifts in salinity and an excessive spin-down of the overturning circulation, the sea surface salinity (SSS) is restored toward monthly mean climatological values. The relaxation time scale is 180 days for the open ocean and 36 days under sea ice. Model outputs are archived as successive 5 day means throughout the whole integration and post-processed to monthly means. More details about the model configuration may be found in Storkey et al. (2010), Barnier et al. (2006) and Penduff et al. (2007).

The sea ice component is the latest public release of the Los Alamos National Laboratory sea ice model CICE version 4.1 (v4.1, Hunke and Lipscomb, 2010). The model includes Elastic–Viscous–Plastic ice dynamics (Hunke and Dukowicz, 1997),

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

energy-conserving thermodynamics (Bitz and Lipscomb, 1999) and multi-category ice thickness (Bitz et al., 2001). The setup of CICE is the same as in the lower-resolution version of HadGEM3 described by Hewitt et al. (2011) with five sea ice thickness categories. Both GO5.0 and HadGEM3 use the zero-layer Semtner thermodynamics scheme (Semtner, 1976). We also note that in both the GO5.0 and HadGEM3 configurations the sea ice model is not on the same grid as the ocean (sea ice is on the Arakawa B-grid and ocean is on the Arakawa C-grid; Arakawa, 1966) and an interpolation routine is used to couple these model components. As in HadGEM3, the ice and ocean components are combined into a single executable, so there is no need for a coupler.

We shall also discuss the preceding version of the Met Office ocean model, GO1. This was based on NEMO version 3.2 (v3.2) and CICE v4.1, and was implemented on the same grid as GO5.0, with the same surface forcing.

The ocean and ice code are managed using the Subversion code-management software, allowing unique identification of the respective code bases using a code version number. Ocean and Ice model code version numbers, compilation keys and namelists are listed in the Appendices.

The model was run on the MonSOON supercomputer, jointly owned by NERC and the Met Office. The ocean and ice models were distributed over 480 cores with the MPI communications harness, and an acceptable throughput of one model year in six hours was achieved.

3 Summary of main physics choices

The main physics changes between GO1 and GO5.0 are changes to the vertical mixing parameters based on the work of Calvert and Siddorn (2013). Vertical mixing in the model is achieved using a turbulent closure scheme with an algebraic mixing length (Gaspar et al., 1990; Madec et al., 1998). In addition to the basic turbulent kinetic energy (TKE) equation, a number of additional parameterisations are included

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

to represent processes such as surface wave breaking. One such term represents the enhanced mixing due to breaking of near-inertial waves as an additional source of TKE exponentially decaying from the surface. Based on the results of sensitivity experiments by Calvert and Siddorn (2013) the vertical length scale for this TKE source term has been reduced from 30 m to 10 m at mid to high latitudes and increased from 0.5 m to 10 m in the tropics. One effect of this is to reduce summer time mixed layer depths at midlatitudes.

Convection in the model is parameterised as an enhanced vertical diffusivity of $10 \text{ m}^2 \text{ s}^{-1}$ for momentum and tracer fields where the water column is unstable. At NEMO v3.2 this enhanced vertical diffusivity was erroneously used in the prognostic equation for the TKE, instead of the vertical diffusivity calculated by the TKE scheme. This was shown to result in a deep bias in wintertime mixed layer depths owing to the non-conservative increase in the calculated TKE. This has since been addressed at NEMO v3.4 and therefore constitutes another difference between GO5.0 and GO1. A further change between v3.2 and v3.4 is the use of the in situ surface salinity, rather than a constant reference salinity of 35 psu used in the former, in the conversion from freshwater fluxes to salt fluxes.

Other changes between GO1 and GO5.0 are: changes to other vertical mixing parameters between GO1 and GO5.0 as noted in Table 1, the inclusion of a double diffusive mixing parameterisation at GO5.0, and the inclusion of a climatological geothermal heating parameterisation at GO5.0.

4 Experimental design

The GO5.0 configuration can be viewed as a set of incremental changes in the model physics relative to the GO1 configuration. In order to evaluate the GO5.0 configuration and to understand the model improvements over GO1, a series of forced ocean-sea ice integrations was performed to assess the effects of each individual change.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A thirty-year integration of GO5.0 was carried out with the final set of modifications and parameter values, from the initial state described above, derived from EN3 climatology. This was compared with the reference integration, from the same initial state and of the same length, of the pre-existing GO1 model based on NEMO v3.2. To estimate the effect of the code change alone, a further thirty-year integration of NEMO 3.4 was made with initial state and all parameters and physics choices identical, or as close as possible, to those of GO1. We compare annual and seasonal means from each of these three integrations and also with observations in the form of the EN3 climatology for subsurface temperature and salinity (Ingleby and Huddleston, 2007), the HadSST3 surface temperature climatology (Kennedy et al., 2011), satellite-derived sea-ice extent (Cavalieri, 1996, updated 2013), the Pan-Arctic Ice Ocean Modeling and Assimilation System (PIOMAS) reanalysis for Northern Hemisphere sea ice volume (Zhang et al., 2003), and measured transports through key straits from a variety of observational studies.

An additional set of ten-year simulations was made to attribute changes between GO1 and GO5.0 to individual changes in configuration. These are summarised as follows:

- The bathymetry was upgraded from the original Drakkar ORCA025 dataset;
- The background vertical diffusivity and viscosity were increased from 1.0×10^{-5} to $1.2 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ and from 1.0×10^{-4} to $1.2 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$, respectively;
- Changes were made to the TKE scheme parameters *rn_ebb*, *rn_mx10* and *nn_htau*;
- Geothermal heat flux and double diffusion of tracers were added;
- Schemes for a bottom boundary layer and lakes were added
- The ice model (CICE) was modified to include a salinity-dependent freezing point. The thermal conductivity of the ice was changed from 2.00 to $2.63 \text{ W m}^{-1} \text{ K}^{-1}$ and

the fixed ice salinity was changed from 4.0 to 8.0 psu, following J. Rae et al. (personal communication, 2013).

The attribution study will compare the above experiments with one another, as well as with the v3.2 model GO1 and the original v3.4 integration with the GO1 parameter set.

We admit that the strategy of adjusting parameters according to individual sensitivity studies may not be the optimal method for finding the most appropriate parameter set, since the parameters and physics choices may interact nonlinearly, but resources were insufficient for a systematic investigation of parameter space such as that carried out with HadCM3 by Williamson et al. (2013).

Table 2 summarises the integrations carried out, including the values of the principal parameter changes at each step. We note that the pair of runs comparing NEMO 3.2 and 3.4 (namely GO1 and N3.4) differ further in one minor respect. The v3.4 parameter *rn_mx10* was erroneously set to 0.001 in the latter experiment to match the value of the parameter *rn_lmin* in v3.2, but the equivalent parameter in v3.2 is in fact *rn_lmin0*, which was set to 0.01. An additional ten-year integration similar to N3.4 was performed, with a value of 0.01: the consequent surface changes were not considered to be significant, with the two simulations being qualitatively the same, with mean surface temperature differences in years 6–10 less than 0.05 °C everywhere.

5 Results

5.1 Validation of GO5.0 against observations

5.1.1 Surface biases and mixed layer depth

Figure 1 shows the surface temperature (SST) and salinity (SSS) errors in years 21–30 of the GO5.0 model, relative to the mean of the Reynolds et al. (2002) and EN3 (Ingleby and Huddleston, 2007) respective monthly climatology over the same period.

There is a cool bias over most of Northern Hemisphere, with the exception of the

GMDD

6, 5747–5799, 2013

GO5.0

A. Megann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

subpolar gyres in the North Atlantic and North Pacific. GO5.0 is too fresh in most of the Atlantic, except in the subpolar gyre, where the salty bias of 0.5–1.0 psu is co-located with the warm bias mentioned above. Generally the areas where there is a surface warm bias (especially in the southern Ocean and the Pacific) correspond to a positive surface salinity error: these may result from forcing errors, but are not inconsistent with an excessive evaporation from surface waters with a warm bias. The exception is in the Arctic, where there is a positive surface salinity error of up to 2 psu, due to the excessive autumn sea ice formation on the Siberian shelves and in the Beaufort Sea (Fig. 1); the reason for this error is unclear, but is most likely to be related to the air temperature and radiative biases in the atmospheric forcing (Barnier et al., 2006). The sea ice biases are discussed further in the Sect. 5.1.4.

Figure 2 shows the annual minimum and maximum mixed-layer depth calculated from monthly data for years 1996–2005, corresponding to the shallowest depth of the mixed layer in the local hemispheric summer and the deepest mixed layer in the local hemispheric winter, alongside the same quantity from the de Boyer Montégut et al. (2004) climatology. The GO5.0 model realistically reproduces the spatial patterns of both summer and winter surface mixing realistically: in particular, the regions of wintertime dense water formation in the North Atlantic and Nordic Seas correspond quite closely to those in the observations, as do the near-zonal bands of deep turbulent mixing in the Southern Ocean (Fig. 2c and d). There is a consistent bias, however, to an unrealistically shallow summer mixed layer over the whole ocean, with maximum values of 30–50 m in the tropics and Southern Ocean in the model, contrasting with a range of 50–70 m in the same regions in the climatology (Fig. 2a and b). This is consistent with the warm surface bias in the same regions seen in Fig. 1. Also, the winter mixing in the dense water formation regions in the North Atlantic and in the Weddell Sea is much deeper than in the climatology, reaching to over 1000 m in many instances. The patch of very deep mixing extending from the Weddell Sea eastwards to 50° E is also seen in HadGEM1 and HiGEM, but not in the observations, where instead deep mixing is strongest in the coastal regions of the Weddell Sea. In GO5.0 this feature develops

after year 20 of the integration, but it does not occur in GO1: it seems to be associated with a gradual modification of the watermasses in the region and the development of the spurious polynya visible in Fig. 6c (discussed in Sect. 5.1.4), which together precondition for the unrealistically deep mixing, but the exact mechanisms are yet unclear.

The surface biases of the model when forced by prescribed surface boundary conditions are to a large degree constrained by the forcing fields, but the subsurface drifts are a stronger test of the model, revealing discrepancies in diapycnal mixing and advection pathways. Figure 3 shows the zonal mean temperature and salinity anomalies in GO5.0 averaged from 1996 to 2005, with reference to the EN3 climatology. The largest biases are in the top 700 m of the water column. Note the cold subsurface bias ($\sim 2^\circ\text{C}$) around Antarctica, the warm salty bias ($\sim 1.5^\circ\text{C}$ and 0.25 psu) between 45 and 60°S , a warm bias in the tropics of up to 2.5°C down to about 200 m, cold, fresh biases in the main thermocline (45°S – 45°N , with maximum discrepancies of 1°C and 0.5 psu), and a warm salty bias in the Northern Hemisphere subpolar gyre regions ($\sim 1^\circ\text{C}$, 0.25 psu).

5.1.2 Atlantic meridional overturning

Figure 4a shows a time series of the North Atlantic overturning strength at 26°N for the full 30 yr integration of GO5.0, alongside that in GO1. Also shown is the potential density averaged over the upper 200 m in the central Labrador Sea (55 – 58°N , 48 – 50°W). Figure 4b illustrates the meridional overturning streamfunction in years 1996 to 2005 in both models. In both GO1 and GO5.0 the overturning circulation reaches a maximum in the second decade of the integration, reducing by 2–3 Sv by year 30; the run length is however not sufficient to determine whether the circulation has settled at that stage. In both runs the decadal scale variability in the MOC is clearly correlated with the Labrador Sea density, which also increases over the first decade and decreases later in the second decade. The overturning strength at 26°N in the final decade is between 21 and 22 Sv, which is at the higher end of the range of 18.5 ± 1 Sv observed between 2004 and 2008 by the RAPID WATCH/MOCHA array (McCarthy et al., 2012). Interestingly, the downward trend we see in the last decade of the model runs (~ 2 – 3 Sv/decade) is

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

similar to that recently reported from the RAPID array (Smeed et al., 2013). The modelled annual means for two years overlapping the observations, namely 2004 (19 Sv) and 2005 (20 Sv) match well with the observations (17.8 Sv and 20.1 Sv respectively), and the strength of the modelled AMOC over latter decade of the run is entirely plausible, particularly since recent studies indicate that a substantial fraction of the variability in the strength of the AMOC originates from surface forcing (Roberts et al., 2013; Blaker et al., 2013). We cannot expect the model to simulate the measured AMOC perfectly, since up to 30 % of the AMOC variability is inherently unpredictable, arising as a consequence of baroclinic wave field and mesoscale eddy field (Hirschi et al., 2013; Thomas and Zhai, 2013). We note that the modelled annual means quoted here are January–December, whilst the observational array figures are April–March.

Figure 4b also shows that the depth of the North Atlantic Deep Water (NADW) return flow is too shallow. At 26° N the depth of the NADW return flow (usually defined as the depth of the zero contour in the streamfunction) is around 3500 m for most of the model run compared to deeper than 4000 m in the RAPID array data. This is a common bias in many ocean GCMs using depth coordinates, and is usually attributed to spurious mixing of overflow waters as they descend from passages in the Greenland–Iceland–Scotland ridges to the deep ocean (Saunders et al., 2008; Danabasoglu, 2010). It is worth noting that substantial variation in the depth profile can arise from the method used to compute the overturning. Computing the overturning from a model using the RAPID array methodology and assuming a geostrophic reference depth of 4740 m can yield a transport profile much more similar to the observations at 26° N than integrating the model velocities (Roberts et al., 2013).

The increase in the AMOC over the first decade of the model run is a phenomenon often seen in ocean GCMs using mixed surface boundary conditions in which the high latitude oceans become overly sensitive to salinity perturbations (Rahmstorf and Willebrand, 1995; Lohmann et al., 1996; Greatbach and Peterson, 1996; Griffies et al., 2009; Yeager and Jochum, 2009). In the GO5.0 model run an initial error in the path of the North Atlantic Current (NAC) causes warm, salty water to be advected into the subpolar

gyre where it joins the Greenland current and enters the Labrador Sea. The prescribed surface air temperature causes excessive surface heat loss in the Labrador Sea, increasing the density of the surface waters and leading to excessive deep water formation in this region. The increase in the MOC causes more warm, salty water to be advected into the subpolar gyre in a positive feedback. Yeager and Jochum (2009) show that stronger sea surface salinity restoring can reduce this feedback mechanism by reducing the surface salinity in the Labrador Sea. This improvement, however, comes at the expense of realistic interannual variability in the global climate.

5.1.3 Critical sill and strait transports

Table 3 lists the volume transports through the major straits and across critical sills, averaged over the last 10 yr of the 30 yr integrations of GO1 and GO5.0, together with recent observed estimates and their sources. Overall, the models simulate these transports acceptably: in particular, the Drake Passage throughflow is much closer to observations than those in the lower-resolution coupled models HadCM3 and HadGEM1, both of which at ~ 200 Sv (Johns et al., 2006) are unrealistically strong. HadGEM2 (Collins et al., 2008) gave a comparable simulated Drake Passage transport of ~ 140 Sv (Meiers et al., 2012). In contrast to these aforementioned coupled models, the $1/4^\circ$ resolution of the ORCA025 grid allows the present model to at least approach an explicit resolution of the narrower passages, including the Bering and Gibraltar Straits, and it can be seen that both GO1 and GO5.0 have transports through each of these of well within a factor of two of the observed values.

The Indonesian Throughflow is too strong in both GO1 and GO5.0, which may be due to insufficient enhancement of the tidal mixing in this region (Koch-Larrouy et al., 2008).

Comparing the model-derived and observation-based estimates of the Arctic–Atlantic exchanges across the Greenland–Scotland Ridge, through Fram and Davis Straits, and through the Barents Sea shows that in both 30 yr model runs the volume transports are within 10–20 % of the long-term mean observed values and within the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

range of the observational uncertainties, except for the Denmark Strait overflow where the model estimates are 45 % (GO1) and 33 % (GO5.0) higher than the observational estimate (Table 3). Although the simulated net outflow from the Arctic Ocean, of 4.8 Sv in GO1 and 4.6 Sv in GO5.0, is very close to the observed value of 4.6 Sv, the model shows a different partitioning of the exports west and east of Greenland: the simulated flow through the Canadian Archipelago is 35–45 % larger than the export through Fram Strait, which is opposite to the observations. The bias is stronger in summer than in winter and is due to excessive Ekman convergence in the Beaufort Sea; this in its turn is caused by the summer sea ice extent being too low (see next section). The simulated Pacific inflow in Bering Strait is higher than in the observations, even considering the recent update in the latter estimate (Woodgate et al., 2012). The excessive Pacific inflow is caused by the gradient in the sea surface height between the Northern Pacific and the Chukchi Sea being too steep. The stronger Pacific inflow brings extra heat into the Arctic Ocean, which may contribute to the excessive sea ice melting (see next section).

Overall, both runs, GO1 and GO5.0, present more vigorous northward flow of the Atlantic water than is observed (“Total Greenland–Scotland inflow” in Table 3 is a proxy for this) and stronger than observed return overflows across the Greenland–Scotland Ridge: the combined overflows through Denmark Strait and through the openings between Iceland and the Faeroes and between the Faeroes and Scotland are 6.3 Sv from the data, 9.3 Sv in GO1 and 8.3 Sv in GO5.0. This is also evident in the stronger simulated AMOC compared to observations.

It should be noted that the observational estimates of the exchange transports into and out of the Arctic should be treated with caution. First, in all straits, except for the moorings in Bering Strait, the hydrographic section in Fram Strait and the one in the Barents Sea between Norway and the Bjørnøya (Barents Sea Opening), uninterrupted records from current meter moorings are no longer than two years. This aliases interannual variations and introduces large uncertainties in the observational transports estimates. Secondly, the instruments were not positioned in the top 50 m and

on the shallow shelves, in order to prevent the moorings being damaged by sea ice keels. Lastly, the distances between the moorings were too great to resolve mesoscale variability of the flows and in Bering Strait the transports were derived from velocity measurements obtained from three separate moorings (e.g., Woodgate et al., 2012).

All this introduces spatial aliasing in the interpolating procedures and uncertainties in the transports. For detailed discussion of uncertainties in observed transports, please refer to e.g. Curry et al., 2011 and Olsen et al., 2008. It also should be noted that, while the model standard deviations in the table represent variability of the transports on synoptic to interannual timescales, the standard deviations of the observational estimates include uncertainty inherent in the estimation methods as well as the variability of the transports, thus rigorous comparison of the variability in the model and data requires additional analysis, not presented here.

5.1.4 Sea ice

Timeseries of the sea ice extent and ice concentration in the Northern and Southern Hemispheres are compared with products from passive microwave satellites SSMR/I and AVHRR (Cavalieri, 1996, updated 2013). In the Northern Hemisphere the simulated annual mean of $11.2 \times 10^6 \text{ km}^2$ and the amplitude of the seasonal cycle of $7 \times 10^6 \text{ km}^2$ are in good agreement with the data ($12.4 \times 10^6 \text{ km}^2$ and $5.8 \times 10^6 \text{ km}^2$, respectively), suggesting good model skill in simulating sea ice extent (Fig. 5a), although the model underestimates summer sea ice extent. The simulated and observed inter-annual trends also agree. Figure 5b compares the modelled Arctic sea ice volumes with these derived from the PIOMAS reanalysis (Zhang et al., 2003). Simulated sea ice volumes are about 60 % of those observed through the annual cycle, with winter (DJF) biases of around 30 % and in summer (JJA) of around 50 %. Despite this bias, the multidecadal trends in the modelled and observed sea ice extents are comparable, showing sea ice extent decline at a rate of $-44 \times 10^3 \text{ km}^2$ per year and $-45 \times 10^3 \text{ km}^2$ per year, respectively. In the Southern Hemisphere the modelled sea ice extent is again in good agreement with observations (Fig. 5c), but with a moderate negative summer

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

bias. At present no published sea ice volume timeseries are available for Antarctica, rendering formal validation of the model skills in simulating sea ice volumes in the Southern Hemisphere impossible. However, comparing simulated sea ice thicknesses around Antarctica for 1996–2005 with the Antarctic Sea Ice Processes and Climate (ASPeCt) data (Worby et al., 2008) for the same period, we conclude that the simulations underestimate long-term mean annual sea ice thickness by about 22 % (0.76 m in the model and 0.89 m for simulations and data respectively). The annual cycle in the model is in good agreement with the observations, with the maximum ice thickness (1.06 m in the model and 1.02 m in the observations) occurring in the austral Summer (DJF) and minimum ice thickness (0.58 m in the model and 0.60 m in the observations) in the austral Winter (JJA). The simulated sea ice extent trend in the Southern Hemisphere is negative and around $-58 \times 10^3 \text{ km}^2 \text{ yr}^{-1}$, in contrast to the positive trend of $13 \times 10^3 \text{ km}^2 \text{ yr}^{-1}$ in the observations. The negative trend in Antarctic sea ice extent is a common feature of global ocean models, and is attributed by Holland and Kwok (2012) to biases in the surface winds around Antarctica.

Comparison between the simulated sea ice concentration fields and those from the HadISST observational dataset (Rayner et al., 2002) show that the simulated winter sea ice distribution in both hemispheres is realistic (Fig. 6a–d), although we note that there is a tongue of reduced ice cover extending eastward from the central Weddell Sea, which has also been seen in HadGEM1 and the higher-resolution HiGEM (Shafrey et al., 2009), and which corresponds to the very deep winter mixing described in Sect. 5.1.1. The summer sea ice concentration in the model is lower than in the data (Fig. 6e–g and i). In the Arctic Ocean this is likely to be caused by the negative bias in the sea ice thickness, which in turn results in lower ice strength, faster ice drift toward the Canadian Arctic Archipelago and thus increased divergence of sea ice in the Central Arctic Ocean. This, combined with the increased sea ice melting in summer due to exposure of the ocean surface to the atmospheric heat, could sustain the lower thicknesses in the Arctic throughout the year. In the present forced simulations without atmosphere-ocean feedbacks, the summer sea ice bias primarily affects polar regions

is also interannual variability in the globally averaged surface temperature and salinity in the upper 200 m, which is probably a signature of ENSO.

It is interesting to relate the drifts in GO1 and GO5.0 to those over the first thirty years of HadGEM1 (Johns et al., 2006) and in CHIME and HadCM3 (Megann et al., 2010). All these except for CHIME (which uses a hybrid isopycnic-coordinate ocean, in contrast to the depth-coordinate ocean model in the other three) have a pronounced freshening in the upper ocean that steadily penetrates into the interior, and this is likely to be a consequence of the numerical diapycnal mixing typical of this model type (Griffies et al., 2000). HadCM3 and HadGEM1 (which shared an ocean model, albeit on a slightly different grid) similarly had a negative surface temperature error over most of the ocean, offset in HadCM3 by a warm bias in the Southern Ocean, while CHIME had a warm surface error, consistent with a reduced drawdown of heat by numerical mixing.

5.2.2 Seasonal cycle of surface temperature and mixed layer depth

Figure 8 shows the mean biases of the sea surface temperature in GO1 and GO5.0 with respect to the Reynolds et al. climatology in the boreal winter and boreal summer seasons, defined as December/January/February and June/July/August periods respectively. It is clear that both configurations have substantial biases in the time-averaged surface fields, and as with the 10 yr mean fields discussed in Sect. 5.1.1, in many regions these biases are very similar: for example, the tropics and Southern Ocean are generally too warm in both configurations, while the northern high latitudes are generally too cold, and there is a warm error in the subpolar North Atlantic with maximum values of 3–4 °C in the boreal winter. There are regions where the seasonal biases in GO1 are smaller than in GO5.0: for example, the cold boreal winter error in the subtropical North Atlantic is larger in GO5.0 south of the separated Gulf Stream (Fig. 8a and b), and in the Southern Ocean there is a substantial coherent warm error in GO5.0 in the austral summer that is not present to the same extent in GO1. Overall, however, there are large scale reductions in seasonal bias, particularly in the northern summer (JJA) season: the cold errors in the North Atlantic and North Pacific are

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



substantially reduced in GO5.0, as are the warm biases in the tropics and the Southern Ocean.

To illustrate the latitude dependence of the large-scale seasonal biases in GO1 and GO5.0, Fig. 9 shows latitude-time plots of the zonally averaged surface temperature bias (referred to the Reynolds et al. climatology) and mixed layer depth error (referred to the de Boyet Montégut et al., 2004 data) in GO1 and GO5.0. This shows more clearly that the boreal summer warm bias in the tropics is reduced in GO5.0, as is also the large summer cold bias in the northern subtropics. As we have already noted, GO5.0 shows systematic biases in both the minimum and maximum mixed-layer depth (Fig. 2): specifically, in both hemispheres winter mixed layers are generally too deep, while summer mixed layers are generally too shallow. The main difference between GO1 and GO5.0 is that mixed layer depths are generally shallower in GO5.0, leading to reduced stratification and hence the warmer summer surface temperatures, especially in the Southern Ocean, seen in Figs. 8b and 9a. The winter mixed layer depth biases, by contrast, are generally reduced in GO5.0.

5.3 Attribution of changes

In this section we refer to the experimental design described in Sect. 4, where a series of shorter (10 yr) integrations are made. The model code is first upgraded from NEMO v3.2 to v3.4, then other changes are then progressively made within v3.4, to attribute the most significant changes in model fields to specific changes in the model physics. These changes are summarised in Table 2. We compare the mean fields in the final five years (1981–1985) of each ten-year integration; the main comparison will be of the surface fields, but the global subsurface biases down to 700 m will also be compared. Our empirical criterion for significance is that the effect of any physics change should be locally large compared with the amplitude of the mesoscale “eddy” signal.

The code changes from NEMO version 3.2 to 3.4 have two main physics components: firstly in the correction to the treatment of convective mixing in the TKE scheme described in Sect. 3, and secondly in the change in the method of calculation of the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



effective salt fluxes applied to the surface level from the surface freshwater fluxes (as described in Sect. 3). This latter modification is expected to lead to changes in the mean surface salinity, since the surface freshwater flux in v3.4 will depend on the local surface salinity: for a given freshwater flux, we would expect a larger effect on the surface salinity where the surface is saltier than in fresher regions. Figure 10 shows that the code upgrade from v3.2 to v3.4 clearly has significant effects on the surface fields: there are basin-scale changes over almost the whole ocean, with warming of 0.1–0.2 °C over the Arctic and the subtropical gyres, but cooling by a similar magnitude on the equator and coastal upwelling regions, in the Southern Ocean and in the North Atlantic subpolar gyre. The surface salinity changes are also predominantly in zonal bands, with the largest increases of 0.2–0.4 psu between 15° S and 30° S and between 15° N and 30° N in the Atlantic and eastern Pacific and a surface freshening over much of the Southern Ocean. The largest of the aforementioned positive salinity changes (in the south Atlantic and south Pacific) are consistent with the change in reference salinity, since these are regions of high salinity and net evaporation. The code change overall, however, has little effect on the RMS surface errors of the model: the RMS SST error reduces from 0.665 to 0.657 °C, while the RMS surface salinity error barely changes from 0.828 to 0.825 psu. There are, however, major subsurface effects resulting from the code upgrade, particularly from the correction to the treatment of convective mixing in the TKE scheme: comparing the temperature changes in the upper 700 m with the mean isopycnal depths (Fig. 11) shows that the upgrade removes much of the warm bias in the thermocline region between 50° S and 60° N, via a mean cooling of up to 1 °C in the depth range from 50 to 250 m over these latitudes. Additionally, the drastic reduction in winter MLD biases between v3.2 and v3.4 observed in Fig. 9 can be directly attributed to the convective mixing correction. The crescent shape of the temperature bias with respect to the observations (and of the difference between v3.4 and v3.4) in Fig. 11 reflects the deepening of the thermocline with increasing latitude.

Upgrading the bathymetry (not shown) leads to small changes in the temperature and salinity in the Arctic, which overall cools by 0.05 °C or less and freshens by around

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

0.05 psu: this is likely to be a consequence of minor modifications to the North Atlantic sill topography. There are large-scale southwards displacements of the path of the topographically-steered ACC, north of the Kerguelen Plateau and north of the Pacific Antarctic Ridge at 140–150° W, along with a depression of the surface elevation in the Southern Ocean by 3–5 cm, which may be associated with alterations in the path and strength of the northward-flowing Antarctic Bottom Water.

Increasing the background vertical diffusivity and viscosity by 20 % (not shown) has a small effect on the surface fields, relative to the other parameter changes. There is a general surface freshening in the Arctic by 0.02–0.04 psu, and a hint of warming north of the ACC, but elsewhere any signal is small compared with the mesoscale noise. It is possible that the explicit representation of mixing processes dominates the background term, while it is also likely that over much of the ocean the numerical mixing is at least as large as that associated with the $1.2 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ explicit background diffusivity, as discussed in Griffies et al., (2000) and Lee et al. (2002).

The changes to the TKE scheme parameters lead to a consistent surface warming of between 0.1 and 0.5 °C north of 30° N and south of 30° S (Fig. 12), while there is a small cooling of around 0.05 °C in the tropics. The pattern of the associated salinity changes is more complex, with freshening of up to 0.2 psu in the Arctic, in the subpolar North Pacific, and to a lesser extent in the tropics and along the path of the ACC; and an increase in salinity in the subtropical zones and, interestingly, in the regions dominated by the Amazon and Congo river plumes. The subtropical surface warming is balanced by a cooling down to 300 m in these latitudes (Fig. 13), consistent with reduced vertical mixing.

The addition of benthic geothermal heat input and double diffusion (not shown) has a comparatively small effect on the surface temperature, with a small localised cooling along the path of the ACC by 0.05 °C, but does produce a freshening of 0.05 psu over much of the Atlantic and the subtropical Pacific. Perhaps surprisingly, this change leads to negligible subsurface changes. The changes to the ice model give a surface cooling (of ~ 0.2 °C) and freshening (of ~ 0.1 psu) in the Southern Ocean and a similar cooling

in the Arctic. The bottom boundary layer and lakes modifications lead to similar but generally opposite regional surface changes, with the addition of a significant cooling of $\sim 0.2^\circ\text{C}$ north of the separated Gulf Stream, which is likely to be a result of a change in path of the deep western boundary current.

In summary, we find that the largest changes result firstly from the ocean code version upgrade from NEMO v3.2 to v3.4; and secondly from the changes to the parameters of the TKE scheme: namely, the parameters *rn_ebb*, *rn_mx10* and *nn_htau*. These have only a small effect on the surface errors, but in combination the two changes result in much more substantial improvement of the subsurface temperature field, as described in Sect. 5.2.

6 Summary and discussion

We have introduced a new ocean model configuration, GO5.0, developed jointly between the Met Office and NERC. This is an implementation of version 3.4 of the NEMO model, on the ORCA025 grid, with horizontal resolution of at least $1/4^\circ$ everywhere, together with the CICE sea ice model on the same grid. The GO5.0 model configuration is derived from the previous GO1 through an upgrade of the NEMO code version from version 3.2, and a set of parameter changes. A 30 yr integration of GO5.0, run with CORE2 surface forcing from 1976 to 2005, has been compared with GO1 with the same forcing. We have additionally described a set of 10 yr sensitivity studies carried out to attribute changes in the model performance to individual changes in the model physics.

The GO5.0 configuration was validated against observations during the final ten years of the 30 yr integration. It was found to have a generally warm surface bias, with respect to the EN3 climatological dataset, of $0.5\text{--}1^\circ\text{C}$ in the tropics, a cool bias of similar magnitude in the extra-tropics and a warm bias of around 2°C in much of the Southern Ocean. The surface salinity biases were again predominantly zonal, being up to 0.2 psu too salty close to the Equator and in subpolar regions and the Arctic, and

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



too fresh in the subtropics. In the Labrador Sea and in the North Atlantic subpolar gyre the surface waters are between 2° and 4 °C too warm, and around 1 psu too salty.

Both GO1 and GO5.0 model configurations showed good skills in simulating oceanic exchanges between North Atlantic, North Pacific and Arctic Oceans. The net oceanic exports from the Arctic Ocean and the contributions from the individual straits are within the uncertainties of the observational estimates. The main model bias is a more vigorous exchange between the Atlantic and Arctic Oceans manifesting itself in too strong (compared to observations) a northward flow of the buoyant warm Atlantic water and too strong a return flow of the dense Arctic water as the overflows across the Greenland–Scotland Ridge. The overturning circulation at 26° N in the Atlantic was correspondingly stronger than that observed, at 21 Sv. The transport in the Antarctic Circumpolar Current was 124 Sv, close to observed estimates, while the Indonesian Throughflow was significantly higher than observations, most likely because of insufficient mixing at the critical straits.

Comparison of the sea ice in the Northern Hemisphere in GO5.0 and in observations show that the model simulates the annual means, the interannual trend and the seasonal cycle well, although the model underestimates summer sea ice extent. In the Southern Hemisphere the sea ice extent again compares well with observations, although the recent rising trend in sea ice cover is not simulated in GO5.0, as is also the case in several other comparable models. Both the GO1 and GO5.0 underestimate sea ice volume in the Northern Hemisphere, with biases larger in summer than in winter. In the Southern Hemisphere the seasonal cycle of sea ice thickness is simulated correctly, with a moderate underestimation (of 22 % for GO5.0) of the hemisphere-averaged sea ice thickness.

The main differences between GO5.0 and GO1 were seen in the penetration of heat and salt into the interior ocean above the thermocline and in the representation of the seasonal cycle. The global mean warming, with a maximum at 200 m depth, was reduced from 0.7 to 0.3 °C, while the steady freshening trend at the same depth was also reduced by 10–20 %. Although the overall reduction in mixed layer depth from

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



GO1 to GO5.0 did not lead to unequivocal improvements in surface biases, wintertime mixed layers were consistently better represented in GO5.0, while the shallow bias in mixed layer depth and consequent warm surface bias in GO1 in tropical latitudes were significantly ameliorated in GO5.0.

To attribute the changes seen between GO1 and GO5.0, the physics modifications were applied incrementally (in most cases individually but some in pairs) starting from the original GO1 configuration. First of all the NEMO source code was upgraded from v3.2 to v3.4; then the model bathymetry was upgraded; the background vertical diffusivity and viscosity were increased; some of the TKE scheme parameters were adjusted; geothermal heat flux and double diffusion of tracers were added; schemes were added to represent a bottom boundary layer and lakes; and finally modifications were made to the ice model. It was found that all of the modifications led to changes with large spatial scales in the model surface and subsurface fields that were distinguishable from the eddy variability, but the dominant effects were traced to the code upgrade and to the TKE changes. These two changes, which affect mainly vertical mixing in the upper few hundred metres, were found to produce most of the reduction of the subsurface temperature and salinity biases of the model, along with the reduced errors in the seasonal cycle.

We conclude that GO5.0 represents a significant improvement in realism over the previous configuration of the Met Office ocean model, GO1. In particular, the improvements in the representation of vertical mixing (associated both with the code upgrade from the NEMO v3.2 and in the modifications to the TKE vertical mixing scheme in v3.4) lead to a more faithful simulation of the annual cycle in surface temperature and mixed layer depth, as well as to reduced subsurface drifts in the depth range 200–400 m.

There are clearly aspects of the GO5.0 configuration that need to be improved further. In particular, the subpolar North Atlantic and the Southern Ocean show substantial errors in both surface and subsurface fields that may be at least partly ascribed to deficiencies in model physics. Process Evaluation Groups (PEGs) have been set up within the JOMP programme specifically to address issues relating to the two aforementioned

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



regions, and work is ongoing in both cases. A further PEG has been initiated to examine the implementation of diapycnal mixing in the ocean model, which clearly has general implications for model performance.

In addition, GO5.0 does not contain several physics upgrades which are currently either available or under development in NEMO, and which offer potentially significant improvements in model realism. These include embedded sea ice (in which the base of the sea ice lies beneath the ocean surface and the ice displaces a non-zero volume of sea water); and the z-tilde modification to the vertical coordinate to reduce numerical mixing from high-frequency vertical motions (Leclair and Madec, 2011). The full non-linear free surface physics is available in NEMO v3.4, but not implemented in GO5.0; it is expected that this, along with z-tilde and the embedded ice, will be included in future implementations of the Global Ocean Model.

Appendix A

Code availability and model trunk and branches

The model code for NEMO v3.4 is available from the NEMO website (<http://nemo-ocean.eu>). On registering, individuals can access the FORTRAN code using the open source subversion software (<http://subversion.apache.org/>). The revision number of the base NEMO code (trunk) used for this paper is 3424. In addition we apply some modifications to the base code (branches). Please contact the authors for more information on these branches and how to obtain them.

The model code for CICE is freely available from the United States Los Alamos National Laboratory (<http://oceans11.lanl.gov/trac/CICE/wiki/SourceCode>), again using subversion. The revision number for the version used for this paper is 430 (trunk). Once again there are some additional modifications (branches) made for the purposes of this paper, and interested readers are requested to contact the authors for details.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



UK users with access to PUMA (cms.ncas.ac.uk/wiki/PumaService) can copy the job details (job id xhimo) and submit a duplicate job using the Met Office Unified Model User Interface (UMUI).

Appendix B

5 FPP keys used in GO5.0 (NEMO and CICE)

key_dynspgflt Filtered free surface
key_ldfslp Rotate diffusion operators (for tracer isopycnal diffusion)
key_traldf_c2d Geographically varying lateral tracer diffusion
key_dynldf_c2d Geographically varying lateral momentum diffusion
10 key_zdftke TKE scheme for vertical mixing
key_zdftmx Include tidal mixing scheme
key_zdfddm Include double diffusive mixing parameterisation
key_trabbl Include bottom boundary layer scheme

Appendix C

15 Ocean and ice namelists for GO5.0

These are included as Supplement.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Appendix D

Surface forcing

These are the CORE-2 forcing dataset (Large and Yeager, 2008), available at <http://rda.ucar.edu/datasets/ds260.2/>.

5 Appendix E

Other input files

Other files such as bathymetry, river runoff mask and interpolation weights for the surface forcing are required to run GO5.0. These can be obtained on request from the authors.

10 **Supplementary material related to this article is available online at**
**[http://www.geosci-model-dev-discuss.net/6/5747/2013/
gmdd-6-5747-2013-supplement.pdf](http://www.geosci-model-dev-discuss.net/6/5747/2013/gmdd-6-5747-2013-supplement.pdf)**

15 *Acknowledgements.* The NOC authors were funded by Natural Environment Research Council (UK) National Capability Funding. Met Office authors were supported by the Joint UK DECC/DEFRA Met Office Hadley Centre Climate Programme (GA01101). Funding support from the European Community's Seventh Framework Programme FP7/2007-2013 under grant agreement no. 283367 (MyOcean2) is gratefully acknowledged. The authors would like to thank Adam Blaker for his comments and suggestions, and Andrew Coward and Chris Harris for assistance with setting up and running the simulations. Ship-based Antarctic sea ice thickness data were provided by the SCAR Antarctic Sea Ice Processes and Climate (ASPeCt) program (20 aspect.antarctica.gov.au).



References

- Aksenov, Y., Bacon, S., Coward, A., and Holliday, N. P.: Polar Outflow from the Arctic Ocean: a high resolution model study, *J. Marine Syst.*, 83, 14–37, 2010.
- Aksenov, Y., Ivanov, V. V., Nurser, A. J. G., Bacon, S., Polyakov, I. V., Coward, A. C., Naveira-Garabato, A. C., and Beszczynska-Moeller, A.: The Arctic Circumpolar Boundary Current, *J. Geophys. Res.*, 116, C09017, doi:10.1029/2010JC006637, 2011.
- Amante, C. and Eakins, B. W.: ETOPO1 1 arc-minute global relief model: procedures, data sources and analysis, NOAA Technical Memorandum NESDIS NGDC-24, 19 pp., 2009.
- Arakawa, A.: Computational design of long-term numerical integration of the equations of fluid motion, *J. Comput. Phys.*, 1, 119–143, 1966.
- Arribas, A., Glover, M., Maidens, A., Peterson, K., Gordon, M., MacLachlan, C., Graham, R., Fereday, D., Camp, J., Scaife, A., Xavier, P., McLean, P., Colman, A., and Cusack, S.: The GloSea4 ensemble prediction system for seasonal forecasting, *Mon. Weather Rev.*, 139, 1891–1910, doi:10.1175/2010MWR3615.1, 2011.
- Barnier, B., Madec, G., Penduff, T., Molines, J.-M., Treguier, A.-M., Le Sommer, J., Beckmann, A., Biastoch, A., Böning, C., Dengg, J., Derval, C., Durand, E., Gulev, S., Remy, E., Talandier, C., Theetten, S., Maltrud, M., McClean, J., and De Cuevas, B.: Impact of partial steps and momentum advection schemes in a global ocean circulation model at eddy permitting resolution, *Ocean Dynam.*, 56, 543–567, doi:10.1007/s10236-006-0082-1, 2006.
- Beckmann, A. and Doscher, R.: A method for improved representation of dense water spreading over topography in geopotential-coordinate models, *J. Phys. Oceanogr.*, 27, 581–591, 1997.
- Bitz, C. M. and Lipscomb, W. H.: An energy-conserving thermodynamic model of sea ice, *J. Geophys. Res.*, 104, 15669–15677, 1999.
- Bitz, C. M., Holland, M., Eby, M., and Weaver, A. J.: Simulating the ice-thickness distribution in a coupled climate model, *J. Geophys. Res.*, 106, 2441–2463, 2001.
- Blaker, A. T., Hirschi, J. J.-M., McCarthy, G., Sinha, B., Taws, S., Marsh, R., Coward, A. C., and de Cuevas, B. A.: Historical analogues of the recent extreme minima observed in the Atlantic meridional overturning circulation at 26° N, *Clim. Dynam.*, in review, 2013.
- Blanke, B. and Delecluse, P.: Variability of the tropical Atlantic Ocean simulated by a general circulation model with two different mixed layer physics, *J. Phys. Oceanogr.*, 23, 1363–1388, 1993.

GMDD

6, 5747–5799, 2013

GO5.0

A. Megann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Bourdalle-Badie, R. and Treguier, A.-M.: A climatology of runoff for the global ocean-ice model ORCA025, Report, Mercator-Ocean, MOO-RP-425-365-MER, 2006.
- Brown, A., Milton, S., Cullen, M., Golding, B., Mitchell, J., and Shelley, A.: Unified modeling and prediction of weather and climate: a 25-year journey, *B. Am. Meteorol. Soc.*, 93, 1865–1877, 2012.
- 5 Calvert, D. and Siddorn, J.: Revised vertical mixing parameters for the UK community standard configuration of the global NEMO ocean model, Hadley Centre Technical Note, 95, 2013.
- Cavalleri, D. J., Parkinson, C. L., Gloersen, P., and Zwally, H.: Updated yearly, sea ice concentrations from Nimbus-7 SMMR and DMSP SSM/I-SSMIS passive microwave data, [indicate subset used], NASA DAAC at the National Snow and Ice Data Center, Boulder, Colorado, USA, 1996.
- 10 Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Hinton, T., Jones, C. D., Lid-dicoat, S., Martin, G., O'Connor, F., Rae, J., Senior, C., Totterdell, I., Woodward, S., Reich-ler, T., and Kim, J.: Evaluation of the HadGEM2 model, Met Office Hadley Centre Technical Note no. HCTN 74, available from Met Office, FitzRoy Road, Exeter EX1 3PB, available at: http://www.metoffice.gov.uk/media/pdf/8/7/HCTN_74.pdf (last access: 18 November 2013), 2008.
- 15 Cunningham, S. A., Alderson, S. G., King, B. A., and Brandon, M. A.: Transport and variability of the Antarctic Circumpolar Current in Drake Passage, *J. Geophys. Res.*, 108, 8084, doi:10.1029/2001JC001147, 2003.
- Cuny, J., Rhines, P., and Kwok, R. R.: Davis Strait volume, freshwater and heat fluxes, *Deep-Sea Res. Pt. I*, 52, 519–542, 2005.
- Curry, B., Lee, C. M., and Petrie, B.: Volume, freshwater, and heat fluxes through Davis Strait, 2004–2005, *J. Phys. Oceanogr.*, 41, 429–462, doi:10.1175/2010JPO4536.1, 2010.
- 25 Dai, A. and Trenberth, K. E.: Estimates of freshwater discharge from continents: latitudinal and seasonal variations, *J. Hydrometeorol.*, 3, 660–687, 2002.
- Danabasoglu, G., Large, G. W., and Briegleb, B.: Climate impacts of parameterized Nordic Sea overflows, *J. Geophys. Res.*, 115, C11005, doi:10.1029/2010JC006243, 2010.
- 30 Dawson, A., Matthews, A. J., Stevens, D. P., Roberts, M. J., and Vidale, P.-L.: Importance of oceanic resolution and mean state on the extra-tropical response to El Nino in a matrix of coupled models, *Clim. Dynam.*, 41, 1439–1452, doi:10.1007/s00382-012-1518-6, 2012.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: an examination of profile data and a profile-based climatology, *J. Geophys. Res.*, 109, C12003, doi:10.1029/2004JC002378, 2004.
- 5 Delworth, T., Rosati, A., Anderson, W., Adcroft, A., Balaji, V., Benson, R., Dixon, K., Griffies, S., Lee, H., Pacanowski, R., Vecchi, G. A., Wittenberg, A. T., Zeng, F., and Zhang, R.: Simulated climate and climate change in the GFDL CM2.5 high-resolution coupled climate model, *J. Climate*, 25, 2755–2781, 2012.
- Farneti, R., Delworth, T. L., Rosati, A. J., Griffies, S. M., and Zeng, F.: The role of mesoscale eddies in the rectification of the Southern Ocean response to climate change, *J. Phys. Oceanogr.*, 40, 1539–1557, doi:10.1175/2010JPO4353.1, 2010.
- 10 Gammelsrod, T., Leikvin, O., Lien, V., Budgell, W. P., Loeng, H., and Maslowski, W.: Mass and heat transports in the NE Barents Sea: observations and models, *J. Marine Syst.*, 75, 56–69, doi:10.1016/j.jmarsys.2008.07.010, 2009.
- Gaspar, P., Grégoris, Y., and Lefevre, J.-M.: A simple eddy kinetic energy model for simulations of the oceanic vertical mixing: tests at Station Papa and long-term upper ocean study site, *J. Geophys. Res.*, 95, 16179–16193, doi:10.1029/JC095iC09p16179, 1990.
- 15 Gnanadesikan, A., Griffies, S. M., and Samuels, B. L.: Effects in a climate model of slope tapering in neutral physics schemes, *Ocean Model.*, 16, 1–16, 2007.
- Gordon, C., Cooper, C., Senior, C. A., Banks, H., Gregory, J. M., Johns, T. C., Mitchell, J. F. B., and Wood, R. A.: The simulation of SST, sea ice extents and ocean heat transports in a version of the Hadley Centre coupled model without flux adjustments, *Clim. Dynam.*, 16, 147–168, 2000.
- 20 Greatbatch, R. J. and Peterson, K. A.: Interdecadal variability and oceanic thermohaline adjustment, *J. Geophys. Res.*, 101, 20467–20482, doi:10.1029/96JC01531, 1996.
- 25 Gregg, M.C., Sanford, T. B., and Winkel, D. P.: Reduced mixing from the breaking of internal waves in equatorial waters, *Nature*, 422, 513–515, 2003.
- Griffies, S. M., Pacanowski, R. C., and Hallberg, R. W.: Spurious diapycnal mixing associated with advection in a z-coordinate ocean model, *Mon. Weather Rev.*, 128, 538–564, 2000.
- 30 Griffies, S. M., Biastoch, A., Böning, C., Bryan, F., Danabasoglu, G., Chassignet, E. P., England, M. H., Gerdes, R., Haak, H., Hallberg, R. W., Hazeleger, W., Jungclaus, J., Large, W. G., Madec, G., Pirani, A., Samuels, B. L., Scheinert, M., Sen Gupta, A., Severijns, C. A., Simmons, H. L., Treguier, A.-M., Winton, M., Yeager, S., Yin, J.: Co-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ordinated Ocean-ice Reference Experiments (COREs), *J. Ocean Model.*, 26, 1–46, doi:10.1016/j.ocemod.2008.08.007, 2008.

Hallberg, R. and Gnanadesikan, A.: The role of eddies in determining the structure and response of the wind-driven Southern Hemisphere overturning: results from the Modeling Eddies in the Southern Ocean (MESO) project, *J. Phys. Oceanogr.*, 36, 2232–2252, 2006.

Hewitt, H. T., Copsey, D., Culverwell, I. D., Harris, C. M., Hill, R. S. R., Keen, A. B., McLaren, A. J., and Hunke, E. C.: Design and implementation of the infrastructure of HadGEM3: the next-generation Met Office climate modelling system, *Geosci. Model Dev.*, 4, 223–253, doi:10.5194/gmd-4-223-2011, 2011.

Hirschi, J. J.-M., Blaker, A. T., Sinha, B., Coward, A., de Cuevas, B., Alderson, S., and Madec, G.: Chaotic variability of the meridional overturning circulation on subannual to interannual timescales, *Ocean Sci.*, 9, 805–823, doi:10.5194/os-9-805-2013, 2013.

Holland, P. R. and Kwok, R.: Wind-driven trends in Antarctic sea-ice drift, *Nature Geosci.*, 5, 872–875, doi:10.1038/ngeo1627, 2012.

Houghton, J. T., Ding, Y., Griggs, D. J., Noguer, M., van der Linden, P. J., and Xiaosu, D.: *Climate Change 2001: The Scientific Basis*, Cambridge University Press, 944 pp., 2001.

Hunke, E. C. and Dukowicz, J. K.: An Elastic–Viscous–Plastic Model for sea ice dynamics, *J. Phys. Oceanogr.*, 27, 1849–1867, 1997.

Hunke, E. C. and Lipscomb, W. H.: CICE: The Los Alamos Sea Ice Model, Documentation and Software User’s Manual, Version 4.1. Tech. Rep. LA-CC- 06-012, Los Alamos National Laboratory, Los Alamos, New Mexico. available at: <http://oceans11.lanl.gov/trac/CICE> (last access: 18 November 2013), 2010.

Ingleby, B. and Huddleston, M.: Quality control of ocean temperature and salinity profiles – historical and real-time data, *J. Marine Syst.*, 65, 158–175, 2007.

IOC, IHO and BODC: Centenary Edition of the GEBCO Digital Atlas, published on CD-ROM on behalf of the Intergovernmental Oceanographic Commission and the International Hydrographic Organization as part of the General Bathymetric Chart of the Oceans, British Oceanographic Data Centre, Liverpool, UK, 2003.

Johns, T. C., Durman, C. F., Banks, H. T., Roberts, M. J., McLaren, A. J., Ridley, J. K., Senior, C. A., Williams, K. D., Jones, A., Rickard, G. J., Cusack, S., Ingram, W. J., Crucifix, M., Sexton, D. M. H., Joshi, M. M., Dong, B.-W., Spencer, H., Hill, R. S. R., Gregory, J. M., Keen, A. B., Pardaens, A. K., Lowe, J. A., Bodas-Salcedo, A., Stark, S., and Searl, Y.: The New

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Hadley Centre Climate Model (HadGEM1): evaluation of coupled simulations, *J. Climate*, 19, 1327–1353, 2006.

Kennedy, J. J., Rayner, N. A., Smith, R. O., Saunby, M., and Parker, D. E.: Reassessing biases and other uncertainties in sea-surface temperature observations measured in situ since 1850 – Part 2: Biases and homogenisation, *J. Geophys. Res.*, 116, D14104, doi:10.1029/2010JD015220, 2011.

Koch-Larrouy, A., Madec, G., Blanke, B., and Molcard, R.: Water mass transformation along the Indonesian throughflow in an OGCM, *Ocean Dynam.*, 58, 289–309, 2008.

Large, W. G. and Yeager, S. G.: The global climatology of an interannually varying air-sea flux data set, *Clim. Dynam.*, 33, 341–364, doi:10.1007/s00382-008-0441-3, 2008.

Leclair, M. and Madec, G.: *Ocean Model.*, 37, 139–152, 2011.

Lee, M.-M., Coward, A. C., and Nurser, A. J. G.: Spurious diapycnal mixing of the deep waters in an eddy-permitting global ocean model, *J. Phys. Oceanog.*, 32, 1522–1535, 2002.

Lohmann, G., Gerdes, R., and Chen, D.: Sensitivity of the thermohaline circulation in coupled oceanic GCM-atmospheric EBM experiments, *Clim. Dynam.*, 12, 403–416, 1996.

Madec, G., NEMO – the OPA9 ocean engine: Note du Pole de Modelisation, Institut Pierre-Simon Laplace, 1:100 pp., available at: <http://www.nemo-ocean.eu> (last access: 18 November 2013), 2008.

McCarthy, G., Frajka-Williams, E., Johns, W. E., Baringer, M. O., Meinen, C. S., Bryden, H. L., Rayner, D., Duchez, A., Roberts, C., and Cunningham, S. A.: Observed interannual variability of the Atlantic meridional overturning circulation at 26.5° N, *Geophys. Res. Lett.*, 39, L19609, doi:10.1029/2012GL052933, 2012.

Megann, A. P., New, A. L., Blaker, A. T., and Sinha, B.: The sensitivity of a coupled climate model to its ocean component, *J. Climate*, 23, 5126–5150, 2010.

Meijers, A. J. S., Shuckburgh, E., Bruneau, N., Sallee, J.-B., T. J. Bracegirdle, and Wang, Z.: Representation of the Antarctic Circumpolar Current in the CMIP5 climate models and future changes under warming scenarios, *J. Geophys. Res.*, 117, C12008, doi:10.1029/2012JC008412, 2012.

Meredith, M. P., Woodworth, P. L., Chereskin, T. K., Marshall, D. P., Allison, L. C., Bigg, G. R., Donohue, K., Heywood, K. J., Hughes, C. W., Hibbert, A., Hogg, A. McC., Johnson, H. L., Jullion, L., King, B. A., Leach, H., Lenn, Y.-D., Morales Maqueda, M. A., Munday, D. R., Naveira Garabato, A. C., Provost, C., Sallée, J.-B., and Sprintall, J.: Sustained monitoring of

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- the Southern Ocean at Drake Passage: past achievements and future priorities, *Rev. Geophys.*, 49, RG4005, doi:10.1029/2010RG000348, 2011.
- Merryfield, W. J., Holloway, G., and Gargett, A. E.: A global ocean model with double-diffusive mixing, *J. Phys. Ocean.*, 29, 1124–1142, 1999.
- 5 Olsen, S. M., Hansen, B., Quadfasel, D., and Østerhus, S.: Observed and modelled stability of overflow across the Greenland–Scotland ridge, *Nature*, 455, 519–522, 2008.
- Østerhus, S., Turrell, W. R., Jónsson, S., and Hansen, B.: Measured volume, heat, and salt fluxes from the Atlantic to the Arctic Mediterranean, *Geophys. Res. Lett.*, 32, L07603, doi:10.1029/2004GL022188, 2005.
- 10 Penduff, T., Le Sommer, J., Barnier, B., Treguier, A.-M., Molines, J.-M., and Madec, G.: Influence of numerical schemes on current-topography interactions in 1/4° global ocean simulations, *Ocean Sci.*, 3, 509–524, doi:10.5194/os-3-509-2007, 2007.
- Penduff, T., Juza, M., Brodeau, L., Smith, G. C., Barnier, B., Molines, J.-M., Treguier, A.-M., and Madec, G.: Impact of global ocean model resolution on sea-level variability with emphasis
- 15 on interannual time scales, *Ocean Sci.*, 6, 269–284, doi:10.5194/os-6-269-2010, 2010.
- Rahmstorf, S. and Willebrand, J.: The role of temperature feedback in stabilizing the thermohaline circulation, *J. Phys. Oceanogr.*, 25, 787–805, doi:10.1175/1520-0485(1995)025<0787:TROTFI>2.0.CO;2, 1995.
- Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P.,
- 20 Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, *J. Geophys. Res.*, 108, 4407, doi:10.1029/2002JD002670, 2003.
- Remy, E., Siefritz, L., Fleury, L., and Bremond, M.: Construction de la bathymetrie pour la configuration modele ORCA025, CERFACS Report, May 2003, 2003.
- 25 Reynolds, R. W., Rayner, N. A., Smith, T. M., Stokes, D. C., and Wang, W.: An improved in situ and satellite SST analysis for climate, *J. Climate*, 15, 1609–1625, 2002.
- Roberts, C. D., Waters, J., Peterson, K. A., Palmer, M. D., McCarthy, G. D., Frajka-Williams, E., Haines, K., Lea, D. J., Martin, M. J., Storkey, D., Blockley, E. W., and Zuo, H.: Atmosphere drives recent interannual variability of the Atlantic meridional overturning circulation at 26.5° N, *Geophys. Res. Lett.*, doi:10.1002/grl.50930, in press, 2013.
- 30 Roberts, M. J., Clayton, A., Demory, M.-E., Donners, J., Vidale, P.-L., Norton, W., Shaffrey, L., Stevens, D. P., Stevens, I., Wood, R. A., and Slingo, J.: Impact of resolution on the tropical pacific circulation in a matrix of coupled models, *J. Climate*, 22, 2541–2556, 2009.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Saunders, P. P., Cunningham, S.A., de Cuevas, B. A., and Coward, A. C.: Comments on “Decadal Changes in the North Atlantic and Pacific Meridional Overturning Circulation and Heat Flux”, *J. Phys. Ocean.*, 38, 2104–2107, 2008.

Scaife, A. A., Copsey, D., Gordon, C., Harris, C., Hinton, T., Keeley, S., O’Neill, A., Roberts, M., and Williams, K.: Improved Atlantic winter blocking in a climate model, *Geophys. Res. Lett.*, 38, L23703, doi:10.1029/2011GL049573, 2011.

Skagseth, Ø., Furevik, T., Ingvaldsen, R., Loeng, H., Mork, K. A., Orvik, K. A., Ozhigin, V.: Volume and heat transports to the Arctic Ocean via the Norwegian and Barents Seas, in: *Arctic–Subarctic Ocean Fluxes: Defining the Role of the Northern Seas in Climate*, edited by: Dickson, R. R., Meincke, J., and Rhines, P., Springer, the Netherlands, 45–64, 2008.

Semtner, A. J.: A model for the thermodynamic growth of sea ice in numerical investigations of climate, *J. Phys. Oceanogr.*, 6, 379–389, 1976.

Shaffrey, L. C., Stevens, I., Norton, W. A., Roberts, M. J., Vidale, P. L., Harle, J. D., Jarrah, A., Stevens, D. P., Woodage, M. J., Demory, M. E., Donners, J., Clark, D. B., Clayton, A., Cole, J. W., Wilson, S. S., Connolley, W. M., Davies, T. M., Iwi, A. M., Johns, T. C., King, J. C., New, A. L., Slingo, J. M., Slingo, A., Steenman-Clark, L., and Martin, G. M.: U.K. HiGEM: The new U.K. high-resolution global environment model – model description and basic evaluation, *J. Climate*, 22, 1861–1896, 2009.

Simmons, H., Jayne, S., Laurent, L. S., and Weaver, A.: Tidally driven mixing in a numerical model of the ocean general circulation, *Ocean Model.*, 6, 245–263, 2004.

Smeed, D. A., McCarthy, G., Cunningham, S. A., Frajka-Williams, E., Rayner, D., Johns, W. E., Meinen, C. S., Baringer, M. O., Moat, B. I., Duchez, A., and Bryden, H. L.: Observed decline of the Atlantic Meridional Overturning Circulation 2004 to 2012, *Ocean Sci. Discuss.*, 10, 1619–1645, doi:10.5194/osd-10-1619-2013, 2013.

Smith, D. M., Cusack, S., Colman, A. W., Folland, C. K., Harris, G. R., and Murphy, J. M.: Improved surface temperature prediction for the coming decade from a global climate model, *Science*, 317, 796–799, doi:10.1126/science.1139540, 2007.

Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L.: *Climate Change 2007: The Physical Science Basis*, Cambridge University Press, 996 pp., 2007.

Soto-Navarro, J, Criado-Aldeanueva, F., García-Lafuente, J., and Sánchez-Román, A.: Estimation of the Atlantic inflow through the Strait of Gibraltar from climatological and in situ data, *J. Geophys. Res.*, 115, C10023, doi:10.1029/2010JC006302, 2010.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Sprintall, J., Wijffels, S. E., Molcard, R., and Jaya, I.: Direct estimates of the Indonesian Throughflow entering the Indian Ocean: 2004–2006, *J. Geophys. Res.*, 114, C07001, doi:10.1029/2008JC005257, 2009.
- Storkey, D., Blockley, E. W., Furner, R., Guiavarc'h, C., Lea, D., Martin, M. J., Barciela, R. M., Hines, A., Hyder, P., and Siddorn, J. R.: Forecasting the ocean state using NEMO: the new FOAM system, *Journal of Operational Oceanography*, 3, 3–15, 2010.
- Tansley, C. E. and Marshall, D. P.: On the dynamics of wind-driven circumpolar currents, *J. Phys. Oceanogr.*, 31, 3258–3273, 2001a.
- Thomas, M. D. and Zhai, X.: Eddy-induced variability of the meridional overturning circulation in a model of the North Atlantic, *Geophys. Res. Lett.*, 40, 1–6, doi:10.1002/grl.50532, 2013.
- Viebahn, J. and Eden, C.: Towards the impact of eddies on the response of the southern ocean to climate change, *Ocean Model.*, 34, 150–165, 2010.
- Williamson, D., Goldstein, M., Allison, L., Blaker, A., Challenor, P., and Jackson, L.: History matching for the quantification and reduction of parametric uncertainty in climate model projections. *Clim. Dynam.*, 41, 1703–1729, doi:10.1007/s00382-013-1896-4, 2013.
- Woodgate, R. A., Weingartner, T. J., and Lindsay, R.: Observed increases in Bering Strait oceanic fluxes from the Pacific to the Arctic from 2001 to 2011 and their impacts on the Arctic Ocean water column, *Geophys. Res. Lett.*, 39, L24603, doi:10.1029/2012GL054092, 2012.
- Worby, A. P., Geiger, C., Paget, M. J., van Woert, M., F Ackley, S., and DeLiberty, T.: Thickness distribution of Antarctic sea ice, *J. Geophys. Res.*, 113, C05S92, doi:10.1029/2007JC004254, 2008.
- Yeager, S. G. and Jochum, M.: The connection between Labrador Sea buoyancy loss, deep western boundary current strength, and Gulf Stream path in an ocean circulation model, *Ocean Model.*, 30, 207–224, 2009.
- Zalesak, S. T.: Fully multidimensional flux corrected transport algorithms for fluids, *J. Comput. Phys.*, 31, 335–362, 1979.
- Zhang, J., Thomas, D. R., Rothrock, D. A., Lindsay, R. W., Yu, Y., and Kwok, R.: Assimilation of ice motion observations and comparisons with submarine ice thickness data, *J. Geophys. Res.*, 108, 3170, doi:10.1029/2001JC001041, 2003.

Table 1. Parameter changes between GO1 and GO5.0.

Parameter	GO1 (where different from GO5)	GO5
Horizontal bilaplacian viscosity	Same as in GO5.0	$-1.5 \times 10^{11} \text{ m}^4 \text{ s}^{-1}$
Isonutral laplacian tracer diffusion	Same as in GO5.0	$300 \text{ m}^2 \text{ s}^{-1}$
Background vertical viscosity	$1.0 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$	$1.2 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$
Background vertical diffusivity	$1.0 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$	$1.2 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$
Energy coefficient for Craig and Banner (1994) surface wave breaking parameterisation	60.0	67.83
Length scale for near-inertial wave breaking parameterisation	0.5 m in tropics, rising to 30 m at midlatitudes	10 m everywhere
Minimum value of surface mixing length scale	0.01 m	0.04 m
Minimum value of interior mixing length scale	0.001 m	0.01 m

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 2. Summary of integrations carried out.

Run	UM job id	NEMO vn.	rn_av0 $\times 10^{-5}$	bathy	rn_mxl0	rn_ebb	nn_htau	nn_geoflx	Run length (yr)
GO1	xexoc	3.2	1.0	G70	n/a	60.0	1	0	30
N3.4	xhiml	3.4	1.0	G70	0.001	60.0	1	0	30
N3.4_bath	xhimj	3.4	1.0	GO5	0.001	60.0	1	0	10
N3.4_vmix	xhkfg	3.4	1.2	GO5	0.001	60.0	1	0	10
N3.4_tke	xhkfi	3.4	1.2	GO5	0.04	67.83	0	0	10
N3.4_geo	xhimp	3.4	1.2	GO5	0.04	67.83	0	2	10
N3.4_ice	xhimm	3.4	1.2	GO5	0.04	67.83	0	2	10
N3.4_bbll	xhimn	3.4	1.2	GO5	0.04	67.83	0	2	10
GO5.0	xhimo	3.4	1.2	GO5	0.04	67.83	0	2	30

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 3. Volume transports (Sv), observed and model mean values and their standard deviations. Model values are means over the last 10 yr of the 30 yr spin up. Model standard deviations are obtained from the 5 day averages. Sign convention is positive northwards and eastwards, and is negative southwards and westwards.

Location	Observed value	GO1	GO5.0
AMOC at 26N	18.5 ± 1^a	21.0 ± 4.2	22.0 ± 4.2
Barents Sea Opening net	$2.8 \pm 0.6^{b,c,d}$	3.3 ± 2.0	3.0 ± 2.0
Fram Strait net	-2.3 ± 4.3^e	-1.9 ± 2.4	-1.6 ± 2.3
Denmark Strait net	$(-6.0 \text{ to } -3.6)^d$	-3.4 ± 3.3	-3.3 ± 3.3
Denmark Strait overflow ^m	-2.9 ± 0.6^f	-5.3 ± 2.9	-4.34 ± 2.1
Iceland–Faeroes net	2.8 ± 0.5^g	2.72 ± 1.2	2.6 ± 1.2
Iceland–Faeroes overflow ^m	-1.0 ± 0.5^f	-0.9 ± 0.5	-0.9 ± 0.5
Faeroes–Scotland net	1.8 ± 0.5^g	1.4 ± 2.3	1.7 ± 2.3
Faeroes–Scotland overflow ^m	-2.4 ± 0.4^f	-3.1 ± 0.8	-3.1 ± 0.9
Total Greenland–Scotland inflow ⁿ	8.5 ± 1.0^g	9.3 ± 1.8	10.0 ± 1.7
Bering Strait net	$0.8 [1.1^i] \pm 0.2^i$	1.3 ± 0.9	1.4 ± 0.9
Davis Strait net ^o	$-2.6 \pm 1.0 \text{ to } -2.3 \pm 0.7^{e,h}$	-2.9 ± 1.2	-3.0 ± 1.1
Drake Passage	135 ± 20^j	119 ± 8	124 ± 8
Indonesian Throughflow	15 ± 4^k	-19.7 ± 5.4	-19.8 ± 5.5

^a McCarthy et al., 2012,

^b Gammelsrod et al., 2009,

^c Skagseth, et al., 2008,

^d Aksenov et al., 2010,

^e Curry et al., 2011,

^f Olsen et al., 2008,

^g Østerhus et al., 2005,

^h Cuny et al., 2005,

ⁱ Woodgate et al., 2012,

^j Cunningham et al., 2003,

^k Sprintall et al., 2009,

^l climatological transport with the estimate for 2011 in parenthesis,

^m southward transport of waters with $\sigma_\theta > 27.8$,

ⁿ Atlantic inflow derived as the residual flow after subtracting the southward transport of waters with $\sigma_\theta > 27.8$,

^o including transports on the West Greenland Shelf.




Fig. 1. Surface biases in years 1996–2005 of GO5.0: **(a)** mean surface temperature bias with respect to the Reynolds et al. (2002) climatology; and **(b)** mean surface salinity bias with respect to the EN3 climatology.




Fig. 2. Seasonal cycle of mixed-layer depth (MLD): **(a)** minimum monthly MLD in years 1996–2005 of GO5.0; **(b)** minimum monthly MLD in the de Boyet Montégut et al. (2004) climatology; **(c)** maximum monthly MLD in years 1996–2005 of GO5.0; and **(d)** maximum monthly MLD in the de Boyet Montégut et al. (2004) climatology.




Fig. 3. Zonal mean (a) temperature and (b) salinity biases in years 1996–2005 of GO5.0. The solid contours are of the zonal mean potential density σ_0 .

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 4. (a) Time series of annual mean Atlantic meridional overturning circulation (AMOC, solid lines) at 26° N in GO1 (blue) and GO5.0 (black), with the potential density σ_0 in the upper 200 m in the central Labrador Sea (dashed lines); and (b) mean Atlantic overturning streamfunction in GO1 (left) and GO5.0 (right).




Fig. 5. Time series of **(a)** area-integrated Arctic sea ice extent; **(b)** Arctic sea ice volume; **(c)** Southern Ocean sea ice extent; and **(d)** Southern Ocean sea ice volume. Monthly and annual mean time series are shown by the thin and thick lines respectively. The GO5.0 simulations are in red, while the observed sea ice extent from passive microwave satellite imagery and the estimated sea ice volume from PIOMAS are in blue.




Fig. 6. Long-term seasonal mean of the simulated sea ice extent in GO5.0 calculated for the last ten years of the integrations 1996–2005 and in the HadISST observational dataset, averaged for the same period: Arctic winter (DJF) ice extent in **(a)** GO5.0 and **(b)** observations; Antarctic winter (JJA) ice extent in **(c)** GO5.0 and **(d)** observations; Arctic summer (JJA) ice extent in **(e)** GO5.0 and **(f)** observations; and Antarctic summer (DJF) ice extent in **(g)** GO5.0 and **(h)** observations.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 7. Subsurface drift in the models as a function of depth: **(a)** GO1 temperature drift; **(b)** GO5.0 temperature drift; **(c)** GO1 salinity drift; and **(d)** GO5.0 salinity drift. The drift is defined as the difference of the zonally-averaged annual mean fields in any given year from that in the first year of integration.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fig. 8. Seasonal sea surface temperature (SST) biases against Reynolds et al. (2002) climatology: boreal winter (DJF) biases in **(a)** GO1 and **(b)** GO5.0; and boreal summer (JJA) biases in **(c)** GO1 and **(d)** GO5.0.




Fig. 9. Monthly sea surface temperature (SST) and mixed layer depth (MLD) biases against Reynolds et al. (2002) and de Boyet Montégut et al. (2002) climatology, respectively, as a function of latitude: **(a)** GO1 SST; **(b)** GO5.0 SST; **(c)** GO1 MLD; and **(d)** GO5.0 (monthly) MLD.




Fig. 10. Effect on sea surface fields in years 1981–1985 of ocean code upgrade from v3.2 (GO1) to v3.4 (N3.4): **(a)** GO1 SST bias; **(b)** N3.4 SST bias; **(c)** N3.4 minus GO1 SST; **(d)** GO1 SSS bias; **(e)** N3.4 SSS bias; and **(f)** N3.4 minus GO1 SSS.




Fig. 11. Effect on zonal mean temperature in years 1981–1985 of code upgrade from NEMO v3.2 (GO1) and v3.4 (experiment N3.4) in years 1981–1985. **(a)** bias in GO1; **(b)** bias in N3.4; and **(c)** difference N3.4 minus GO1. The solid contours are of the zonal mean potential density σ_0 in N3.4.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 12. Effect on sea surface fields in years 1981–1985 of TKE scheme changes (from experiment N3.4_vmix to N3.4_tke): **(a)** N3.4_vmix SST bias; **(b)** N3.4_tke SST bias; **(c)** N3.4_tke minus N3.4_vmix SST; **(d)** N3.4_vmix SSS bias; **(e)** N3.4_tke SSS bias; and **(f)** N3.4_tke minus N3.4_vmix SSS.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 13. Effect on zonal mean temperature in years 1981–1985 of TKE scheme changes (from experiment N3.4_vmix to N3.4_tke). **(a)** bias in N3.4_vmix; **(b)** Bias in N3.4_tke; and **(c)** difference N3.4_tke minus N3.4_vmix. The solid contours are of the zonal mean potential density σ_0 in N3.4_tke.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)