



MOMBA 1.1 – a high-resolution Baltic Sea configuration of GFDL’s modular ocean model

H. Dietze, U. Löptien, and K. Getzlaff

GEOMAR, Helmholtz Centre for Ocean Research Kiel, Düsternbrooker Weg 20,
24105 Kiel, Germany

Received: 18 March 2014 – Accepted: 20 March 2014 – Published: 3 April 2014

Correspondence to: H. Dietze (hdietze@geomar.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We present a new coupled ocean circulation – ice model configuration of the Baltic Sea. The model features, contrary to most existing configurations, a high horizontal resolution of ≈ 1 nautical mile which is eddy resolving over much of the domain. The vertical discretisation comprises a total of 47 vertical levels. Results from a 1987 to 1999 hind cast simulation show that the model's fidelity is competitive. As suggested by a comparison with sea surface temperatures observed from space, this applies especially to near-surface processes. Hence, the configuration is well suited to serve as a nucleus of a full-fledged coupled ocean circulation biogeochemical model (which is yet to be developed). A caveat is that the model fails to reproduce major inflow events. We trace this back to spurious vertical circulation patterns at the sills which may well be endemic to high resolution models based on geopotential coordinates. Further, we present indications that – so far neglected – eddy/wind effects exert significant control on wind-induced up- and downwelling.

1 Introduction

The Baltic is a shallow sea with an average depth of less than 60 m, situated in central northern Europe between 53 to 67° N and 10 to 30° E. It is a brackish sea where intermittent salt water inflows from the North Sea mix with river runoff and a precipitation which, on average, exceeds evaporation. 85 million people distributed over 14 countries live in its drainage basin (e.g., Leppäranta and Myrberg, 2009), and there is growing concern about their effect on the pelagic ecosystem (e.g., Neumann et al., 2012).

A major problem is anthropogenic eutrophication (e.g., Pawlak et al., 2009; Wasmund and Uhlig, 2003). Nutrients, such as phosphate and nitrate, stemming from, e.g., sewage and fertiliser usage in agriculture, drive enhanced primary production by phytoplankton at the sun-lit surface of the Baltic Sea. The associated export of organic matter

GMDD

7, 2065–2124, 2014

MOMBA 1.1

H. Dietze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

to depth drives increased oxygen consumption (e.g., Karlson et al., 2002). The implications of the resulting decreased oxygen concentrations at depth are manifold. One major problem, e.g., is that decreasing oxygen reduces the volume of seawater which hosts the specific combination of density and dissolved oxygen that is so indispensable for cod eggs to survive. This hampers the recruitment of cod (e.g., MacKenzie et al., 2000), which is an ecologically and economically important fish species in the Baltic.

An additional problem is that depleted oxygen concentrations in bottom waters can trigger fundamental changes in organic matter cycling when certain thresholds are reached. Among the changes are two, which are thought to set the scene for a dominance of cyanobacteria: (1) denitrification and, (2) the release of dissolved inorganic phosphorous (bound to iron oxyhydroxides) from the sediments to the water column. This is of concern because cyanobacteria can release toxins and thereby constrain recreational use of the water body.

It is considered possible to reduce the nutrient load entering the Baltic Sea so that its assets (e.g., fisheries and tourism) are sustained for generations to come. But, such a reduction calls for either increased cost for sewage treatment facilities or reduced profits from crops and cattle breeding. Because these are large scale economic decisions effecting highly industrialised countries, the development of projection tools appears mandatory. Further, because any decision effecting significant impact must be taken in concert with all the countries in the drainage basin (not all of which actually border on the sea), the development of, and access to these tools, should be as transparent as possible.

This paper describes a cutting-edge Baltic Sea configuration of an ocean circulation model applied to simulate a hindcast for the period 1987 to 1999. With a competitive horizontal resolution of ≈ 1 nm it is eddy-permitting in most of the Baltic (Fennel et al., 1991). We envision to develop, based on this nucleus, a full-fledged coupled ice–ocean circulation–biogeochemical model. The biogeochemical module will be added in forthcoming projects; step by step as advised by Arhonditsis and Brett (2004) (and bearing in mind the harsh criticism by Flynn, 2005; Anderson, 2005), and starting with simple

approaches similar to the one described in Lehmann et al. (2014b). The underlying framework is the MOM4p1 release of NOAA's Geophysical Fluid Dynamics Laboratory Modular Ocean Model (Griffies, 2009). The code was chosen because of its exceptional documentation, maturity, modularity and user community.

In this study we aim to demonstrate that our configuration of the coupled ice–ocean circulation model is competitive by comparing our hind cast with observations. Further, we give guidance on the research questions which could be addressed with the model in its present state. The configuration is distributed, free of charge, by the Biogeochemical Modelling group at the Helmholtz Centre for Ocean Research, Kiel, Germany (<http://www.geomar.de/en/research/fb2/fb2-bm/>).

The project is dubbed **MOMBA**, a configuration of GFDL's **Modular Ocean Model** version 4p1 for the **BAltic** Sea. It is accessible via <http://www.baltic-ocean.org>.

2 Model setup

2.1 Grid and bathymetry

The model domain covers the area from 4.2 to 30.3° E, and 53.8 to 66.0° N (Fig. 1a). There are no open boundaries. The North Sea is represented by a simplified basin with solid walls in the North, South and West. The horizontal discretisation is an Arakawa B-grid (Arakawa and Lamp, 1977) and comprises 603 × 588 tracer grid boxes.

The longitudinal resolution is a function of longitude, starting with 0.18° at the western boundary, increasing to 0.0375° at 9.36° E and decreasing slightly thereafter to 0.03998 at the eastern boundary. The meridional resolution is 0.02°. Expressed as the harmonic mean of longitudinal and meridional resolution, this corresponds to a horizontal resolution of 0.9 to 1.3 nautical miles in the Baltic Sea, and up to 2 nautical miles in the North Sea. This resolution is exceptionally high and resolves almost the full range of first baroclinic Rossby radii in the Baltic Sea (e.g., Fennel et al., 1991).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The vertical discretisation comprises a total of 47 levels. Figure 2 shows the nominal depth and thickness of each level. We use the z^* (z^*) coordinate (Stacey et al., 1995; Adcroft and Campin, 2004) in the vertical, i.e., the depth and thickness of each level varies with time. z^* is calculated as a function of nominal depth (z), water depth (H) and sea surface height (η) (which varies with time because we use a free surface):

$$z^* = H \left(\frac{z - \eta}{H + \eta} \right) \quad (1)$$

(Eq. 6.6 in Griffies, 2009). The approach overcomes the problem with vanishing surface grid boxes which appears in generic z -level discretisation when sea surface height variations are of similar magnitude than the thickness of the uppermost grid box. Hence, the z^* approach allows for higher vertical resolution at the surface than the traditional z -level coordinates.

The model bathymetry is interpolated from the high-resolution data base compiled by Seifert et al. (2001) (<http://www.io-warnemuende.de/topografie-der-ostsee.html>). We use partial cells. The interpolation scheme is bilinear and takes the respective 10 nearest neighbours into account. Water columns shallower than 3 grid boxes are deepened to 3 grid boxes in order to ensure numerical stability. The model bathymetry is smoothed with a filter similar to the Shapiro filter (Shapiro, 1970). The filter weights are 0.25, 0.5, 0.25. The filtering procedure can only decrease the bottom depth, i.e. essentially, it fills rough holes. The filter is applied three times consecutively. The resulting bathymetry contained lakes which we filled after visual inspection. In addition, we filled narrow inlets which received large river runoffs and developed numerical oscillations (and river mouths were shifted accordingly thereafter).

In total, MOMBA has 1 483 236 wet tracer grid boxes which is still insufficient to resolve the complex bathymetry and coastline (Fig. 1b–d). This is especially problematic in the narrow and shallow Danish Straits (Øresund, Great Belt, Little Belt) which control the exchange of water between the North Sea and the Baltic Sea (Fig. 1b). Aggravating in that respect is that the horizontal Arakawa B-grid discretisation necessitates a

“no-slip” boundary condition for velocity grid point bordering the bottom. We suspect that this dampens our simulated transport through the Danish Straits. In order to compensate for this spurious effect, we deepen the bathymetry at several grid points in the Straits as documented in Table 1.

2.2 Initial conditions and sponges

Situated in central northern Europe, the Baltic Sea is considered to be among the best-sampled regions worldwide. Even so, because of its high temporal and spacial variability it is not straightforward to derive initial conditions for a high-resolution ocean circulation model which features more than one million grid boxes.

We use temperature and salinity bottle data archived by the International Council for the Exploration of the Sea (hereafter ICES, <http://ocean.ices.dk>). The data were extracted upon request by E. J. Green at ICES in February 2009. All data, from 1961 to 2008, were binned into monthly boxes of 6 nm horizontal resolution with a vertical thickness corresponding to the model grid (Sect. 2.1). These monthly binned data were subsequently averaged over all months of the whole 1961 to 2008 period. In a third step we applied an objective mapping with a horizontal decay length scale of 10 nm in order to smooth and extrapolate the data. Our approach results in a smooth 3-dimensional, plausible temperature and salinity distribution throughout the whole Baltic Sea. In order to identify a date at which these smooth temperature and salinity fields resemble actual conditions, we compared the smoothed fields visually to observations of near-bottom salinity (as provided within the Shark database (<http://produkter.smhi.se/pshark/>) at SMHI (Swedish Meteorological and Hydrological Institute, Norrköping, Sweden)). This comparison, based on the monitoring stations BY5, BY15 and BY31 (cf. Fig. 1a) suggests, that our smoothed temperature and salinity fields are similar to actual conditions on 1 January 1987. Hence, we start our simulation with the smoothed fields on 1 January 1987.

West of 9.3° E temperatures (and salinities) are restored to prescribed temperatures (salinities) using an exponential law with a restoring timescale of 1 year (1 day). Salinity



is restored to 35 PSU throughout the water column. Temperature is restored to the temperature field described above (i.e., to the initial conditions) below 80 m. Above 80 m we do not apply temperature restoring such that a seasonal cycle can freely evolve.

5 2.3 Sea surface height restoring

On average, the Baltic Sea exports fresh water at the surface to the North Sea, and receives deep, saline inflows, from the North Sea. The inflow events are intermittent and show a strong correlation with atmospheric circulation patterns (e.g., Schinke and Matthäus, 1998; Wyrski, 1954) because the winds drive differences in sea surface height between the Baltic Sea and the North Sea. Their associated pressure gradients, in turn, drive the exchange between the Baltic Sea and the North Sea.

As described in Sect. 2.1, the North Sea is represented by a simplified basin. This retards the simulation of sea surface height (SSH) in the North Sea which, in turn, leads to an unrealistic exchange between the Baltic Sea and the North Sea. In order to compensate for this we restore sea surface height between the western edge of the model domain and 6.895° E (the latter longitude refers to the centre of the tracer grid box). The respective restoring timescale is prescribed as a linear function of longitude, starting with 1 h at the western boundary and increasing to 12 h at 6.895° E.

We do not restore directly to observations of sea surface height because this would limit the application of the model: coupled ocean atmosphere earth system models are not (yet) up to the task of providing reliable sea surface height projections at the entrance to the Baltic Sea. Hence, if we would restore directly to sea surface height we would be constrained to the past where we have observations. Instead, we restore to SSH derived from a proxy which can reliably be simulated. Specifically we restore to SSH as calculated by a regression to the atmospheric sea level pressure gradient between Oslo in Norway and Szczecin in Poland. The ratio behind the approach is the high correlation between the normalised atmospheric pressure gradient, dubbed Baltic

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea Index (BSI), and the exchange between the Baltic and the North Sea observed by Lehmann et al. (2002).

We restore to SSH (η) calculated as:

$$\eta = (333.393 \cdot \text{BSI} + 17.225) \times 10^{-3} \quad (2)$$

The respective coefficients are from a regression of monthly mean PSMSL (PSML, 2012) tide gauge data against the BSI (A. Lehman, personal communication, 2013). The BSI used here is identical to the one presented in Lehmann et al. (2002).

2.4 River runoff

The river runoff applied in this study is a crossover from observations (Bergström and Carlsson, 1994) and a large-scale hydrological model (Graham, 2004). It is identical to the one applied in Meier and Faxen (2002), Meier and Kauker (2003), Hordoir and Meier (2010), and Löptien and Meier (2011). Figure 3 shows the total discharge to the model domain. We resolve 30 discharge locations (rivers and groundwater inflows).

In high-resolution Baltic Sea configurations, river runoff is prone to cause numerical problems because the salinity is close to zero in the vicinity of some rivers. This can, at times, due to the dispersive nature inherent to all state-of-the-art advection schemes (other than the out-dated upstream scheme) induce negative salinities for which the equation of state is not defined. We solved this problem by distributing the runoff over the 6 grid boxes closest to the actual position of the river mouth. The distribution among the 6 grid boxes is determined by weights (W_i),

$$W_i = \frac{\exp(-D_i\tau^{-1})}{\sum_{i=1}^6 \exp(-D_i\tau^{-1})}, \quad (3)$$

calculated with an exponential decay length scale $\tau = 3$ km as a function of the distance of respective grid boxes to the actual river mouth (D_i).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.5 Air-sea fluxes

The atmospheric boundary conditions driving our ocean-ice model are identical to those applied to the “RCO” ocean-ice model in Meier and Faxen (2002), Meier and Kauker (2003), Hordoir and Meier (2010), and Löptien and Meier (2011). They are based on downscaled ERA-40 reanalysis (Uppala et al., 2005, and many more to follow) which, compared to the original ERA-40 reanalysis, features additional regional details which considerably effect the solution of standalone ocean models of the Baltic Sea (Meier et al., 2011). The downscaling procedure takes ERA-40 reanalysis data as boundary conditions for the Rossby Centre Regional Atmosphere model version 3 (hereafter RCA3) which features an enhanced (relative to ERA-40) horizontal resolution of 25 km (Jones et al., 2004; Samuelsson et al., 2011). As shown by Samuelsson et al. (2011), the approach provides a very realistic climatology. A shortcoming, however, is that RCA3 does not employ techniques such as spectral nudging (von Storch et al., 2000) which constrain simulated cyclone paths to match actual tracks. This is of some concern because a systematic analysis of cyclone tracks (e.g. similar to analysis carried out by Löptien et al., 2008) simulated with RCA3 has not been carried out yet.

We use cloud cover, precipitation, near-surface humidity, pressure, temperature and wind speed from the RCA3 model at a temporal resolution of 3 h as provided by the Rossby Centre at SMHI, Norrköping, Sweden.

Solar shortwave radiation entering the ocean is calculated with a solar constant of 1353Wm^{-2} , an atmospheric turbidity of 0.95 and cloud cover as described by Meier et al. (1999). All shortwave radiation is absorbed in the uppermost wet grid box which has a nominal thickness of 3 m (Sect. 2.1). This is a pragmatic decision because a prognostic model of water turbidity is yet to be developed. Note that this induces an error since some of the incoming radiation reaches below the surface mixed layer depth under certain conditions (e.g., Löptien and Meier, 2011).

Long wave radiation entering the ocean ($\text{LWRAD}_{\text{down}}$) is calculated from cloud cover as in the RCO model used by Meier and Faxen (2002), Meier and Kauker (2003),

GMDD

7, 2065–2124, 2014

MOMBA 1.1

H. Dietze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hordoir and Meier (2010), and Löptien and Meier (2011). The formulation is unusual since it does neither contain the water vapour pressure nor the dew point temperature. However, given the high uncertainty of these empirical relationships which generally exceeds 10Wm^{-2} (e.g., Zapadka et al., 2007) we use it, nevertheless:

$$5 \quad \text{LWRAD}_{\text{down}} = \sigma T_{\text{air}}^4 (0.7829(1 + 0.2232 \cdot \text{CLCO}^{2.75})), \quad (4)$$

where $\sigma = 5.6703 \times 10^{-8} \text{Wm}^{-2} \text{K}^4$ is the Stefan–Boltzmann Constant, T_{air} is the air temperature in 2 m height, and CLCO is the cloud cover expressed as a number between 0 and 1.

10 Long wave radiation leaving the ocean is calculated following Large and Yeager (2004) assuming an emissivity of 1.

The turbulent air–sea fluxes, i.e., the wind-stress, the evaporation, the sensible and the latent heat flux are calculated with bulk formulas according to Eqs. (4a–d) in Large and Yeager (2004), respectively. We account for ocean currents in the bulk parameterisation of the wind stress which significantly affects the model solution as explained in the forthcoming Sect. 3.2.

15 The calculation of ocean–atmosphere fluxes which are actually applied to the ocean every time step is updated (based on linearly interpolated 3 hourly atmospheric boundary conditions described above and the contemporary ocean state) every 36 min, and kept constant in-between.

20 We do account for the effect of atmospheric bottom pressure gradients on ocean model pressure gradients.

3 Results

25 To serve as a nucleus of a fully coupled biogeochemical circulation model of the Baltic Sea, the ocean circulation module has to perform with a certain level of quality. To this end we present a choice of model assessments:

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Surface circulation (Sects. 3.1 and 3.2) which, e.g., effects the wind-driven upwelling of nutrients (as we will put forward) and subsequent pathways of nutrients.
- Sea surface temperature (SST, Sect. 3.3) which can be observed from space and, therefore, is available in an unrivalled (compared to in-situ measured properties) spacial and temporal resolution. SST serves as a proxy for the realism of the surface mixed layer depth whose dynamics is a major process involved in the supply of nutrients from depth to the sun-lit surface ocean.
- Inflow of saline oxygen-rich waters from the North Sea to the Baltic and its subsequent spread at depth (Sect. 3.4) which a major agent resupplying the Baltic Sea with oxygen at depth and, in its turn, determines if organic matter sinking to depth is denitrified rather than being remineralized under oxic conditions.
- Halocline depth (Sect. 3.5) which is a bottleneck for the diffusive supply of nutrients (oxygen) to the surface (abyss).
- Sea ice (Sect. 3.6) which caps the ocean from the atmosphere throughout almost half of the time up North and thereby controls the air–sea gas exchange of climate-relevant species such as carbon and N_2O .

We close this section with a discussion of technical issues such as grid scale noise (Sect. 3.7), computational uncertainty (Sect. 3.8) and cost (Sect. 3.9).

3.1 Average surface circulation

The circulation in the Baltic Sea is driven by the wind exerting stress at the sea surface, sea surface tilt effected by e.g. river runoff, thermohaline horizontal gradients of density and tidal forces. Overall, this results in a predominantly wind-driven, weak, cyclonic circulation with surface speeds of some 5 cm s^{-1} which, at places, are quite persistent (Leppäranta and Myrberg, 2009; Sjöberg, 1992). To this end our simulation is consistent with the observations as is suggested by Fig. 4. Unfortunately a more quantitative

model assessment is hindered by the lack of current measurements available to us, and thus we are restricted to a comparison with sea surface salinity and other model simulations.

Sea surface salinity (SSS) is determined by the surface circulation mixing river runoff and salty water of North Sea origin. Hence, as put forward by Eilola and Stigebrandt (1998) and exploited by Lehmann et al. (2002), a realistic SSS is a proxy for a realistic surface circulation. Figure 5 shows a comparison between simulated and observed SSS which suggest that the simulated surface circulation is realistic.

As concerns other simulations we find that patterns of the magnitudes of mean surface velocities are comparable, although weaker in amplitude, with recent results from Jedrasik et al. (2008) based on a model that covers the entire Baltic. The most pronounced features in our simulation which are different to Jedrasik et al. (2008) are: (1) lower velocities in the Archipelago Sea east of Åland, (2) a distinct band of high velocities in the north eastern Sea of Bothnia west of Ulvö Deep, and (3) a distinct cyclonic gyre southeast of Gotland.

1. As regards the Archipelago Sea it is save to assume that both models are deficient because their spacial resolution does not allow to resolve the multitude of islands east of Åland (Fig. 1d).
2. As regards west of Ulvö Deep our model is consistent with the simulation of the Gulf of Bothnia by Myrberg and Andrejev (2006) which also features high southwards velocities. Furthermore these authors report, consistent with our results, a high persistency R of these currents.

The persistency R is defined as the ratio between vector and scalar mean speeds:

$$R = \frac{|\langle \mathbf{U} \rangle|}{\langle U \rangle}, \quad (5)$$

where vector $\langle \mathbf{U} \rangle$ is mean vector speed and $\langle U \rangle$ the mean scalar speed. Figure 4 shows a map of simulated R : in summary and consistent with results from Jedrasik et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(2008), Lehmann et al. (2002), Myrberg and Andrejev (2006), and Andrejev et al. (2004) we find currents of a persistency exceeding 80 % along closely packed isobaths (Fig. 4) – a feature that can be described in terms of a depth integrated vorticity balance (e.g., Lehmann et al., 2002).

3.2 Transient surface circulation

Even though the mean circulation can provide insight into tracer dynamics (such as, e.g., transports of salinity, pollutants and nutrients from river mouths), the concept is limited by the highly transient nature of the Baltic Sea circulation. Figure 6 shows a measure of this transiency, the simulated eddy kinetic energy (E) defined as:

$$E = 0.5 \overline{(u'^2 + v'^2)}, \quad (6)$$

where u' and v' are deviations of the zonal and meridional surface velocities from their time mean over the 1987 to 1999 period, respectively. The overbar denotes a temporal average over the same period and, u' and v' are calculated from daily means. Typical eddy kinetic energies are around $50 \text{ cm}^2 \text{ s}^{-1}$ with hotspots exceeding $200 \text{ cm}^2 \text{ s}^{-1}$. For comparison, this is an order of magnitude less than the up to $2000 \text{ cm}^2 \text{ s}^{-1}$ found, e.g., off Japan where the Kuroshio separates from the coast; but still substantial when compared with typical open ocean conditions away from worlds' most powerful boundary currents (Dietze and Kriest, 2012; Liu et al., 2010).

A comparison between persistency (Fig. 4) and eddy kinetic energy (Fig. 6) suggests, however, that the term eddy kinetic energy is somewhat misleading because some regions of high eddy kinetic energy are characterised by high current persistency (e.g., west of Ulvö Deep). This is apparently inconsistent with the generic concept of whirling eddies. The solution to this conundrum is that eddy kinetic energy (Eq. 6) can be effected by intermittent *unidirectional* currents which score a high persistency (Eq. 6). Hence, our model predicts that some of the regions which host the most powerful surface current variability are characterised by intermittent but unidirectional currents which are steered along isobaths (Figs. 6 and 4).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

As concerns the small scale transient eddying circulation we report, in the remainder of this subsection, on a subtlety, the so-called eddy/wind effects (Martin and Richards, 2001): our model configuration explicitly includes the effect of the ocean surface circulation on the stress exerted by the winds on the sea surface. To this end our configuration is more realistic than those studies who neglect this effect (e.g., Jedrasik et al., 2008, to name one of the very few studies who document this aspect of their wind-forcing). The eddy/wind effect introduces a non-linearity to the dynamics, since the wind stress is a function of the difference between the wind speed and the movement of water at the surface, which in turn is determined by the wind stress. Recent open ocean studies suggested that this non-linear effect is significant (e.g., Duhaut and Straub, 2006; Zhai and Greatbatch, 2007; Xu and Scott, 2008) even though surface currents are typically an order of magnitude smaller than the winds. The reason is that the spacial scales of transient circulation are much smaller in the ocean than in the atmosphere (which is a consequence of the stronger stratification in the atmosphere resulting in larger Rossby radii). Thus, even though small in magnitude, ocean currents can significantly alter the curl of the wind stress (which is a function of spacial derivatives of stress) exerted on the sea surface. As put forward by Martin and Richards (2001) this causes vertical Ekman velocities w_e ,

$$w_e = -\text{curl} \left(\frac{\mathbf{T}}{\rho f} \right) \quad (7)$$

(where \mathbf{T} is the vector wind stress, ρ the density of water and f the Coriolis parameter) of the order of $\approx 0.5 \text{ m day}^{-1}$ in typical open-ocean eddies. The associated nutrient transports from depth to the sun-lit surface ocean can be substantial (Martin and Richards, 2001; McGillicuddy et al., 2007; Ledwell et al., 2008; Eden and Dietze, 2009). It is straightforward to assume that this holds, even more so, in the Baltic Sea where shallow water depths reduce the Rossby radii to a couple of kilometres (compared to the several tens of km typical for open ocean conditions). Figure 7, indeed, supports this hypothesis: Ekman upwelling (downwelling) w_e calculated by taking into

account the surface velocity in the calculation of the wind stress, is correlated with small-scale anticyclonic (cyclonic) features. Since these features are smaller than the resolution of the atmospheric forcing they have to be significantly influenced by the surface circulation modulating the wind stress exerted on the ocean's surface. Further we find, consistently with the Rossby radii being an order magnitude smaller than w_e is of an amplitude of several meters – almost an order of magnitude larger than the $\approx 0.5 \text{ m day}^{-1}$ observed in the open ocean by, e.g., Ledwell et al. (2008).

3.3 Sea surface temperature

We use the KPP boundary layer parameterization (Large et al., 1994) with the same parameters applied in eddy-permitting global configurations of Dietze and Krist (2012), Dietze and Löptien (2013), and Liu et al. (2010). Remarkably, the simulated SSTs are very realistic which is a real asset of MOMBA. Figure 8 shows a comparison with SHARK (Swedish Oceanographic Data Centre at SMHI) data which suggest, that the only flaw of the model is, at times, an underestimation of maximum summer temperatures. We speculate that this is the consequence of:

1. the vertical discretisation which assigns a nominal (c.f. Eq. 1) thickness of 3 m to the uppermost grid-box (Fig. 2) which (implicitly) imposes a lower bound of 3 m on our simulated surface mixed layer depths. This, however, can be unrealistically high during low-wind summer conditions.
2. The neglected effects of phytoplankton blooms which can shift the absorption of solar radiation in the water column upwards which, in turn, increases the SST (e.g. Löptien et al., 2009; Löptien and Meier, 2011).

Observations from space confirm the extraordinary quality of our SST hindcast: Fig. 9 shows a comparison with weekly composites of quality controlled Advanced Very High Resolution Radiometer data from NOAA and the european MetOp satellites distributed by the Bundesamt für Seeschiff und Hydrographie, Germany via

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



http://www.bsh.de/de/Meeresdaten/Beobachtungen/Fernerkundung. The data is referred to as BSH SST data in the following. For the following model-data comparison we set all BSH SST data below -0.5 to -0.5°C in order to prevent for accidentally comparing simulated SSTs with observed sea ice temperatures. Typically, we find that the model explains more than 90 % of the variance in the BSH SST data. Exceptions with lower correlations (which will be discussed in the following) are the (1) Skagerrak, (2) the Bothnian Bay, (3) the Archipelago Sea east of Åland and (4) those regions hosting frequent upwelling (Lehmann et al., 2012) such as the Swedish south and east coasts, south of Gotland and off Finland in the Gulf of Finland:

1. we speculate that simulated temperatures in the Skagerrak are deteriorated by restoring to an annual mean climatology (Sect. 2.2).
2. The reason for the low correlations in the Bothnian Bay is probably associated with sea ice melting too late in spring (Sect. 3.6).
3. The low correlation in the Archipelago Sea is not surprising given the unrealistic land-water distribution owed to scant resolution (Fig. 1d).
4. The low correlations in the upwelling regions are not necessarily indicative of a poor representation of associated processes. To the contrary, visual inspections of SST maps such as shown in Fig. 10 reveals striking similarities between the model and the BSH SST. However, on a grid-point to grid-point basis there is a mismatch. First, because we compare weekly composites (assembled during cloud free periods) with simulated daily averages of upwelling events some of which have a lifetime of a few days only. Second, upwelling is highly non-linear (also because of the subtle circulation/wind effects described at the end of Sect. 3.2). This non-linearity makes it difficult (or even impossible) to retrace the dynamics of single upwelling events on a one-to-one basis.

GMDD

7, 2065–2124, 2014

MOMBA 1.1

H. Dietze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.4 Deepwater renewal

The assimilation of inorganic carbon by algae in the sun-lit surface ocean and its subsequent export as sinking particulate organic carbon fuels oxygen consumption at depth. This consumption is balanced by resupply of oxygen-rich waters of North Sea origin which enter the Baltic intermittently via the Danish Straits. When this resupply falls short of the consumption, the oxygen concentrations can be depleted below certain thresholds. In turn, denitrification and the release of dissolved organic phosphorous bound to feoxyhydroxides from the sediments may set in. This is (although discussed controversially, Pahlow et al., 2013) of concern, because the associated change in the N/P ratio is thought to promote the growth of potentially harmful cyanobacteria (e.g., Vahtera et al., 2007).

Unfortunately, MOMBA features a spuriously hampered deepwater renewal: Fig. 11 shows a comparison between simulated and observed near-bottom salinities at stations BY2, BY5 and BY15. The further into the Baltic, away from the Danish Straits, the weaker the simulated salinity signature of the inflow events. This holds especially for the massive 1993 event which is not reproduced at all at station BY15.

The underestimation of deep salinities is an infamous feature of z-level models which do not apply data assimilation techniques (e.g., Fu et al., 2012). However, the especially bad performance of MOMBA came as a surprise because of the model's the high and competitive spacial resolution and its – as suggested in the following sea surface height analysis – realistic exchange of water through the Danish Straits:

Sea surface height variations are effected by, e.g, (1) variations of air pressure, wind and density in the Baltic, (2) internal eigenoscillations and standing waves (so called seiches), and (3) “external” forcing by varying sea level outside the Baltic (and, to a lesser extent, by freshwater supply to the Baltic). The relative contribution of these mechanisms depends on the actual location and the frequencies considered, also because the seiches have distinct spatial characteristics both in space and in the frequency domain. To this end the area around Stockholm/Landsort is special, because it is situated

GMDD

7, 2065–2124, 2014

MOMBA 1.1

H. Dietze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

at the node of major seiches, and therefore sea level readings are rather unaffected by internal oscillations. Consistently, Samuelsson and Stigebrandt (1996) report that the variability at the Stockholm/Landsort gauges is highly correlated with “external” forcing. For this reason, a realistic simulation of the sea surface height at Stockholm/Landsort is generally understood as a proof of a realistic water exchange through the Danish Straits (e.g., Neumann and Schernewski, 2008).

Figure 12 shows the simulated daily mean sea surface height along with daily mean observations (which comprises the world’s longest continued series of sea level observations Ekman, 1988; Woodworth and Player, 2003; PSML, 2012). Specifically we use data retrieved on 5 June 2013 from the Permanent Service for Mean Sea Level at <http://www.psml.org/data/obtaining/> (Holgate et al., 2013). Expressed in terms of a correlation, we reach 0.78 and 0.85 at Stockholm and Lansort, respectively. The number of observations is 4748 in both cases. Compared to previous studies which typically report correlations higher than 0.9 (e.g. Neumann and Schernewski, 2008; Jedrasik et al., 2008) this is on the lower end of published models, but not extraordinarily poor. To this end it is noteworthy, that our rather low correlations are not necessarily indicative of a poor model dynamics, but may well be preliminarily caused by our boundary conditions: in contrast to other studies we do not restore to observed sea levels in the Kattegat/Skagerrak but prescribe sea surface height derived from a correlation with a normalised atmospheric pressure gradient (Sect. 2.3) further west, at the edge of our model domain. Our approach is adopted from Lehmann et al. (2002) who report a *weekly* (monthly) correlation of 0.83 (0.82) of their simulation with weekly (monthly) sea level data from Landsort. Compared to their performance we reach a competitive 0.86 (0.88) calculated based on weekly (monthly) data at Landsort.

In line with the results from the sea surface height analysis, a more direct comparison with an observational estimate from the 1993 inflow in Table 2 confirms that the simulated transport through the Danish Straits is, indeed, realistic. As concerns the associated salinity transport it is save to assume that it is not biased low because we restore salinity to a rather high (and unrealistic) 35 PSU in the North Sea (Sect. 2.2).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Overall, the combination of a rather high salinity transport to the Baltic and its missing salinity signatures at depth leaves us with the question where the salty and dense inflow gets lost on its way to the Baltic Proper. It is straightforward to assume that the problem of an apparently hampered deepwater renewal is associated with our choice of the bottom boundary layer parameterization (Döscher and Beckmann, 2000). However, excessive tuning exercises (not shown) suggested that this is not the case which is, maybe, not so surprising because Fig. 13a suggests that most of the topographic slopes are actually well resolved, which in turn, should prevent for much of the infamous mixing at sills in Boussinesq-approximated geopotential coordinate models.

Analysis based on temporal high resolution output finally revealed the cause of the spurious dilution of salty inflows and their subsequent absence in the deeper basins: Fig. 14 shows a section along the Hamare Strait (as indicated in the inlayed map) which is typical for the situation during strong inflows at the sills in MOMBA. On timescales of hours and spacial scales of several nautical miles, vertical circulation cells with alternating up- and downward velocities develop during stronger inflows. The magnitude of these circulation patterns is of the order of several tens of meters per day and they intermittently cross the surface mixed layer. Thus, the upward branches of these patterns transport deep saline bottom water to the surface (and, likewise, lighter surface waters to the bottom) where it is spread horizontally by the prevailing turbulent circulation (Fig. 6). This can cause vertical instabilities as dense bottom water is shifted horizontally over lighter subsurface waters which, in turn, drives further vertical mixing. The overall effect is one of enhanced vertical mixing at the sills during strong inflow events. The salt associated to the inflow events is mixed upwards to the surface and eventually flushed out of the Baltic by the net positive freshwater supply rather than reaching the deeper basins further away from the Danish Straits.

We speculate that the vertical circulation cells during strong inflows at the sills (as the one shown in Fig. 14 for the Hamare Strait) are numerical artefacts caused by the divergence of horizontal currents interacting with the step-like representation of topography. At the least, even if effects like that exist (although we have found no supporting

evidence in the literature) their mixing effect is obviously unrealistically strong in our model. Further, we speculate that the problem is more pronounced in high-resolution configurations like ours, because the spacial scale of up and downwards jets is several grid-boxes only (Fig. 14). Previous Bousinesq-approximated configurations on geopotential coordinates with resolutions of several nautical miles might just have been lucky enough to not “resolve” these spurious cells.

The remaining question is how to dampen these cells. To this end we performed an extensive set of tuning experiments including modified winds, surface mixed layer depth and bottom boundary layer parameterizations, initial conditions, all branching off the reference simulation on 1 January 1993. As regards improved realism, all were to no avail. Figure 15 shows the most promising approaches: increased viscosity dampens the spurious circulation cells. Hence, higher viscosity reduces the spurious mixing and comes along with increased simulated near-bottom salinities at station BY5. When the viscosities get too high, however, the saline inflows are decelerated, and even less salt finds its way to the deep Baltic. Note, that the apparently optimal solution with an added vertical viscosity of $10^{-2} \text{ m}^2 \text{ s}^{-1}$ which reaches realistic values in January 1993 at station BY5 in Fig. 15b, features an overall worse performance than the reference simulation. Similar behaviour, actually, applied to all our tuning experiments: while quite some of them reduced local misfits to observations none of them achieved global improvement.

We conclude that spuriously damped deepwater renewal is endemic to MOMBA. The underlying mechanism is apparently related to the high spacial resolution, a finding which is consistent with results from Ezer and Mellor (2004) and tuning exercises with the BaltiX model configuration (Hordoir et al., 2013): R. Hordoir, personal communication (2012) reports that increasing viscosity at the surface only, increases near-bottom salinities in the deep Baltic basins up to realistic values leading to an improved global performance. We speculate that this is the case because the higher near-surface viscosity dampens that part of the spurious circulation pattern which effectively mixes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



across the surface mixed layer and, at the same time, does not slow down the near-bottom overflows which transport the salt from the Danish Straits far into the Baltic.

Additional and pragmatic remedies other than applying non-homogenous viscosities could be adopted from previous efforts to model the Denmark strait overflow: Kösters et al. (2005), e.g., parameterize the transport across the (Denmark Strait) sill as a function of the large-scale conditions and then – directly – impose this transport thereby overriding the effects of resolved spurious circulation on the sill. It is noteworthy that their improved and apparently more realistic model features significantly differing climate projections (Köller et al., 2010). We anticipate a similar behaviour in our case and conclude that a comprehensive study on this subject should preclude any attempts to make climate projections with the current modeling framework.

3.5 Halocline

A permanent halocline which separates the dense saline near-bottom waters of North Sea origin from the surface layer which comprises fresh river runoff, is a characteristic feature of the Baltic. Its depth is determined by the balance between inflowing saline waters and its subsequent diapycnal upwards transport (via upwelling and mixing processes). Thus, a realistic simulation of the halocline depth is a necessary precondition for simulating vertical nutrient and oxygen fluxes.

The simulated median salinity profile at station AE (Fig. 1a) in the Kattegat is of a shape very similar to the observed one (Fig. 16a). However, there is an offset, probably effected by the rather high (and unrealistic) salinity value to which we restore in the North Sea (35 PSU, Sect. 2.2). Further into the Baltic, away from the Danish Straits, e.g., at station BY5 (Fig. 16b) and BY15 (Fig. 16c) salinities are no longer biased high. To the contrary, as outlined in Sect. 3.4, near-bottom salinities are low because of spurious vertical circulation patterns at the sills during major inflow events. Even so, we conclude, based on Fig. 16 (and corresponding analysis at all other stations indicated in Fig. 1 which we omit here for the sake of brevity), that the transition from the surface mixed layer to the stratified halocline is simulated realistically (i.e., within ≈ 10 m of the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



observations). This puts confidence in the simulated diapycnal transport mechanisms at the base of the surface mixed layer which is an important asset of the model since the associated nutrient fluxes drive primary production and subsequent respiration at depth (e.g., Dietze et al., 2004).

3.6 Sea ice

The Baltic Sea is seasonally ice covered. To first order approximation this cover acts as a lid, hindering the exchange of heat and moisture between the ocean and the atmosphere, thereby essentially decoupling the systems. Further, it prevents for the exchange of climate relevant trace gases such as CO₂ and N₂O. To this end sea ice dynamics is of interest, maybe even more so in a warming world of ever retreating ice cover.

Here we compare our model results with digitised ice charts from the operational ice service provided by SMHI. The charts are based on information from ship crews, monitoring stations blended and complemented with measurements from space. The major consumer of this information has been commercial shipping and it remains to be shown that they provide a realistic ground truth to which models can be compared. That said, we assume in the following that the operational ice charts are, to date, the most reliable large-scale source of information for the respective time-period.

Figure 17 shows that the simulated ice extend is biased high compared to the observations. This spurious feature is further highlighted by exemplary snapshots for particularly warm and cold winter conditions in mid February (typically the time of maximum extend, Leppäranta and Myrberg, 2009) in Fig. 18.

Figure 17 suggest that, although biased high, the simulated inter-annual variability of the ice extent is realistic. This is confirmed by an empirical orthogonal function (EOF) analysis of both simulated and observed mean seasonal mean ice concentrations: The leading EOF explains 71 % (68 %) of the variability of observed (simulated) ice concentrations and thus covers the major signal. The corresponding leading principle components (PCs) are almost identical (correlation: 0.92) and show (Fig. 19), furthermore,

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a strong correlation (0.8) with the PC-based North Atlantic Oscillation (NAO) index (as applied, e.g., in Hurrell, 1995; Hurrell and Deser, 2009; Löptien and Ruprecht, 2005). This is in agreement with published state-of-the-art model studies (e.g., Tinz, 1996; Jevrejeva and Moore, 2001; Jevrejeva et al., 2003; Koslowski and Loewe, 1994; Löptien et al., 2013).

As regards spacial patterns of variability, the similarity between the leading EOFs of the simulation and the observations in Fig. 20 suggests, that the underlying dynamics which drive the ice are – overall – realistically reproduced by the model. However, as a consequence of the model’s notorious overestimation of ice concentrations the corresponding EOF-pattern are more pronounced in the simulation. This holds particularly in the Bothnian Sea.

Additional model deficiencies are apparent in Fig. 20 which shows spuriously reversed phase relations off Germany and up north in Bothnian Bay and the Bothnian Sea. Off Germany this is the consequence of the simulated bias in ice extent. As for up north, we speculate that the spuriously reversed sign in the EOF pattern is related to unresolved wind-ice interaction: cold winters are often related to persistent northerly winds which can cause frequent leads (Löptien et al., 2013). The reproduction of these areas of open water is, probably, hindered by model’s spacial resolution (even though we run on a competitive 1 nm).

3.7 Grid scale noise

Ocean circulation models which use an Arakawa B-grid spacial discretisation (Arakawa and Lamp, 1977) (like MOMBA) are prone to catch numerically induced grid scale noise in the simulated velocity field and sea surface height (Mesinger, 1973; Killworth et al., 1991; Pacanowski and Griffies, 1999; Griffies et al., 2001). These so-called checkerboards are especially pronounced in coarsely represented enclosed embayments (Griffies, 2009) when the applied viscosity is rather small (e.g., Jochum, 2008). Given that, (1) the Baltic Sea is almost enclosed, (2) our spacial discretisation is still rather coarse even though we have a resolution of 1 nm (compare Fig. 1b, c and d) and,

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(3) that we aim too keep the explicit viscosity as low as possible such that we have no unrealistic damping of the circulation, special attention was dedicated to the checkerboard problem. Especially because Löptien and Meier (2011) report checkerboards in simulated temperature in an Arakawa B-grid model of the Baltic Sea with a horizontal resolution of 2 nm.

We use the dissipative predictor-corrector barotropic time stepping scheme with a dissipation parameter of 0.2. In addition we smooth the sea surface height spatially (1) at each barotropic time step with a Laplacian operator and (2) at each longer, baroclinic, time step with a more scale selective biharmonic operator. In both cases (i.e, the Laplacian and the biharmonic operation) the associated parameter is a velocity of 0.5 cm s^{-1} . The applied laplacian (biharmonic) diffusivity is then calculated by multiplication with the (cube of the) harmonic mean of the zonal and meridional grid length. Details of the sea surface height filtering (and, consistently to the tracer equation) applied in MOMBA are described in Sect. 8.3 of Griffies (2009). Visual inspection of snapshots of all prognostic variables in MOMBA revealed no evidence of checkerboard wheresoever.

3.8 Computational uncertainty

Model integrations are not necessarily reproducible. This applies when switching from one hardware to another, changing the compiler, or changing compiler switches. Among the reasons is, that round-off errors generally violate commutative laws. Hence, the order of execution – which is tailored by the compiler for optimal performance on the respective hardware – can influence the results. Note that reproducibility is not even guaranteed if the identical executable is executed on the identical hardware (as is the case with aggressive floating point optimizations set by Intel’s -fp-model compiler option).

Reproducibility issues with numerical integrations of highly nonlinear systems have been reported elsewhere (e.g., Song et al., 2012) and can be disturbing: Fig. 21 shows results from two “identical” integrations applying the default (but rather aggressive) Intel

-fp-model compiler option. The difference in surface properties seem to be insignificant (compare panel (a) with panel (b) in Fig. 21). However, the effects of the highly nonlinear vertical circulation patterns above the sills during inflow events (Sect. 3.4) amplify to a stunning difference in near-bottom salinities at station BY5 in early 1993 (panel (c) and (d) in Fig. 21).

3.9 Computational cost

MOM4 runs on a wide range of machines. E.g., we successfully tested coarser configurations on a 8-core MacPro (Intel Xeon, 2.5 GHz, 12 MB L3 cache, 16 GB RAM) before we switched to higher performing hardware.

The configuration presented in this study was benchmarked on three different high performance computing clusters (HPCs), all of them interconnected with InfiniBand. The respective underlying computational nodes were (1) Intel Xeon Harpertown E5472, 8 core, (2) Intel Xeon Gainestown X5570, 8 core, and (3) Intel Xeon Sandybridge, 2 × 8 core. In line with Schmidt (2007) we find that the code is suitable for parallel processing up to several hundred tasks (Fig. 22). Unfortunately, however, we found comparably bad scalability on our main workhorse, the Xeon Sandybridge (RZ-Kiel/GEOMAR): as shown in Fig. 22 performance drops dramatically at around 100 cores, probably as a consequence of an IO bottleneck during initialisation.

In addition to tests on HPCs we ran preliminary benchmarks (albeit running the MOM5 kernel which may, or may not be comparable) on a contemporary inexpensive (\approx EUR10 000,-) system consisting of two 32-core workstations, based on 6320 AMD Opteron (Abu-Dhabi) (8-Core CPU, 2.80 GHz, 16MB L3 Cache, DDR3 1600) and interconnected with a QDR InfiniBand. Wall clock times for the integration of one model year are less than 4 days and the associated cost (write off and power consumption) are below EUR100,- (based on a machine lifecycle of three years). If processing speed and cost will remain to be linked to Moore's law, we expect similar cost for a fully coupled biogeochemical ocean circulation model in two years time from now.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Summary and conclusions

We set out to develop a competitive Baltic Sea ocean-circulation model configuration, dubbed MOMBA, which is intended to serve as a nucleus of a coupled ocean-circulation biogeochemical model. To this end our main focus was a realistic simulation of surface mixed layer dynamics and diapycnal near-surface transport processes because these processes are key controls on pelagic biogeochemistry (e.g., Sverdrup, 1953; Eppley and Peterson, 1979, respectively).

MOMBA features a horizontal resolution of ≈ 1 nautical mile and a total of 47 vertical layers. It is based on a recent release of GFDL's modular ocean model. The computational cost associated to integrating one year is substantial but, yet, decadal or even centennial are computationally feasible: expressed in terms of wall clock time the cost ranges, depending on the hardware ("low cost" or high performance computing) from 4 days to several hours.

The high spacial resolution of MOMBA permits the resolution of eddies. Overall, we find an impressively eddying surface circulation. The associated eddy kinetic energy is, locally, even comparable to open ocean conditions which showcases the highly non-linear characteristics of the circulation field in the Baltic Sea. As a peculiarity we find strong evidence that this nonlinearity is further amplified by eddy/wind effects which modulate the wind-induced up- and downwelling. Rough scaling suggests that these eddy/wind effects may well be among the major processes transporting deep nutrient rich waters to the surface.

Despite the overall transient characteristics of the flow field we simulate local, highly persistent circulation features which are correlated with topographic slopes. If this topographic steering of surface currents is indeed as dominant as indicated by the model is, however, yet to be investigated.

A real asset of MOMBA is its realistic simulation of sea surface temperature which we interpret as indicative of a realistic simulation of surface mixed layer dynamics. Additional evidence indicating that the simulation of near-surface diabatic processes

GMDD

7, 2065–2124, 2014

MOMBA 1.1

H. Dietze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in general, is realistic is deduced from the model's realistic halocline depths. Overall, this suggests that the simulated physics of MOMBA is of a sufficient fidelity so as to provide a sound base for research focused on environmental near-surface pollution (such as in, e.g., Lehmann et al., 2014a), or on biogeochemical questions such as, e.g., the controls of cyanobacteria (which are discussed controvesially Pahlow et al., 2013; Ward et al., 2013).

Caveats are (1) the simulated sea ice extent, which is reasonable as concerns temporal and spacial variability, but is apparently biased high relative to observations, and (2) the failure of MOMBA to reproduce major inflow events. The latter problem is associated with spurious vertical circulation cells developing at the sills during strong inflows. These cells effectively dilute the saline deep inflows with lighter surface waters and thereby inhibit deep water renewal in the deep Baltic Basins. A remedy for this model artefact, which is probably a consequence of the high spacial resolution, is still to be developed. We suggest to investigate approaches, similar to the one developed by (Kösters et al., 2005) for the Denmark straight overflow.

5 Model code and data repositories

We use the original code MOM4p1 as distributed by NOAA's Geophysical Fluid Dynamics Laboratory (<http://www.gfdl.noaa.gov/fms>) except for in the first year of spin-up, where we use a modified version of `ocean_tracer.F90` which caps outlying temperatures and salinities to enforce numerical stability. The model configuration described here, including respective namelists, the forcing, initial conditions and the modified `ocean_tracer.F90` are accessible via <http://www.baltic-ocean.org>. In addition we distribute a modified version of `memutils.F90` which runs on OS X 10.8.5.

Acknowledgements. Discussions with L. Axell, C. Dieterich, A. Hoeglund, R. Hordoir and G. Väli are appreciated. H. and U. thank A. Oschlies for essential long-term support. Integrations were performed at the Norddeutscher Verbund für Hoch- und Höchstleistungsrechnen, and at the Rechenzentrum at Kiel University, Germany. We thank the Rossby Center at SMHI,

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Norrköping, Sweden for sharing their atmospheric re-analysis which drives our ocean model. The International Council for the Exploration of the Sea (ICES) is acknowledged (along with all data contributors) for archiving and distributing temperature and salinity data. We thank the Bundesamt für Seeschifffahrt und Hydrographie, Germany for the distribution of satellite SST, and SMHI for in-situ temperature and salinity observations. The “Permanent Service for Mean Sea Level” is acknowledged for providing tide gauge data. The ice charts were provided by SMHI. The NAO-index is from from Hurrell, James, National Center for Atmospheric Research Staff, as of 8 October 2013. This work is a contribution of the Sonderforschungsbereich 754 “Climate-Biogeochemistry Interactions in the Tropical Ocean” (www.sfb754.de), which is funded by the German Science Foundation (DFG).

The service charges for this open access publication have been covered by a Research Centre of the Helmholtz Association.

References

- Adcroft, A. and Campin, J.-M.: Rescaled height coordinates for accurate representation of free-surface flows in ocean circulation models, *Ocean Model.*, 7, 269–284, 2004. 2069
- Anderson, T.: Plankton functional type modelling: running before we can walk?, *J. Plankton Res.*, 27, 1073–1081, 2005. 2067
- Andrejev, O., Myrberg, K., Alenius, P., and Lundberg, P. A.: Mean circulation and water exchange in the Gulf of Finland – a study based on three-dimensional modelling, *Boreal Environ. Res.*, 9, 1–16, 2004. 2077
- Arakawa, A. and Lamb, V. R.: Computational design of the basic dynamical processes of the UCLA general circulation model, in: *Methods in Computational Physics*, edited by: Chang, J., Vol. 17, Academic Press, New York, 173–265, 1977. 2068, 2087
- Arhonditsis, G. and Brett, M.: Evaluation of the current state of mechanistic aquatic biogeochemical modeling, *Mar. Ecol. Prog.-Ser.*, 271, 13–26, 2004. 2067
- Bergström, S. and Carlsson, S.: River runoff to the Baltic Sea: 1950–1990, *Ambio*, 23, 280–287, 1994. 2072

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Dietrich, G. and Schott, F.: Wasserhaushalt und Strömungen, in: *Meereskunde der Ostsee*, edited by L. Maagrad and G. Rheinheimer, Springer-Verlag Berlin, 33–41, 1974.
- Dietze, H. and Kriest, I.: ^{137}Cs off Fukushima Dai-ichi, Japan – model based estimates of dilution and fate, *Ocean Sci.*, 8, 319–332, doi:10.5194/os-8-319-2012, 2012. 2077, 2079
- 5 Dietze, H. and Löptien, U.: Revisiting “nutrient trapping” in global coupled biogeochemical ocean circulation models, *Global Biogeochem. Cy.*, 27, 1–20, doi:10.1002/gbc.20029, 2013. 2079
- Dietze, H., Oschlies, A., and Kähler, P.: Internal-wave-induced and double-diffusive nutrient fluxes to the nutrient-consuming surface layer in the oligotrophic subtropical North Atlantic, *Ocean Dynam.*, 54, 1–7, doi:10.1007/s10236-003-0060-9, 2004. 2086
- 10 Döös, K., Meier, H. E. M., and Döscher, R.: The Baltic haline conveyor belt or the overturning circulation and mixing in the Baltic, *Ambio*, 33, 258–262, 2004.
- Döscher, R. and Beckmann, A.: Effects of a bottom boundary layer parameterization in a coarse-resolution model of the North Atlantic Ocean, *J. Atmos. Ocean. Tech.*, 17, 698–707, 2000. 2083
- 15 Duhaut, T. H. A. and Straub, D. N.: Wind stress dependence on ocean surface velocity: implications for mechanical energy input to ocean circulation, *J. Phys. Oceanogr.*, 36, 202–211, 2006. 2078
- Eden, C. and Dietze, H.: Effects of mesoscale eddy/wind interactions on biologic new production and eddy kinetic energy, *J. Geophys. Res.*, 114, C05023, doi:10.1029/2008JC005129, 2009. 2078, 2109
- 20 Eilola, K. and Stigebrandt, A.: Spreading of juvenile freshwater in the Baltic Propoer, *J. Geophys. Res.*, 103, 27795–27807, 1998. 2076
- Ekman, M.: The world’s longest continued series of sea level observations, *Pure Appl. Geophys.*, 127, 73–77, 1988. 2082
- 25 Eppley, R. W. and Peterson, B. J.: Particulate organic matter flux and planktonic new production in the deep ocean, *Nature*, 282, 677–679, 1979. 2090
- Ezer, T. and Mellor, G. L.: A generalized coordinate ocean model and a comparison of the bottom boundary layer dynamics in terrain-following and in z-level grid, *Ocean Model.*, 6, 379–403, 2004. 2084
- 30 Fennel, W., Seifert, T., and Kayser, B.: Rossby radii and phase speeds in the Baltic Sea, *Cont. Shelf. Res.*, 11, 23–36, 1991. 2067, 2068

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Fennel, W., Radtke, H., Schmidt, M., and Neumann, T.: Transient upwelling in the central Baltic Sea, *Cont. Shelf. Res.*, 30, 2015–2026, 2010.
- Flynn, K.: Castles built on sand: dysfunctionality in plankton models and the inadequacy of dialogue between biologists and modellers, *J. Plankton Res.*, 27, 1205–1210, 2005. 2067
- 5 Fu, W., She, J., and Dobrynin, M.: A 20-year reanalysis experiment in the Baltic Sea using three-dimensional variational (3DVAR) method, *Ocean Sci.*, 8, 827–844, doi:10.5194/os-8-827-2012, 2012. 2081
- Gerdes, R.: A primitive equation ocean circulation model using a general vertical coordinate transformation: 2. Application to an overflow problem, *J. Geophys. Res.*, 93, 14703–14726, 1993.
- 10 Graham, L. P.: Climate change effects on river flow to the Baltic Sea, *Ambio*, 33, 235–241, 2004. 2072
- Griffies, S. M.: Elements of MOM4p1. GFDL Ocean Group Technical Report No. 6, NOAA/Geophysical Fluid Dynamics Laboratory, 444 pages, Version prepared on 16 December 2009, 2009. 2068, 2069, 2087, 2088
- 15 Griffies, S. M., Pacanowski, R. C., Schmidt, M., and Balaji, V.: Tracer Conservation with an Explicit Free Surface Method for z-coordinate Ocean Models, *Mon. Weather Rev.*, 129, 1081–1098, 2001. 2087
- Holgate, S. J., Matthews, A., Woodworth, P. L., Rickards, L. J., Tamisiea, M. E., Bradshaw, E., Foden, P. R., Gordon, K. M., Jevrejeva, S., and Pugh, J. P.: New data systems and products at the Permanent Service for Mean Sea Level, *J. Coastal Res.*, 29, 493–504, doi:10.2112/JCOASTRES-D-12-00175.1, 2013. 2082
- 20 Hordoir, R. and Meier, H. E. M.: Freshwater fluxes in the Baltic Sea: a model study, *J. Geophys. Res.*, 115, C08028, doi:10.1029/2009JC005604, 2010. 2072, 2073, 2074
- 25 Hordoir, R., Dieterich, C., Basu, C., Dietze, H., and Meier, H. E. M.: Freshwater outflow of the Baltic Sea and transport in the Norwegian current: a statistical correlation analysis based on a numerical experiment, *Cont. Shelf. Res.*, 64, 1–9, doi:10.1016/j.csr.2013.05.006, 2013. 2084
- Hurrell, J. W.: Decadal trends in the North Atlantic Oscillation: regional temperatures and precipitation, *Science*, 269, 676–679, 1995. 2087
- 30 Hurrell, J. W. and Deser, C.: North Atlantic climate variability: the role of the North Atlantic Oscillation, *J. Marine Syst.*, 78, 28–41, doi:10.1016/j.jmarsys.2008.11.026, 2009. 2087

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Jakobsen, F.: The major inflow to the Baltic Sea during January 1993, *J. Marine Syst.*, 6, 227–240, 1995. 2102
- Jedrasik, J., Cieslikiewicz, W., Kowalewski, M., Bradtke, K., and Jankowski, A.: 44 years hindcast of the sea level and circulation in the Baltic Sea, *Coast. Eng.*, 55, 849–860, doi:10.1016/j.coastaleng.2008.02.026, 2008. 2076, 2078, 2082
- Jevrejeva, S. and Moore, J. C.: Singular spectrum analysis of Baltic Sea ice conditions and large-scale atmospheric patterns since 1708, *Geophys. Res. Lett.*, 28, 4503–4506, 2001. 2087
- Jevrejeva, S., Moore, J. C., and Grinsted, A. G.: Influence of the Arctic Oscillation and El Niño–Southern Oscillation (ENSO) on ice conditions in the Baltic Sea: The wavelet approach, *J. Geophys. Res.*, 108, 4677, doi:10.1029/2003JD003417, 2003. 2087
- Jochum, M., Danabasoglu, G., Holland, M., Kwon, Y.-O., and Large, W. G.: Ocean viscosity and climate, *J. Geophys. Res.*, 113, C06017, doi:10.1029/2007JC004515, 2008. 2087
- Jones, C. G., Willén, U., Ullerstig, A., and Hansson, U.: The Rossby Centre Regional Atmospheric Climate Model Part I: Model climatology and performance for the present climate over Europe, *Ambio*, 33, 199–210, 2004. 2073
- Karlson, K., Rosenberg, R., and Bonsdorff, E.: Temporal and spatial large-scale effects of eutrophication and oxygen deficiency on benthic fauna in Scandinavian and Baltic waters: a review, *Oceanogr. Mar. Biol.*, 40, 427–489, 2002. 2067
- Killworth, P. D., Stainforth, D., Webb, D. J., and Paterson, S. M.: The development of a free-surface Bryan–Cox–Semtner ocean model, *J. Phys. Oceanogr.*, 21, 1333–1348, 1991. 2087
- Köller, M., Käse, R. H., and Herrmann, P.: Interannual to multidecadal variability and predictability of North Atlantic circulation in a coupled earth system model with parameterized hydraulics, *Tellus A*, 62, 569–578, 2010. 2085
- Koslowski, G. and Loewe, P.: The western Baltic sea ice season in terms of a mass-related severity index: 1879–1992, Part 1: Temporal variability and association with the North Atlantic Oscillation, *Tellus A*, 269, 66–74, 1994. 2087
- Kösters, F., Käse, R. H., Schmittner, A., and Herrmann, P.: The effect of Denmark Strait overflow on the Atlantic meridional overturning circulation, *Geophys. Res. Lett.*, 32, L04602, doi:10.1029/2004GL022112, 2005. 2085, 2091
- Large, W. G. and Yeager, S. G.: Diurnal to decadal global forcing for ocean and sea-ice models: the data sets and flux climatologies, NCAR Technical Note, TN-460, Boulder Colorado, USA, 2004. 2074

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Large, W. G., McWilliams, J. C., and Doney, S. C.: Oceanic vertical mixing – A review and a model with nonlocal boundary-layer parameterization, *Rev. Geophys.*, 32, 363–403, 1994. 2079
- Ledwell, J. R., McGillicuddy, D. J., and Anderson, L. A.: Nutrient flux into an intense deep chlorophyll layer in a mode-water eddy, *Deep-Sea Re. II*, 55, doi:10.1016/j.dsr2.2008.02.005, 2008. 2078, 2079
- Lehmann, A., Kraus, W., and Hinrichsen, H.-H.: Effects of remote and local atmospheric forcing on circulation and upwelling in the Baltic Sea, *Tellus A*, 54, 299–316, 2002. 2072, 2076, 2077, 2082
- Lehmann, A., Myrberg, K., and Höflisch, K.: A statistical approach to coastal upwelling in the Baltic Sea based on the analysis of satellite data for 1990–2009, *Oceanologica*, 54, 369–393, doi:10.5697/oc.54-3.369, 2012. 2080
- Lehmann, A., Hinrichsen, H.-H., and Getzlaff, K.: Identifying potentially high risk areas for environmental pollution in the Baltic Sea, *Boreal Environ. Res.*, 19, in press, 2014a. 2091
- Lehmann, A., Hinrichsen, H.-H., Getzlaff, K., and Myrberg, K.: Quantifying the heterogeneity of hypoxic and anoxic areas in the Baltic Sea by a simplified coupled hydrodynamic-oxygen consumption model approach, *J. Marine Syst.*, 134, 20–28, doi:10.1016/j.jmarsys.2014.02.012, 2014b. 2068
- Leppäranta, M. and Myrberg, K.: *Physical Oceanography of the Baltic Sea*, Springer, ISBN: 978-3-540-79702-9, 85, 296–85, Berlin, Heidelberg, New York, 2009. 2066, 2075, 2086
- Liu, N., Eden, C., Dietze, H., Wu, D., and Lin, X.: Model-based estimates of the heat budget in the East China Sea, *J. Geophys. Res.*, 115, C08026, doi:10.1029/2009JC005869, 2010. 2077, 2079
- Löptien, U. and Meier, H. E. M.: The influence of increasing water turbidity on the sea surface temperature in the Baltic Sea: a model sensitivity study, *J. Marine Syst.*, 88, 323–331, 2011, 133, 2894–2904, 2011. 2072, 2073, 2074, 2079, 2088
- Löptien, U. and Ruprecht, E.: Effect of Synoptic Systems on the Variability of the North Atlantic Oscillation, *Mon. Weather Rev.*, 133, 2894–2904, 2005. 2087
- Löptien, U., Zolina, O., Gulev, S., Latif, M., and Soloviev, V.: Cyclone life cycle characteristics over the Northern Hemisphere in coupled GCMs, *Clim. Dynam.*, 31, 507–532, doi:10.1007/s00382-007-0355-5, 2008. 2073

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Löptien, U., Eden, C., Timmermann, A., and Dietze, H.: Effects of biologically induced differential heating in an eddy-permitting coupled ocean-ecosystem model, *J. Geophys. Res.*, 114, C06011, doi:10.1029/2008JC004936, 2009. 2079

Löptien, U., Mårtensson, S., Meier, H. E. M., and Höglund, A.: Long-term characteristics of simulated ice deformation in the Baltic Sea (1962–2007), *J. Geophys. Res.*, 23, 279–290, 2013. 2087

Martin, A. P. and Richards, K. J.: Mechanisms for the vertical nutrient transport within a North Atlantic mesoscale eddy, *Deep-Sea Res. Pt. II*, 48, 757–773, 2001. 2078

McGillicuddy, D. J., Anderson, L. A., Bates, N. R., Bibby, T., Buesseler, K. O., Carlson, C. A., Davis, C. S., Ewart, C., Falkowski, P. G., Goldthwait, S. A., Hansell, D. A., Jenkins, W. J., Johnson, R., Kosnyrev, V. K., Ledwell, J. R., Li, Q. P., Siegel, D., and Steinberg, D. K.: Eddy/wind interactions stimulate extraordinary mid-ocean plankton blooms, *Science*, 316, 1021–1026, doi:10.1126/science.1136256, 2007. 2078

MacKenzie, B. R., Hinrichsen, H.-H., Plikhs, M., Wieland, K., and Zezera, A. S.: Quantifying environmental heterogeneity: habitat size necessary for successful development of cod *Gadus morhua* eggs in the Baltic Sea, *Mar. Ecol.-Prog. Ser.*, 193, 143–156, 2000. 2067

Meier, H. E. M.: Modeling the pathways and ages of inflowing salt- and freshwater in the Baltic Sea, *Estuar. Coast. Shelf S.*, 74, 610–627, 2007.

Meier, H. E. M. and Faxen, T.: Performance analysis of a multiprocessor coupled ice–ocean model for the Baltic Sea, *J. Atmos. Ocean. Tech.*, 19, 114–124, 2002. 2072, 2073

Meier, H. E. M. and Kauker, F.: Modeling decadal variability of the Baltic Sea: 2. Role of freshwater inflow and large-scale atmospheric circulation for salinity, *J. Geophys. Res.*, 108, 3368, doi:10.1029/2003JC001799, 2003. 2072, 2073

Meier, H. E. M., Döscher, R., Coward, A. C., Nycander, J., and Döös, K.: RCO – Rossby Centre regional Ocean climate model: model description (version 1.0) and first results from the hind cast period 1992/1993, *SMHI Reports Oceanography*, 26, 1999. 2073

Meier, H. E. M., Höglund, A., Döscher, R., Andersson, H., Löptien, U., and Kjellström, H.: Quality assessment of atmospheric surface fields over the Baltic Sea from an ensemble of regional climate model simulations with respect to ocean dynamics, *Oceanologica*, 53, 193–227, 2011. 2073

Meier, H. E. M., Andersson, H. C., Arheimer, B., Blenckner, T., Chubarenko, B., Donnelly, C., Eilola, K., Gustafsson, B. G., Hansson, A., Havenhand, J., Höglund, A., Kuznetsov, I., Mackenzie, B. R., Müller-Karulis, B., Neumann, T., Niiranen, S., Piwowarczyk, J., Raud-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

sepp, U., Reckermann, M., Ruoho-Airola, T., Savchuk, O. P., Schenk, F., Schimanke, S., Väli, G., Weslawski, and, Zorita, E., J.-M.: Comparing reconstructed past variations and future projections of the Baltic Sea excosystem – first results from multi-model ensemble simulations, *Environ. Res. Lett.*, 7, 034005, doi:10.1088/1748-9326/7/3/034005, 2012

5 Mesinger, F.: A method for construction of second-order accurate difference schemes permitting no false two-grid-interval waves in the height field, *Tellus*, 25, 444–457, 1973. 2087

Myrberg, K. and Andrejev, O.: Modelling of the circulation, water exchange and water age properties of the Gulf of Bothnia, *Oceanologica*, 48, 55–74, 2006. 2076, 2077

Neumann, T. and Schernewski, G.: Eutrophication in the Baltic Sea and shifts in nitrogen fixation analysed with a 3D ecosystem model, *J. Marine Syst.*, 74, 592–602, 2008. 2082

10 Neumann, T., Eilola, K., Gustafsson, B., Müller-Karulis, B., Kuznetsov, I., Meier, H. E. M., and Savchuk, O. P.: Extremes of temperature, oxygen and blooms in the Baltic Sea in a changing climate, *Ambio*, 41, 574–585, 2012. 2066

Pacanowski, R. C. and GRIFFIES, S. M.: *The MOM3 Manual*, NOAA/Geophysical Fluid Dynamics Laboratory, Princeton, USA, 680 pp., 1999. 2087

Pahlow, M., Dietze, H., and Oschlies, A.: Optimality-based model of phytoplankton growth and diazotrophy, *Mar. Ecol.-Prog. Ser.*, 489, 1–16, doi:10.3354/meps10449, 2013. 2081, 2091

15 Pawlak, J. F., Laamanen, M., and Andersen, J. H.: Eutrophication in the Baltic Sea – an integrated thematic assessment of the effects of nutrient enrichment in the Baltic Sea region, An executive summary, *Baltic Sea Environment Proceedings*, 115A, Helsinki Commission (Baltic Marine Environment Protection Commission), 1–18, 2009. 2066

Permanent Service for Mean Sea Level: “Tide Gauge Data”, available at: www.psmsl.org/data/obtaining/ (last access: 12 September 2012), 2012. 2072, 2082

Roberts, M. J. and Wood, R. A.: Topographic sensitivity studies with a Bryan-Cox-type ocean model, *J. Phys. Oceanogr.*, 27, 823–836, 1997.

25 Samuelsson, M. and Stigebrandt, A.: Main characteristics of the long-term sea level variability in the Baltic Sea, *Tellus A*, 48, 672–683, 1996. 2082

Samuelsson, P., Jones, C. G., Willén, U., Ullerstig, A., Gollvik, S., Hansson, U., Jansson, C., Kjellström, E., Nikulin, G., and Wyser, K.: The Rossby Centre Regional Climate model RCA3: model description and performance, *Tellus A*, 63, 4–23., 2011. 2073

30 Sayin, E. and Kraus, W.: A numerical study of the water exchange through the Danish Straits, *Tellus A*, 48, 324–341, 1996.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Schinke, H. and Matthäus, W.: On the cause of major Baltic inflows – an analysis of long time series, *Cont. Shelf. Res.*, 18, 67–97, 1998. 2071
- Schmidt, M.: A benchmark for the parallel code used in FMS and MOM-4, *Ocean Model.*, 17, 49–67, 2007. 2089
- 5 Seifert, T., Tauber, F., and Kayser, B.: A High-Resolution Spherical Grid Topography of the Baltic Sea, revised Edn., 2001. 2069
- Shapiro, R: Smoothing, filtering, and boundary effects, *Rev. Geophys.*, 8, 359–387, 1970. 2069
- Sjöberg, B.: Sea and coast, national atlas of Sweden, SMHI Norrköping, SNA Publishing, 128 pp., 1992. 2075
- 10 Song, Z., Qiao, F., Lei, X., and Wang, C.: Influence of parallel computational uncertainty on simulations of the Coupled General Climate Model, *Geosci. Model Dev.*, 5, 313–319, doi:10.5194/gmd-5-313-2012, 2012. 2088
- Stacey, M. W., Pond, S., and Nowak, Z. P.: A numerical model of the circulation in Knight Inlet British Columbia, Canada, *J. Phys. Oceanogr.*, 25, 1037–1062, 1995. 2069
- 15 Sverdrup, H. U.: On conditions for the vernal blooming of phytoplankton, *Journal du Conseil International our l'Exploration de la Mer*, 18, 287–295, 1953. 2090
- Tinz, B.: On the relation between annual maximum extent of ice cover in the Baltic Sea and sea level pressure as well as air temperature field, *Geophysica*, 32, 319–341, 1996. 2087
- Uppala, S. M., Kallberg, P. W., Simmons, A. J., Andrae, U., Bechtold, V. D., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Holm, E., Hoskins, B. J., Isaksen, L., Janssen, P. A. E. M., Jenne, R., McNally, A. P., Mahfouf, J.-F., Morcrette, J.-J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The ERA-40 re-analysis, *Q. J. Roy. Meteor. Soc.*, 131, 621, doi:10.1256/qj.04.176, 2005. 2073
- 20 Vahtera, E., Conley, D. J., Gustafsson, B. G., Kuosa, H., Pitkänen, H., Savchuk, O. P., Tamminen, T., Viitasalo, M., Voss, M., Wasmund, N., and Wulff, F.: Internal ecosystem feedbacks enhance nitrogen-fixing cyanobacteria blooms and complicate management in the Baltic Sea, *Ambio*, 36, 186–194, 2007. 2081
- 30 von Storch, H., Langenberg, H., and Frauke, F.: A spectral nudging technique for dynamical downscaling purposes, *Mon. Weather Rev.*, 123, 3664–3673, 2000. 2073

- Ward, B. A., Dutkiewicz, S., Moore, C. M., and Follows, M. J.: Iron, phosphorous, and nitrogen supply ratios define the biogeography of nitrogen fixation, *Limnol. Oceanogr.*, 58, 2059–2075, doi:10.4319/lo.2013.58.6.2059, 2013. 2091
- 5 Wasmund, N. and Uhlig, S.: Phytoplankton trends in the Baltic Sea, *ICES J. Mar. Sci.*, 60, 177–186, 2003. 2066
- Woodworth, P. L. and Player, R.: The permanent service for mean sea level: an update to the 21st century, *J. Coastal Res.*, 19, 287–295, 2003. 2082
- Wyrтки, K.: Der große Salzeinbruch in der Ostsee im November und Dezember 1951, *Kieler Meeresforschung*, 10, 19–25, 1954. 2071
- 10 Xu, Y. and Scott, R. B.: Subtleties in forcing eddy resolving ocean models with satellite wind data, *Ocean Model.*, 20, 240–251, 2008. 2078
- Zhai, X. and Greatbatch, R.: Wind work in a model of the northwest Atlantic Ocean, *Geophys. Res. Lett.*, 34, L04606, doi:10.1029/2006GL028907, 2007. 2078
- 15 Zapadka, T., Woźnick, B., and Jerzy, D.: A more accurate formula for calculating the net long-wave radiation in the Baltic Sea, *Oceanologia*, 49, 4, 449–470, 2007. 2074

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)**Table 1.** List of model grid boxes where the bathymetry was deepened to increase the salt water supply to the Baltic Sea.

Longitude [° E]	Latitude [° N]	Straight	New depth [m]
12.571	56.035	Øresund	8.0987
12.533	55.955–56.055	Øresund	9.0331
12.495	56.075	Øresund	9.0331
10.917–10.992	54.695–54.915	Great Belt	17.2539
10.842	55.555–55.795	Great Belt	21.4247
10.879	55.555–55.755	Great Belt	16.8284

MOMBA 1.1

H. Dietze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 2.** Volume transport [km^3] through the Danish Straits from 5 to 26 January 1993 to the Baltic.

	Øresund	Big Belt	Small Belt
Observation (Jakobsen, 1995)	73	188	27
Model (MOMBA)	71	141	30




Fig. 1. Horizontal model grid and bathymetry. Panel (a) shows the entire model domain with red crosses indicating internationally renowned monitoring stations. Panel (b), (c) and (d) are close-ups of the Danish Straits, Gulf of Finland and the region between Åland and Finland, respectively. The coloured bins denote the water depth of each model water column. The white areas are considered as land by the model. The actual coastline is denoted by a black line.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 2. Vertical model grid. The nominal box thicknesses (Δz) as a function of depth.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. Total river runoff applied in MOMBA in units Sv ($= 10^6 \text{ m}^3 \text{ s}^{-1}$).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 4. Simulated mean (1987 to 1999) surface circulation. **(a)** The colour and arrows denote the magnitude and direction of the velocity, respectively. Note that the actual spatial resolution is 20 times higher than indicated by the arrows, both in zonal and meridional direction. **(b)** shows the persistency R of the surface circulation (in colour, as defined by Eq. 5) along with bathymetry (black contours every 25 m).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 5. Sea surface salinity averaged over 1987 to 1999. Panel (a) shows a composite based on ICES bottle observations (upper 10 m, binned monthly into boxes of 6 nm resolution and subsequently averaged over time). Panel (b) refers to MOMBA.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. Eddy kinetic energy, calculated based on surface velocities according to Eq. (6) in units $\text{cm}^2 \text{s}^{-1}$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 7. Daily mean Ekman pumping calculated from the curl of the wind stress in units day^{-1} (in color) overlaid by surface currents (arrows) for an exemplary time slice. Strong upwelling is denoted in red. The calculation of the wind stress takes the ocean currents into account (as explained, e.g., in Eden and Dietze, 2009). Note, that the Ekman pumping features variability correlated with the surface currents on spacial scales much smaller than those that could possibly be effected by the ≈ 25 km resolution wind fields.




Fig. 8. Sea surface temperature at station AE, BY15 and F9 in panels (a), (b) and (c), respectively. The positions of the respective stations are indicated in Fig. 1. Observations are from the Swedish Oceanographic Data Centre at SMHI (SHARK).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 9. Assessment of simulated sea surface temperature with weekly composites observed from space throughout 1990 to 1999 (BSH SST, Sect. 3.3). The color shows the variance explained by the model in units %.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 10. Observed (left panel, weekly composite observed from space, BSH SST, Sect. 3.3) and simulated (right panel, daily mean) sea surface temperature in °C. Note that areas of missing observations are flagged white in both the left and right panel.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 11. Near-bottom salinities at station BY2, BY5 and BY15 in panels (a), (b) and (c), respectively. The positions of the respective stations are indicated in Fig. 1. Observations are from the Swedish Oceanographic Data Centre at SMHI (SHARK).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 12. Detrended daily mean sea surface height at Stockholm and Landsort (upper and lower panel, respectively). The black (red) line denotes observed (simulated) heights.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 13. Topographic slope and thickness of lowermost wet grid box in MOMBA. Panel (a) shows the difference of the number of grid boxes representing neighbouring water columns. Of the respective, up to four, differences (calculated from all neighbours) panel (a) shows the maximum difference. Differences smaller than two indicate “resolved” topographic slopes and are masked by white areas. Panel (b) shows the thickness of the lowermost wet grid box (note that we use partial cells).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 14. Simulated deep saline inflow into the Bornholm Basin through Hamrre Strait along the transect indicated in the inlayed map. The color denotes salinity. The thick black dashed line is the surface mixed layer depth as calculated by the KPP scheme. The white (grey) contours denote upwards (downwards) velocities in m day^{-1} . All properties are averaged over one hour.




Fig. 15. Maximum near-bottom salinity at station BY5 in the period 9 January to 18 April 1993. Panel (a) and (b) show maximum simulated salinities as function of horizontal and vertical Laplacian viscosity, respectively. The dashed black line in panel (a) shows the maximum salinity simulated with the “standard” horizontal Laplacian viscosity of 0. Observed salinities peak at 18.5 PSU (Fig. 11b).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 16. Median salinity profiles for the period 1987–1999 with 10th- and 90th-percentiles at stations AE, BY5 and BY15 in panels (a), (b) and (c) respectively. The locations of the stations are indicated in Fig. 1. Observations refer to SHARK.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 17. Ice covered area in the Baltic Sea. The red line denotes an estimate based on digitised ice-charts from the operational ice service provided by SMHI. The black line shows daily mean simulated ice extent.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 18. Snapshots of observed (left column) and simulated (right column) ice concentrations at exemplary days in mid February, during a particularly warm (1989, upper panels) and cold winter (1996, lower panels). The units are %.




Fig. 19. Leading principle components of observed (red line) and simulated (black line) seasonal mean ice concentrations. The blue line (and right y axis) depicts the (unitless) NAO-index for comparison.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 20. Leading EOFs of (a) observed and (b) modeled seasonal mean ice concentrations [%].

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fig. 21. (a) Simulated surface salinity (color) and sea surface height (black contours in units m). (b) Supposedly identical to (a) but slightly different due to “computational uncertainty”. (c) temporal evolution of sea surface salinity at station BY2. The black and red line correspond to identical setups which differ due to “computational uncertainty”. (d) Temporal evolution of salinity at 80 m depth at station BY5. Black and red lines are coded like in (c). The location of station BY2 and BY5 are shown in the upper panels and in Fig. 1a, respectively.




Fig. 22. Computational performance (wall clock hours required to integrate one year) as a function of allocated computational cores. Black crosses, grey circles and grey x refer to architectures based on Sandybridge, Gainstown and Harpertown CPUs, respectively. The red line depicts a theoretic, ideal parallel performance of the code on the Sandybridge architecture.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
