



This discussion paper is/has been under review for the journal Geoscientific Model Development (GMD). Please refer to the corresponding final paper in GMD if available.

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Earth System Science Department, UC Irvine California, USA

Received: 29 April 2015 – Accepted: 6 May 2015 – Published: 27 May 2015

Correspondence to: M. J. Prather (mprather@uci.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

GMDD

8, 4051–4073, 2015

Photolysis rates in
correlated
overlapping cloud
fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

A new approach for modeling photolysis rates (J values) in atmospheres with fractional cloud cover has been developed and implemented as Cloud-J – a multi-scattering eight-stream radiative transfer model for solar radiation based on Fast-J. Using observed statistics for the vertical correlation of cloud layers, Cloud-J 7.3 provides a practical and accurate method for modeling atmospheric chemistry. The combination of the new maximum-correlated cloud groups with the integration over all cloud combinations represented by four quadrature atmospheres produces mean J values in an atmospheric column with root-mean-square errors of 4% or less compared with 10–20% errors using simpler approximations. Cloud-J is practical for chemistry-climate models, requiring only an average of 2.8 Fast-J calls per atmosphere, vs. hundreds of calls with the correlated cloud groups, or 1 call with the simplest cloud approximations. Another improvement in modeling J values, the treatment of volatile organic compounds with pressure-dependent cross sections is also incorporated into Cloud-J.

1 Introduction

Photolysis, the dissociation of molecules upon absorbing sunlight, drives atmospheric chemistry and controls the composition of the air we breathe. Photolysis rates are governed by the intensity of sunlight, which is altered by scattering and absorption processes within the atmosphere. Clouds, aerosols, and gases control these processes; but ambiguity in the representation of clouds in atmospheric models is currently the largest source of uncertainty in photolysis rates. This paper presents a new, pragmatic approach for representing the overlap of clouds derived from observations and cloud models, and then provides several practical approximations with marginal computational costs that can be readily incorporated in atmospheric chemistry models. This computer code is a major expansion of Fast-J (Wild et al., 2000; Bian and Prather, 2002; Neu et al., 2007) and is presented here as Cloud-J 7.3 (Cloud-J contains Fast-J

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and thus continues that numbering sequence, for which Fast-J 7.2 was the last released version.)

Clouds can increase photolysis rates through scattered sunlight, but they can greatly reduce them by shadowing. Modeling the scattering by cloud layers in a column atmosphere and resulting photolysis rates is practical, as in Fast-J, if the layers are horizontally uniform across the modeled air parcel (defined typically as a rectilinear box bounded by latitude, longitude and pressure surfaces). Clouds layers, however, have horizontal scales of a few kilometers (Slobodda et al., 2015), and thus are represented in global and regional models as fractional coverage in each parcel. In calculating the average photolysis or heating rates through the column atmosphere, one must know how the cloud fractions overlap. Early modeling assumed that model layers consisted of maximally overlapped groups (MAX) that would be randomly overlapped relative to one another (MAX-RAN) (Briegleb, 1992; Feng et al., 2004). A more accurate description of cloud overlap is that clouds are correlated throughout the column atmosphere with a correlation length ranging from 1.5 to 3 km in height (Pincus et al., 2005; Naud and DelGenio, 2006; Kato et al., 2010; Oreopoulos et al., 2012).

A practical application of this cloud overlap information, merging maximally overlapped groups that are correlated with each other (MAX-COR) is defined in Sect. 2, where the impact of cloud overlap models on photolysis rates (J values) is also shown. Cloud overlap models generate statistics that lead to a large number of weighted independent column atmospheres (ICAs), where the number is too large to be used directly to calculate photolysis or heating rates in global models. Section 3 looks at the simple cloud models and the approaches to approximate the sum over ICAs, examining their errors. Another recent development in modeling photolysis rates included with Cloud-J is the treatment of volatile organic compounds with pressure-dependent cross sections, presented in Sect. 4. Recommendations for the cloud-overlap model and the ICA-approximation method are discussed in Sect. 5.

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2 Overlap models for fractionally cloudy atmospheres

Typically, meteorological forecasts or climate models used in atmospheric chemistry models report fractionally cloudy atmospheres (FCAs) in each grid-square. Computation of the photolysis or heating rates in an FCA requires knowledge of how the clouds in each layer overlap. The calculation of J values in most atmospheric chemistry models today involves solving the radiative transfer equations in a plane-parallel atmosphere where the vertical layers can be highly inhomogeneous but the horizontal planes are uniform (Stamnes et al., 1988; Wild et al., 2000; Tie et al., 2003). Thus, the only workable method (other than 3-D radiative transfer) is to represent the FCA by a number of independent column atmospheres (ICAs) where each ICA is either 100% cloudy or clear in each layer. The fractional cloud overlap model determines the layer-structure, weighting, and number of ICAs. Other simple cloud models approximated overlap by: (i) ignoring clouds entirely (clear sky), (ii) averaging the fractional cloud, f , over each layer, conserving total cloud water (average clouds); and (iii) increasing the cloud fraction by using $f^{3/2}$, increasing the total cloud water in proportion, and then averaging over the layer (Briegleb, 1992). These methods will be compared with cloud overlap models in Sect. 3. Here, we focus on how the ICAs differ across cloud overlap models.

2.1 Random overlap (RAN)

The ways in which fractionally cloudy layers can overlap is shown schematically in Fig. 1. One assumption is random overlap (RAN). In this case the likelihood (fractional weight, w) of having the cloud in layer L1 fall below the cloud in layer L2 is random and hence equals f^{L1} . This particular pairing – cloud below cloud – becomes ICA #1. Superscripts in the equations below refer to atmospheric layers. The likelihood for the

GMDD

8, 4051–4073, 2015

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



clear layer under the cloudy layer is by default the complement.

$$w^{L1}(\#1) = f^{L1} \quad (1)$$

$$w^{L1}(\#2) = 1 - f^{L1} \quad (2)$$

The likelihood of the cloudy layer in L2 above is

$$w^{L2}(\#1) = w^{L2}(\#2) = f^{L2} \quad (3)$$

The total weight W for each ICA #1 and #2 is then the product of w^{L1} and w^{L2} .

$$W^{L1-L2}(\#1) = w^{L1}(\#1)w^{L2}(\#1) = f^{L1}f^{L2} = 0.15 \times 0.20 = 3\% \text{ (from Fig. 1)} \quad (4)$$

$$W^{L1-L2}(\#2) = w^{L1}(\#2)w^{L2}(\#2) = (1 - f^{L1})f^{L2} = 0.85 \times 0.20 = 17\% \quad (5)$$

Similar rules apply to ICAs #3 and #4,

$$w^{L1}(\#3) = f^{L1} \quad \text{and} \quad w^{L2}(\#3) = 1 - f^{L2} \quad (6)$$

$$w^{L1}(\#4) = 1 - f^{L1} \quad \text{and} \quad w^{L2}(\#4) = 1 - f^{L2}. \quad (7)$$

and thus

$$W^{L1-L2}(\#3) = w^{L1}(\#3)w^{L2}(\#3) = f^{L1}(1 - f^{L2}) = 0.15 \times 0.80 = 12\% \text{ (Fig. 1)} \quad (8)$$

$$W^{L1-L2}(\#4) = w^{L1}(\#4)w^{L2}(\#4) = (1 - f^{L1})(1 - f^{L2}) = 0.85 \times 0.80 = 68\% \quad (9)$$

15 ICAs #1 and #3 are tagged as cloudy in L1, and ICAs #2 and #4 are tagged as clear in L1. The sum of cloudy fractions in L1 must be conserved: $3\% + 12\% = 15\% = f^{L1}$. One of the problems in implementing a full RAN model is that the number of ICAs scales as 2^{NL} , where NL is the number of cloudy layers in the RAN group.

GMDD

8, 4051–4073, 2015

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Correlated overlap (COR)

When correlated, the likelihood of a cloudy layer underlying another cloud is greater than random, $w^{L1}(\#1) > f^{L1}$, by a factor $g > 1$. The correlation coefficient cc ranges from 0 (random) to 1 (maximal overlap).

$$g = 1 + cc(1/f^{L2} - 1), \text{ subject to } g \leq 1/f^{L1} \text{ and } g \leq 1/f^{L2} \quad (10)$$

Hence for $cc > 0$ we have increased likelihood of the cloud in L1 falling underneath the cloud in L2. For the example in Fig. 1, $cc = 1/2$ and $g = 3$.

$$w^{L1}(\#1) = gf^{L1} = 3 \times 0.15 = 45\% \quad (11)$$

$$w^{L1}(\#2) = 1 - gf^{L1} = 1 - 0.45 = 55\% \quad (12)$$

The likelihood of clouds in L1 falling below the clear section in L2 are reduced and calculated from the requirement that the sum of cloudy fractions in L1 is still f^{L1} .

$$w^{L1}(\#3) = f^{L1}(1 - gf^{L2})/(1 - f^{L2}) = 0.15 \times (1 - 3 \times 0.20)/0.80 = 7.5\% \quad (13)$$

By complement, the weighting of clear sky in layer L1 under clear sky in layer L2 is

$$w^{L1}(\#4) = 1 - w^{L1}(\#3) = 1 - 0.075 = 92.5\% \quad (14)$$

Note that if $cc = 0$, or $f^{L2} = 1$, or $f^{L1} = 1$, then $g = 1$ and COR defaults to RAN. The two additional limits on g in Eq. (10) are required to keep $w^{L1}(\#2)$ and $w^{L1}(\#3)$ positive. The w^{L2} weights remain simply

$$w^{L2}(\#1) = w^{L2}(\#2) = f^{L2} \quad (15)$$

$$w^{L2}(\#3) = w^{L2}(\#4) = (1 - f^{L2}). \quad (16)$$

The combined ICA weights are

$$W^{L1-L2}(\#1) = w^{L1}(\#1)w^{L2}(\#1) = gf^{L1}f^{L2} = 3 \times 0.15 \times 0.20 = 9\% \quad (17)$$

$$W^{L1-L2}(\#2) = w^{L1}(\#2)w^{L2}(\#2) = (1 - gf^{L1})f^{L2} = 0.55 \times 0.20 = 11\% \quad (18)$$

$$W^{L1-L2}(\#3) = w^{L1}(\#3)w^{L2}(\#3) = f^{L1}(1 - gf^{L2}) = 0.15 \times 0.40 = 6\% \quad (19)$$

$$W^{L1-L2}(\#4) = w^{L1}(\#4)w^{L2}(\#4) = 1 - f^{L2} - f^{L1}(1 - gf^{L2}) = 1 - 0.20 - 0.06 = 74\% \quad (20)$$

As in RAN, ICAs #1 and #3 are tagged as cloudy in layer L1, and ICAs #2 and #4 are tagged as clear in layer L1, and the sum of cloudy fractions is conserved ($9\% + 6\% + 15\% = f^{L1}$), but with different weightings. The COR model also has ICAs scaling as 2^{NL} .

2.3 Maximal overlap (MAX)

For maximal overlap of clouds (MAX) as in Fig. 1, the two layers L1 and L2 form a MAX group G1 consisting of 1 clear-sky column (80% fractional coverage) and 2 cloudy columns – one with clouds in both layers (15%) and one with a cloud only in the upper layer L2 (5%). The clear-sky column does not occur if any of the MAX layers has a cloud fraction of 100%. For continuous cloud fractions, the number of ICAs equals the number of different cloud fractions present (plus 1 if clear sky present), and thus it scales as NL, the number of layers in the MAX group. A MAX group is characterized by the number of unique cloudy fractions (f_1, f_2, f_3, \dots) and their weights (w_1, w_2, w_3, \dots) with a cloudy sum $F = \sum f_i \leq 1$ and a (possible) clear-sky column of fraction $1 - F$. As in the earlier Fast-J work (Neu et al., 2007), the cloud fractions in Cloud-J are quantized to limit the number of ICAs in a MAX group. The examples here use 10 bins, and hence cloud fractions are limited to 0, 10, 20, 30, ..., 100%. With this binning, the in-cloud water content is scaled to conserve the cloud water content in each layer. This approximation is then resolution independent in terms of the number of model layers and limits each MAX group to 10 ICAs.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.4 Maximal groups with correlated overlap (MAX-COR)

The MAX-COR model generates ICAs from upper and lower layers that are MAX groups. For a general approach, we will assume that the upper group G2 consists of N2 cloudy columns (members) with fractions $f_{J2=1:N2}^{G2}$ totaling F^{G2} and 1 clear-sky column (member) of fraction $1 - F^{G2}$. Likewise, the lower group G1 has N1 cloudy members with fractions $f_{J1=1:N1}^{G1}$ totaling F^{G1} and a clear-sky with fraction $1 - F^{G1}$. Each of the cloudy members in group G1, $f_{J1=1:N1}^{G1}$, will be paired with N2 (cloudy) + 1 (clear) member above. The number of ICAs will be $(N1 + 1)(N2 + 1)$, assuming that there are clear-sky members in both groups. The ICA sequence (J1, J2) is then

$$(1, 1), (2, 1), (3, 1), \dots, (N1 + 1, 1), (1, 2), (2, 2), (3, 2), \dots, (N1 + 1, N2 + 1) \quad (21)$$

such that ICA #M is composed of members

$$J1 = (M - 1) \bmod (N1 + 1) + 1 \quad (22)$$

$$J2 = \text{integer}((M - 1)/(N1 + 1)) \bmod (N2 + 1) + 1 \quad (23)$$

The correlation factor is same for all members, derived from the total cloudy fractions F^{G1} and F^{G2} .

$$g = 1 + cc(1/F^{G2} - 1), \text{ subject to } g \leq 1/F^{G1} \text{ and } g \leq 1/F^{G2} \quad (24)$$

For convenience denote $J1 \leq N1$ as cloudy^{G1}, $J1 = N1 + 1$ as clear^{G1}, $J2 \leq N2$ as cloudy^{G2}, and $J2 = N2 + 1$ as clear^{G2}. Then the weightings for the G1 members are

$$w^{G1}(\text{cloudy}^{G1}, \text{cloudy}^{G2}) = gf_{J1}^{G1} \quad (25)$$

$$w^{G1}(\text{clear}^{G1}, \text{cloudy}^{G2}) = 1 - \sum_{J1=1:N1} gf_{J1}^{G1} = 1 - gF^{G1} \quad (26)$$

Conserving each cloudy group member's fractional area in G1 gives the weights under G2 clear sky.

$$w^{G1}(\text{cloudy}^{G1}, \text{clear}^{G2}) = f_{j1}^{G1}(1 - gF^{G2}) / (1 - F^{G2}) \quad (27)$$

$$w^{G1}(\text{clear}^{G1}, \text{clear}^{G2}) = 1 - F^{G1}(1 - gF^{G2}) / (1 - F^{G2}) \quad (28)$$

- 5 All of these formulae work also if $F^{G1} > F^{G2}$, and if $F^{G1} = 0$ or 1 (same for F^{G2}). A special case of MAX-COR is MAX-RAN when $cc = 0$. With the cloud fractions binned into 10 intervals, then the number of ICAs for MAX-COR or MAX-RAN models scales as 10^{NG} , where NG is the number of MAX groups.

2.5 J value errors

10 Our recommended cloud overlap model uses the information on vertical correlations (Pincus et al., 2005; Naud and DelGenio, 2006; Kato et al., 2010; Oreopoulos et al., 2012), which shows correlations lengths of order 1.5 km in the lowest layers increasing to 3 km or more in the upper troposphere. Since a true COR model will scale as 2^{NL} and becomes rapidly impractical for high-resolution models, we define 6 MAX groups as those layers within a correlation length: 0–1.5 km altitude, 1.5–3.5, 3.5–6, 6–9, 9–13, 15 and > 13 km. The MAX groups collapse if there are no cloud fractions within the layers of the group. In looking at how this model aligned the clouds for realistic FCAs, we found that extensive cirrus fractions in the uppermost layers prevented the correlation of small-fraction cumulus below. Thus a 7th MAX group was added if there was a cirrus shield (defined from top down as adjacent ice-only clouds with $f > 0.5$). We chose a correlation factor $cc = 0.33$, similar to one e-fold, between groups, as the groups are chosen to be separated by about one correlation length. This model is denoted G6/.33. 20 Two other G6 models were tested: $cc = 0.00$ corresponds to MAX-RAN (G6/.00); and $cc = 0.99$ is close to one large MAX group (G6/.99). Because of the cloud-fraction bins and the fixed correlation-length groups, the number of ICAs is bounded by 5×10^6 25 (including the cirrus shield). This limit is resolution independent and was never reached

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in any FCAs examined here (highest number of ICAs for one FCA was 3500). The major computational cost comes with the Fast-J computation, and the methods for approximating the average of J values over all ICAs (Sect. 3) requires at most 4 Fast-J calculations no matter how many ICAs.

Two other cloud overlap models tested here are the MAX-RAN groupings, the same or similar to earlier work (Feng et al., 2004; Neu et al., 2007). The model G0 declares all adjacent cloudy layers to be a MAX group, and all such groups separated by a clear layer to be RAN overlapped. This model has the potential to be unstable with increasing layers as alternating clear and cloud layers results in $2^{NL/2}$ ICAs. In our tests using meteorological data with $NL = 36$, the maximum number of ICAs, 375, was well below this limit. The model G3 declares MAX groups by 3 atmospheric regimes: 0–1.5 km altitude as stratus-like clouds, 1.5 km to the uppermost mixed-phase clouds as cumulus-like clouds, and all ice-only cirrus-like clouds. With cloud-fraction bins, this model is limited to 10^3 ICAs, independent of vertical resolution, and had a maximum of 288 ICAs in our tests.

Our best cloud overlap model is G6/.33 since it is based on the observed-modeled cloud correlations. For a given FCA, we treat the J values calculated by summing Fast-J over all the ICAs generated by G6/.33 as the correct value. We calculate errors for the other cloud-overlap models (here) or various ICA-approximation models using the G6/.33 model (Sect. 3). The errors in photolysis rates are calculated for different cloud overlap models by generating all the ICAs, using Fast-J to calculate J values, and computing the weighted sum of J 's. This study focuses on two J values that are critical in tropospheric chemistry and emphasize different wavelength ranges from near 300 nm, where O_3 absorption and molecular scattering are important, to 600 nm, where clouds are the predominant factor. $J-O^1D$ refers to the photolysis rate of $O_3 + h\nu \Rightarrow O_2 + O(^1D)$; and $J-NO_3$ includes both channels of the rate $NO_3 + h\nu \Rightarrow NO + O_2$ and $NO_2 + O$. We tested other key J values like those of HNO_3 and NO_2 , but found that their errors fell between the first two.

has smaller rms errors for the cases in Table 1, but the original MdQCA method may have a better profile for the mean error. Ran-3 is computationally comparable to the QCAs and has a reasonable mean bias, but the rms error is much worse, typically 10% more.

4 Cloud-J and volatile organic compounds

Volatile organic compounds (VOCs) cover a wide class of gaseous species containing C, H, O and sometimes N or S. They play a major role in the chemical reactivity of the troposphere, including production and loss of O₃ and loss of CH₄ (e.g., Jacob et al., 1993; Horowitz et al., 1998; Ito et al., 2009; Emmons et al., 2010), plus the formation of secondary organic aerosols (SOA, e.g., Ito et al., 2007; Fu et al., 2008; Galloway et al., 2011). For most VOCs (and H₂O₂) photolysis is the dominant loss, see Fig. 4. Daily photolysis rates (loss frequencies) range from 0.03 to 20 per day and some vary greatly with altitude. For 9 of the 14 species shown in Fig. 4, the photolysis rates are larger or comparable to the loss rates for reaction with OH (given in the legend). Thus, accurate calculation of their J values is important in atmospheric chemistry models.

VOCs present a particular problem for any photolysis code that averages over wavelength intervals. For most chemical species, cross sections including quantum yields are parameterized as a function of wavelength (ν) and temperature (T) (e.g., Atkinson et al., 2008; Sander et al., 2011). In this case, Fast-J calculates solar-flux-weighted, average cross sections for each wavelength bin (Wild et al., 2000; Bian and Prather, 2002). These tables are created for a set of fixed T s, and then the cross section used for each bin in each atmospheric layer is interpolated in T . Many VOCs have complex, pressure-dependent quantum yields (e.g., Blitz et al., 2006) that follow the Stern–Volmer formulation where photolysis cross sections (for dissociation) are a function of wavelength, temperature, and pressure (P), typically of the form $A(T, \nu)/(1 + B(T, \nu)P)$, where A and B can be rational polynomial functions of T and ν (see Sander et al., 2011). For most VOCs the pressure dependence changes across the wavelengths

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



chemical reactivity across cloudy and clear. Nevertheless, for species with photolysis rates that are less than the frequency at which clouds form and air is processed through them ($\sim 24 \text{ day}^{-1}$), the average J is the relevant quantity for chemistry modeling.

The next step would be to model at high-enough resolution so that air parcels are either cloudy or clear. This could resolve the 3-D correlation of clouds at scales of 1–4 km, which would in turn require a 3-D radiative transfer model (Norris et al., 2008; Davis and Marshak, 2010). A more interesting approach that is practical with typical global model resolution is the treatment of inhomogeneous cloud fields as being composed of independently scattering cloudlets (Petty, 2002). This cloudlet approximation could be readily integrated into the plane-parallel framework of Fast-J.

The added computational cost with G6/.33+AvQCA occurs with the additional calls to Fast-J, as the MAX-COR model and sorting of ICAs is fast. Computing photolysis rates 2.8 times per atmospheric column instead of once may add to the overall computational burden, but Fast-J is efficient and the costs should still be much less than the overall chemistry-solver and tracer-transport codes.

Code availability

The most recent version of Cloud-J and earlier versions of Fast-J can be found at <ftp://128.200.14.8/public/prather/fastJX/>. Cloud-J 7.3 as described here is included as a zip-file with the supplementary material of this publication. Subscribe to the maillist UCI-Fast-J@uci.edu or check the ftp site for updates on cross sections following new evaluations of photochemical data.

The Supplement related to this article is available online at [doi:10.5194/gmdd-8-4051-2015-supplement](https://doi.org/10.5194/gmdd-8-4051-2015-supplement).

GMDD

8, 4051–4073, 2015

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. Research at UCI was supported by NASA grant NNX13AL12G and DOE BER award DE-SC0012536.

References

- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., Troe, J., and Wallington, T. J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume IV – gas phase reactions of organic halogen species, *Atmos. Chem. Phys.*, 8, 4141–4496, doi:10.5194/acp-8-4141-2008, 2008.
- Bian, H. S. and Prather, M. J.: Fast-J2: accurate simulation of stratospheric photolysis in global chemical models, *J. Atmos. Chem.*, 41, 281–296, 2002.
- Blitz, M. A., Heard, D. E., and Pilling, M. J.: Study of acetone photodissociation over the wavelength range 248–330 nm: evidence of a mechanism involving both the singlet and triplet excited states, *J. Phys. Chem. A*, 110, 6742–6756, doi:10.1021/Jp056276g, 2006.
- Briegleb, B. P.: Delta-Eddington Approximation for solar-radiation in the NCAR Community Climate Model, *J. Geophys. Res.*, 97, 7603–7612, 1992.
- Davis, A. B. and Marshak, A.: Solar radiation transport in the cloudy atmosphere: a 3-D perspective on observations and climate impacts, *Rep. Prog. Phys.*, 73, 026801, doi:10.1088/0034-4885/73/2/026801, 2010.
- Emmons, L. K., Apel, E. C., Lamarque, J.-F., Hess, P. G., Avery, M., Blake, D., Brune, W., Campos, T., Crawford, J., DeCarlo, P. F., Hall, S., Heikes, B., Holloway, J., Jimenez, J. L., Knapp, D. J., Kok, G., Mena-Carrasco, M., Olson, J., O'Sullivan, D., Sachse, G., Walega, J., Weibring, P., Weinheimer, A., and Wiedinmyer, C.: Impact of Mexico City emissions on regional air quality from MOZART-4 simulations, *Atmos. Chem. Phys.*, 10, 6195–6212, doi:10.5194/acp-10-6195-2010, 2010.
- Feng, Y., Penner, J. E., Sillman, S., and Liu, X.: Effects of cloud overlap in photochemical models, *J. Geophys. Res.*, 109, D04310, doi:10.1029/2003JD004040, 2004.
- Fu, T. M., Jacob, D. J., Wittrock, F., Burrows, J. P., Vrekoussis, M., and Henze, D. K.: Global budgets of atmospheric glyoxal and methylglyoxal, and implications for formation of secondary organic aerosols, *J. Geophys. Res.*, 113, D15303, doi:10.1029/2007JD009505, 2008.

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Galloway, M. M., Loza, C. L., Chhabra, P. S., Chan, A. W. H., Yee, L. D., Seinfeld, J. H., and Keutsch, F. N.: Analysis of photochemical and dark glyoxal uptake: implications for SOA formation, *Geophys. Res. Lett.*, 38, L17811, doi:10.1029/2011GL048514, 2011.

Horowitz, L. W., Liang, J. Y., Gardner, G. M., and Jacob, D. J.: Export of reactive nitrogen from North America during summertime: sensitivity to hydrocarbon chemistry, *J. Geophys. Res.*, 103, 13451–13476, 1998.

Hsu, J. N. and Prather, M. J.: Is the residual vertical velocity a good proxy for stratosphere–troposphere exchange of ozone?, *Geophys. Res. Lett.*, 41, 9024–9032, doi:10.1002/2014gl061994, 2014.

Ito, A., Sillman, S., and Penner, J. E.: Effects of additional nonmethane volatile organic compounds, organic nitrates, and direct emissions of oxygenated organic species on global tropospheric chemistry, *J. Geophys. Res.*, 112, D06309, doi:10.1029/2005JD006556, 2007.

Ito, A., Sillman, S., and Penner, J. E.: Global chemical transport model study of ozone response to changes in chemical kinetics and biogenic volatile organic compounds emissions due to increasing temperatures: sensitivities to isoprene nitrate chemistry and grid resolution, *J. Geophys. Res.*, 114, D09301, doi:10.1029/2008jd011254, 2009.

Jacob, D. J., Logan, J. A., Yevich, R. M., Gardner, G. M., Spivakovsky, C. M., Wofsy, S. C., Munger, J. W., Sillman, S., Prather, M. J., Rodgers, M. O., Westberg, H., and Zimmerman, P. R.: Simulation of summertime ozone over North-America, *J. Geophys. Res.*, 98, 14797–14816, 1993.

Joseph, J. H., Wiscombe, W. J., and Weinman, J. A.: Delta-Eddington Approximation for radiative flux-transfer, *J. Atmos. Sci.*, 33, 2452–2459, 1976.

Kato, S., Sun-Mack, S., Miller, W. F., Rose, F. G., Chen, Y., Minnis, P., and Wielicki, B. A.: Relationships among cloud occurrence frequency, overlap, and effective thickness derived from CALIPSO and CloudSat merged cloud vertical profiles, *J. Geophys. Res.*, 115, D00H28, doi:10.1029/2009JD012277, 2010.

Naud, C. and DelGenio, A. D.: Cloud Overlap Dependence on Atmospheric Dynamics, Sixteenth ARM Science Team Meeting Proceedings, 27–31 March 2006, Albuquerque, NM, 2006.

Neu, J. L., Prather, M. J., and Penner, J. E.: Global atmospheric chemistry: integrating over fractional cloud cover, *J. Geophys. Res.*, 112, D11306, doi:10.1029/2006JD008007, 2007.

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Models for cloud overlap and approximation of ICAs including errors in J values.


Cloud overlap models to generate ICAs		ICAs ^a	avg err 0–1 km		rms err 0–1 km		rms err 0–16 km	
			J-O ¹ D	J-NO ₃	J-O ¹ D	J-NO ₃	J-O ¹ D	J-NO ₃
G0	MAX-RAN with MAX groups bounded by layers with CF = 0	19	+2%	+2%	21%	17%	6%	11%
G3	3 MAX-RAN groups split at 1 km and at the ice-only cloud level	21	+2%	+2%	15%	15%	5%	7%
G6/.00	6 MAX-COR groups, cc = 0.00	169	–1%	–1%	5%	4%	2%	3%
G6/.33	6 MAX-COR groups, cc = 0.33 ^b	169						
G6/.99	6 MAX-COR groups, cc = 0.99	169	+2%	+1%	11%	8%	4%	7%
Simple cloud models		ICAs						
CISky	clear sky, ignore clouds	1	+14%	+10%	24%	20%	14%	23%
AvCld	average fractional cloud across layer	1	–5%	+1%	11%	11%	8%	15%
CF3/2	increase CF to CF ^{3/2} and average over layer	1	+7%	+11%	10%	15%	5%	8%
ICA approximations		J calls						
AvDir	average direct beam from all ICAs	1	+5%	+11%	6%	13%	3%	7%
MdQCA	Quadrature Column Atmospheres uses mid-point in each QCA	2.8	+1%	0%	4%	4%	4%	5%
AvQCA	QCAs, uses average in each QCA	2.8	–1%	0%	3%	2%	2%	4%
Ran-3	Select 3 ICAs at random	3	+2%	+1%	12%	12%	9%	12%

^a Average number of ICAs for a tropical atmosphere, see Fig. 2.


^b Recommended cloud overlap model and reference model for calculation of errors.

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather



MAXimum overlap (connected layers become a MAX-Group)



RANdom overlap (between layers here, or between MAX-Groups)



CORrelated overlap, with $cc = \frac{1}{2}$ (between layers here, or between MAX-Groups)




Figure 1. Schematic of overlapping fractional-cloud layers. See text.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 2. Number of Independent Column Atmospheres (ICAs) generated by three different cloud overlap models (G0, G3, G6) from 640 different tropical fractionally cloudy atmospheres (FCAs) and sorted in order of increasing ICA number. The different correlation coefficients used in the G6 model do not change the number of ICAs, only their weights. The average number of ICAs per FCA is given in the legend. See text for definition of models.

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

[Title Page](#)

[Abstract](#) | [Introduction](#)

[Conclusions](#) | [References](#)

[Tables](#) | [Figures](#)

[◀](#) | [▶](#)

[◀](#) | [▶](#)

[Back](#) | [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Figure 3. Profile of the average bias in J value approximations relative to the J value calculated from the weighted average of all ICAs using model G6/.33. Values here are the average of 640 FCAs (108,125 ICAs) derived from the equatorial statistics (all longitudes) of cloud fraction, liquid water content, and ice water content from a snapshot of a T319L60 meteorology from the European Centre for Medium-range Weather Forecasts. Three simple cloud methods (dashed lines) do not use any cloud-overlap model, and three approximations for the ICAs (solid lines) use the G6/.33 model described here. The MdQCA ICA approximation was developed in Neu et al. (2007); the AvQCA and AvDir approximations are developed in this paper. The $J\text{-O}^1\text{D}$ refers to the photolysis rate of $\text{O}_3 + h\nu \Rightarrow \text{O}_2 + \text{O}^1\text{D}$, with average values of 4 ($z = 0$ km) to 9 ($z = 16$ km) $\times 10^{-5} \text{ s}^{-1}$; and $J\text{-NO}_3$, to all channels of the rate $\text{NO}_3 + h\nu \Rightarrow$, with average values of 2 to 4 $\times 10^{-1} \text{ s}^{-1}$. These two J values emphasize sunlight from 310 to 600 nm, respectively, and thus span the typical range of errors in tropospheric photolysis rates.

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 4. Volatile organic compounds (VOCs) and related species photolysis rates (day^{-1}) as a function of altitude (km). The complex structure with altitude is due to a combination of increasing UV-radiation with altitude and Stern–Volmer pressure dependences on quantum yields. Changes in slope occur at the interpolation points, temperature or pressure, of the cross sections. We assume that the noon-time J 's (clear-sky, tropical atmosphere, albedo = 0.10, SZA = 15°) apply for 8 of 24 h. Equivalent rates for OH loss are shown with the species name in the legend and assume a noontime OH density of $6 \times 10^6 \text{ cm}^{-3}$. Asterisks denote species for which photolysis loss is greater than or comparable to OH loss. VOC abbreviations are: MGlyxl = methyl glyoxal; Glyxl = glyoxal; PropAld = propionaldehyde; GlyAld = glycol aldehyde; ActAld = acetaldehyde; MEKeto = methylethyl ketone; MeVK = methylvinyl ketone; MeOOH = CH_3OOH ; MeAcr = methacrolein; MeNO₃ = methyl nitrate; PAN = peroxyacetyl nitrate.

Photolysis rates in correlated overlapping cloud fields: Cloud-J 7.3

M. J. Prather

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

