

Assessment of valley cold pools and clouds in a very high resolution NWP model.

J.K. Hughes¹, A.N. Ross¹, S.B. Vosper², A.P. Lock², and B.C. Jemmett-Smith¹

¹School of Earth and Environment, University of Leeds, UK

²Met Office, Exeter, UK

Correspondence to: J.K.Hughes
(J.K.Hughes@leeds.ac.uk)

Abstract. The formation of cold air pools in valleys under stable conditions represents an important challenge for numerical weather prediction (NWP). The challenge is increased when the valleys which dominate cold pool formation are on scales unresolved by NWP models, which can lead to substantial local errors in temperature forecasts. In this study a two-month simulation is presented using a nested model configuration with a finest horizontal grid spacing of 100m. The simulation is compared with observations from the recent COLPEX project and the model's ability to represent cold pool formation and the surface energy balance is assessed. The results reveal a bias in the model long-wave radiation which results from the assumptions made about the sub-grid variability in humidity in the cloud parametrization scheme. The cloud scheme assumes relative humidity thresholds below 100% to diagnose partial cloudiness, an approach common to schemes used in many other models. The biases in radiation, and resulting biases in screen temperature and cold pool properties are shown to be sensitive to the choice of critical relative humidity, suggesting that this is a key area which should be improved for very high resolution modelling.

1 Introduction

The stable boundary layer presents a difficult challenge for numerical simulation (?), ~~it being as~~ it is associated with a rich variety of complex dynamical processes. The dominant eddy scales are smaller than those typical of convective boundary layers, turbulence is weak and intermittent (?) and the presence of internal gravity waves can have an important influence on drag and mixing. In complex terrain the flows are further complicated by drainage flows and the formation of valley cold pools which can result in minimum temperatures which are significantly lower than those above the

surrounding higher terrain. Cold pools can lead to localized road icing and fog formation, presenting hazards to road users and frost damage to agriculture.

Previous studies of drainage flow and cold pools have tended to focus on large scale mountain valleys (e.g. ??) or isolated bowls and sinkholes (e.g. ??). However, small scale valleys are also of practical importance and these are often unresolved even in modern NWP regional models which have grid spacing of a few kilometers.

These small scale valleys can nonetheless lead to significant spatial temperature variations (?). The COLd air Pooling Experiment (COLPEX) was an investigation of the formation of cold air pools in such small-scale ~~UK~~ United Kingdom orography (?). It combined high resolution modelling and observations for a field campaign in the Clun valley, Shropshire, UK from January 2009 to April 2010. The Clun valley is 1–2 km wide with a depth of 100–200 m and as such is not resolved in the Met Office regional UK forecast model, either at the 4 km horizontal resolution used operationally at the time, or at the current 1.5 km operational resolution. Cold air pools in this region can lead to local temperature differences of 5–10 K between hill top and valley bottom. Details of the observational campaign are described in ?. ? analysed the observed cold pool structure and related the frequency and strength of cold pools to previous idealised studies of cold pool formation (?). Modelling work has used a nested version of the Met Office Unified Model (MetUM), with a horizontal grid spacing of 100 m. Over the lowest 1km the vertical resolution is on average 27m. The basic model configuration and initial comparison with observations is given by ?. The results show that the high horizontal and vertical resolution is required in order to sufficiently resolve the orography and obtain a good level of agreement between the simulated and observed temperature variations across the cold pools in clear-sky cases. Subsequently, ? focussed in detail on a single Intensive Observation Period (IOP) which took place on the night of 04 March, 2010 (IOP 16). The latter study demonstrated the skill of the MetUM in reproducing the observed temperatures across the valley and provided a detailed analysis of the heat budget within the cold pool.

Previous modelling studies in COLPEX, and in other field campaigns, have used a case study approach, simulating short (2–3 day) events where cold air pools are known to form. The current study takes a ~~different~~ longer simulation period approach with a two-month simulation at 100 m resolution in order to provide a longer term assessment of cold pool formation over the COLPEX region. This allows a more systematic and objective validation of the model representation of cold pools over a range of conditions. It also allows a quantification of the frequency, strength and drivers of cold air pools. Given the density of observations obtained in COLPEX this provides a good opportunity to validate the behaviour of the MetUM at high resolution over a range of conditions. The simulation also allows an investigation of the importance of spinning up the high-resolution model and can also be used to provide a more complete analysis of the synoptic scale influences on cold pool formation.

Two months is not sufficiently long to generate a true climatology, however the computational resources required to undertake a multi year simulation at this resolution are beyond what was available for this project. The current study provides a useful intermediate step towards this. ~~Although not truly a climatology, we will refer to the two month simulation as a “climatology simulation” to contrast it from shorter, 1-2 day long case study simulations.~~

In the following sections the model configuration, the simulations and forcing data are described. Following this a discussion of the importance of model spin up is presented. From the ~~climatological~~ long term simulation an analysis of the temperature bias relative to observations is then given. Based on this analysis, an additional simulation is then presented and analysed. A comparison of simulated and observed cloud cover is then made with reference to unresolved humidity variability. The paper concludes with a more general discussion of the results.

2 Two month simulation over the COLPEX region

2.1 Model setup

The COLPEX simulations were conducted using the MetUM Model with a double-nested setup, with simulations running with horizontal grid spacings of 4 km, 1.5 km and 100 m (hereafter referred to as $\Delta 4\text{km}$, $\Delta 1.5\text{km}$ and $\Delta 100\text{m}$ models). ? describe the modelling setup in detail so only a brief summary is given here. The innermost $\Delta 100\text{m}$ domain uses a non-uniform horizontal grid with the resolution decreasing from 100 m to 1.5 km at the boundaries (?). The domain covers a region of 80 km by 80 km, centred on the Clun Valley, Shropshire (UK) with a 30 km by 30 km inner domain of constant 100 m resolution. Figure 1 shows the innermost ~~part~~ 10km by 10km of the domain including the Clun valley as represented in the 100 m simulations. Analysis data from the Met Office three-dimensional variational assimilation scheme (3D-VAR) is used to initialise the 4 km model and updates the $\Delta 4\text{km}$ state every three hours. The $\Delta 4\text{km}$ simulation then produces the lateral boundary conditions for $\Delta 1.5\text{km}$ which in turn produces lateral boundary conditions for the $\Delta 100\text{m}$ simulation. The nesting is one-way, with no feedback to lower resolutions. $\Delta 1.5\text{km}$ and $\Delta 100\text{m}$ simulations are ‘free-running’, i.e. are not re-initialised during the simulation. This allows aspects such as soil moisture to develop spatial patterns consistent with the high resolution terrain data during the simulation rather than being constrained by relatively coarse analysis fields. The simulations presented here are all initialised at 12 UTC, allowing the model to spin up prior to the onset of stable conditions in the evening.

Other details of the model setup are broadly similar between resolutions, however there are a few key differences. Firstly, in the $\Delta 100\text{m}$ simulations the number of vertical levels is increased from 70 to 140. Secondly, different turbulence parametrisations are used at different resolutions. The $\Delta 4\text{km}$ model uses a 1D boundary-layer scheme, while the $\Delta 1.5\text{km}$ uses a 2D Smagorinsky turbulence parameterisation for horizontal mixing and a 1D boundary layer scheme for vertical mixing, and the




Figure 1: Orography of the Clun valley and surrounding region, as represented in the COLPEX $\Delta 100\text{m}$ simulations. The main observational tower sites are marked as squares and labelled. Hobo stations are marked with circles (see ?). The main sites are Duffryn (D; main valley, height 243m), Springhill (S; local ridgetop, height 404m) and Burfield (B; secondary valley, height 303m). It should be noted that the full domain of the $\Delta 100\text{m}$ simulation is more extensive than is shown here.

$\Delta 100\text{m}$ model uses a 3D Smagorinsky scheme (?). Finally, the vertical profile of critical relative humidity (RH_{crit}) necessary for cloud formation is increased, to take into account the reduction in grid box size (see ?). The critical relative humidity threshold approach allows non-zero cloud cover to be diagnosed when grid box relative humidity is below saturation. This assumes that there is unresolved sub-grid scale variability in temperature and moisture which results in regions of cloud cover. This approach to sub-grid scale variability was first proposed by ? and ?. It was then developed into the form used here by ?. The critical relative humidity is a function of the standard deviation of sub-grid scale variability, such that when temperature and moisture variability are fully resolved the standard deviation of sub-grid scale variability is zero and RH_{crit} is 100%. For the $\Delta 1.5\text{km}$ model the vertical profile of RH_{crit} is set to 91% at the surface, reducing to 80% at 845 m and constant at 80% above. At $\Delta 100\text{m}$ this profile is set to 99% in the lowest 500 m, decreasing linearly to 91% at 3.4 km and constant (90%) above this level. These profiles are shown in figure 2.

2.1.1 The simulations

The COLPEX observational field campaign ran between January 2009 and April 2010 (?). A number of short simulations have been completed, corresponding to IOPs, some of which have been the




Figure 2: Vertical profiles of RH_{crit} used in the different nested models. The $\Delta 100m$ and $\Delta 1.5km$ profiles are shown in black and red, respectively.

focus of previous analysis (??, ?). In addition to these, an extended simulation has been completed, for the period 17 August to 15 October 2009 (the climatology-long term simulation). This period was chosen because it includes multiple IOP cases, two of which (09–10 September, IOPs 4 and 5) have already been investigated in detail (??). The IOPs during this period include cases where the model reproduces the observations well (e.g. IOPs 4 and 5), and those where the model performs less well (16–17 September, IOP 6 and 7). In terms of investigating the mechanisms involved in cold pool formation it is appropriate to focus on IOPs where the model performs well, however when the more general performance of the model is of interest it is important to consider a wider range of conditions over the whole two month period.

2.2 Continuity of 4 km analysis forcing data

The COLPEX simulations are reliant on archived operational 4 km UK analyses data from the Met Office 3D-VAR assimilation scheme. Unfortunately during 2009-2010 there are occasional instances

of missing data from the archive. During the ~~climatology~~long term simulation there were missing
120 data during 05-07 September (18-14 UTC), 26-28 September (00-14 UTC) and on 04 October (09
to 14 UTC). It was beyond the resources of the current project to recreate the missing data. It was
therefore necessary to understand and minimize the impact of the data gaps.

In order to retain as much of the system memory as possible, when the model is restarted after a pe-
riod of missing forcing data the soil properties (soil moisture content and temperature) are initialised
125 from the model state at the last equivalent time of day prior to the missing data. For example, when
restarting the simulation at 12 UTC on 29 September the soil properties are initialised with those at
12 UTC on 25 September. Other, faster components of the simulation are simply downscaled, i.e. air
temperature in the $\Delta 100\text{m}$ simulation at 12 UTC 29 September, is re-initialized with the $\Delta 4\text{km}$ air
temperature. In tests this approach was shown to minimize the impact of the data gaps on the model
130 evolution. To test this a period without data gaps was chosen. Temperature and moisture timeseries
at the main observation sites of Spring Hill(hill-top) and Duffryn (bottom of the Clun valley) from
the long $\Delta 100\text{m}$ simulation were compared with equivalent timeseries from simulations where the
model was initialized using either the $\Delta 100\text{m}$ model state from 48 hours previously, or by interpo-
lating the $\Delta 4\text{km}$ solution to the 100 m grid. This analysis (not included here for brevity) led to the
135 above approach.

2.3 Impact of spinning up the model state

One of the motivations for completing the ~~climatology~~long term simulation is to quantify the im-
portance of spinning up slower components of the model state (e.g. ~~the soil properties~~soil moisture
and temperature) within the $\Delta 100\text{m}$ simulation. In the context of these simulations, spin up is the
140 process of integrating the model forwards in time to reduce the impact of uncertainty in the initial
conditions and to allow different components of the simulation to reach a consistent state. Com-
pleting multi-month simulations with this setup is currently very computationally expensive and the
benefits of avoiding this are substantial. It is important to note, however, that soil properties in the
shorter simulations are already spun up at the $\Delta 4\text{km}$ resolution. In downscaling to the $\Delta 100\text{m}$ sim-
145 ulation the soil temperatures are interpolated onto the $\Delta 100\text{m}$ grid and then adjusted relative to the
height difference between orographic heights in the different resolutions using an assumed lapse rate
of 6 K km^{-1} . This lapse rate was chosen from an analysis of soil temperature and orographic height
variability in the $\Delta 4\text{km}$ output. Soil moisture is simply interpolated onto the $\Delta 100\text{m}$ grid.

Figure 3 shows the $\Delta 100\text{m}$ domain-averaged soil temperatures throughout the ~~climatology~~long
150 term run. The top soil cools throughout August despite warm synoptic conditions (mean temperature
at Clun was 292.5 K ; ?, 91-92). At the start of September there is a ~~significant~~-warming trend (just
prior to the first missing data period), before resuming the cooling trend. September was dominated
by anticyclonic conditions with cyclonic conditions developing in October (?, 92). The lowest soil
level shows an underlying trend which results from the extremely slow response time of this level.

Figure 4 shows the domain-average soil moisture content throughout the ~~climatology-long term~~ simulation on each soil level. Figures 3 and 4 show that there exists long term variability in the soil properties.

During the ~~climatology-long term~~ simulation there are three pairs of IOPs: 09 and 10 September (IOP 4 and IOP 5), 16 and 17 September (IOP 6 and IOP 7), and 13 and 14 October (IOP 8 and IOP 9). Markers on figures 3 and 4 show domain-averaged soil temperature and moisture content from the IOP simulations. The initial domain-mean IOP soil temperatures are close to those in the ~~climatology-long term~~ simulation. Figure 4 shows some difference in soil moisture between the IOP and ~~climatology-long term~~ simulations, which is most pronounced after the ~~moist-precipitation~~ event at the start of September. This suggests that there are differences in the hydrological cycle between the two resolutions since in the IOP simulations the soil moisture content remains similar to the downscaled $\Delta 4\text{km}$ content throughout the simulation.

Figure 5 shows a comparison between the screen-level air temperature and specific humidity at the Duffryn site in the IOP and ~~climatology-long term~~ simulations for the three IOP pairs. Figure 5 shows that in general the ~~climatology-long term~~ simulation is not required in order to simulate the IOP cases, i.e. that the atmosphere adjusts relatively rapidly ~~and that the temperature and humidity fields are not sensitive to the difference between soil properties~~ in the $\Delta 4\text{km}$ and $\Delta 100\text{m}$ simulations. This supports the validity of earlier COLPEX work (e.g. ?) and also suggests that the missing data periods in the ~~climatology-long term~~ simulation do not undermine the analysis presented here.

3 Model temperature bias

Figure 6 shows the frequency distribution of biases in the daily minimum and maximum model screen-level (1.5 m) air temperature from the ~~climatology-long term~~ simulation at the main observation sites, Duffryn (valley site) and Spring Hill (hill top site; see figure 1). All data are hourly averaged and differences are defined as model minus observed. Results from $\Delta 100\text{m}$ and $\Delta 1.5\text{km}$ resolutions are shown. Figure 6a shows that for $\Delta 100\text{m}$ the daily minimum temperature at Duffryn shows a stronger cold bias than at Spring Hill, i.e. at Duffryn there is a higher number of days where the minimum temperature is more than 3 K lower than observed. The mean bias at Duffryn is -1.6 ± 0.3 K and at Spring Hill it is -0.7 ± 0.2 K, where all errors stated are calculated as the standard error of the mean. For $\Delta 1.5\text{km}$ there is an apparent warm bias of 1.4 ± 0.3 K at Duffryn and 0.4 ± 0.2 K at Spring Hill. This warm bias at Duffryn is consistent with the fact that the valley is poorly resolved in the $\Delta 1.5\text{km}$ simulation.

Figure 6b shows that the simulated daily maximum temperatures have a general cold bias. For $\Delta 100\text{m}$ this cold bias is -0.6 ± 0.3 K at Duffryn and -0.9 ± 0.3 K at Spring Hill. In the $\Delta 1.5\text{km}$ simulation the mean biases are -1.1 ± 0.1 K at Duffryn and -0.1 ± 0.1 K at Spring Hill. If the Spring Hill temperatures are representative of temperatures outside the valley this suggests that there is a




Figure 3: Evolution of the domain mean soil temperature. ~~Day number corresponds to~~, in the day since the start of the climatology $\Delta 100\text{m}$ simulation. The soil levels shown are 0.05 m (black), 0.225 m (green), 0.675 m (dark blue) and 2 m (light blue). Shaded circles show the values from corresponding IOP simulations. Vertical dashed lines mark 01 September and 01 October.

190 widespread cold bias in the $\Delta 100\text{m}$ simulation that is not caused by bias in the lateral boundary conditions or present in the lower resolution simulations. The cold bias in $\Delta 1.5\text{km}$ at Duffryn is most likely due to unresolved sheltering effects, which would give rise to local daytime warming within the valley.

The difference between hill top (Spring Hill) and valley bottom (Duffryn) screen temperature gives a measure of the strength of cold air pooling in the Clun valley [?](#). Using this difference, figure 6c shows the frequency distribution of daily maximum cold pool strength in the $\Delta 100\text{m}$ and $\Delta 1.5\text{km}$ simulations and in the observations. [It should be noted that this analysis includes all data from the simulation period, including stable and unstable conditions.](#) For $\Delta 100\text{m}$ the model simulates more strong cold pool events than observed. The simulated mean $\Delta 100\text{m}$ cold pool strength is 2.4 ± 0.3 K, compared to an observed mean strength of 1.6 ± 0.3 K. Figures 6a and 6b show this difference is due to the stronger cold bias at Duffryn. For $\Delta 1.5\text{km}$ the valley is not properly resolved and the apparent cold pool strength is typically close to zero; the mean is 0.4 ± 0.1 K.




Figure 4: Evolution of the domain mean layer soil moisture content at different levels. Day number corresponds to, in the day since the start of the climatology $\Delta 100\text{m}$ simulation. The soil levels shown are 0.05 m (black), 0.225 m (green), 0.675 m (dark blue) and 2 m (light blue). Shaded circles show the values from corresponding IOP simulations. Panels a-d correspond to increasing soil depths of 0.05m, 0.225m, 0.675m and 2.0m. Vertical dashed lines mark 01 September and 01 October.

3.1 Daily minimum temperature bias

Figure 7 shows the distribution of $\Delta 100\text{m}$ model daily minimum temperatures against observations at Duffryn. The strongest cold biases exist during relatively warm nights, whilst there is good agreement between model and observations during the coldest nights. The coldest nights are clear sky cases, suggesting that perhaps errors in the simulated cloud cover may contribute to the cold bias seen at higher temperatures. Previous COLPEX studies have focused on clear sky cases, since this is a prerequisite for strong cold pool formation (e.g. ?) however figure 7 shows that a more general evaluation of the modelling setup should include other cases.

Whilst there were no in-situ direct measurements of cloud properties during this time at Duffryn or Spring Hill, downward longwave radiative flux is available and directly quantifies the impact of




Figure 5: Difference between shorter IOP simulations and the longer ~~climatology term~~ simulation (IOP minus ~~climatology~~long term simulation) for screen temperature (a,c,e) and relative percentage differences in screen-level specific humidity (b,d,f) at Duffryn for IOPs 4-5 (a,b), IOPs 6-7 (c,d), IOPs 8-9 (e,f).

cloud on the nocturnal energy balance. Figure 8 shows the agreement between simulated ($\Delta 100\text{m}$) and observed longwave radiative flux at Spring Hill. The relationship to the temperature bias is
 215 illustrated by colouring the markers based on the observed hourly temperature bias. Each panel shows all observations from the 2-month simulation in the daily 3-hour window, with each circle showing a single hourly comparison. At both Spring Hill and Duffryn (not shown) sites there are a substantial number of days where downward longwave radiation is under predicted. This bias develops during the evening and persists through the morning. The size of the bias in longwave
 220 radiation is similar at both sites, with a mean daily bias of $-21.7 \pm 2.7 \text{ Wm}^{-2}$ at Duffryn and $-23.3 \pm 3.2 \text{ Wm}^{-2}$ at Spring Hill. The bias is smallest in the period 13-16 UTC at both sites when there is a mean model bias of $-6.0 \pm 2.9 \text{ Wm}^{-2}$. There is a systematic increase in this bias during the night and morning, increasing from $-24.4 \pm 2.9 \text{ Wm}^{-2}$ (19-22 UTC) to $-33.7 \pm 2.7 \text{ Wm}^{-2}$ (07-10 UTC). There is a substantial cold bias in all time periods in figure 8, however figure 8h in particular
 225 suggests that this cold bias is associated with the under predicted downwelling longwave radiation.




Figure 6: Bias in the model screen-level (1.5 m) temperatures (model minus observed). a) Frequency distributions of daily ~~hourly~~ minimum temperature bias compared to observations at Spring Hill and Duffryn, b) Frequency distributions of daily ~~hourly~~ maximum temperature bias, c) Frequency distributions of cold pool strength, defined as the maximum daily difference between Spring Hill and Duffryn. The data are binned into intervals of width 0.5 K. Daily minimum and maximum temperatures are calculated from hourly average temperatures.

In the $\Delta 1.5\text{km}$ simulation the bias in downward longwave flux is smaller than seen in the $\Delta 100\text{m}$ model, the mean being $-10.1 \pm 2.4 \text{ Wm}^{-2}$, at Duffryn a reduction of 53% compared to the $\Delta 100\text{m}$ model. At Spring Hill this value is $-7.2 \pm 2.4 \text{ Wm}^{-2}$ (a reduction of 69%).

At this point we conclude that the daily minimum temperature in the $\Delta 100\text{m}$ simulation has a systematic cold bias, most likely caused by too little cloud cover in the $\Delta 100\text{m}$ simulation itself (in contrast to biases introduced via transport through the lateral boundaries). The fact that Duffryn exhibits a greater cold bias than Spring Hill is most likely due to the effects of resolved cold pool formation, which exacerbate the bias.

3.2 Daily maximum temperature

During the daytime shortwave radiation dominates the surface energy budget. Figure 9 shows the agreement between the simulated and observed downward shortwave radiative flux at Spring Hill,




Figure 7: Simulated daily minimum $\Delta 100\text{m}$ screen temperatures versus observations at Duffryn.




Figure 8: Bias in $\Delta 100\text{m}$ simulated downward longwave radiation and its relationship to bias in the screen level temperature, at Spring Hill. The temperature bias is represented as the colour and size of the marker (simulated - observed). Each marker represents mean hourly values.




Figure 9: Bias in $\Delta 100\text{m}$ simulated downward shortwave radiation and its relationship to bias in the screen level temperature, at Spring Hill. The temperature bias is represented as the colour and size of the marker (simulated - observed).

combined with the screen temperature bias in the $\Delta 100\text{m}$ model. As shown previously, the simulated mean daily maximum temperature exhibits a cold bias. Figure 9 is consistent with this. Averaged between both stations, 89% of data show a cold bias at the start of the day (07–10 UTC). This fraction decreases throughout the day, dropping to 71% by 16–19 UTC. In contrast to this cold bias, downward shortwave radiation is over predicted on average, consistent with the reduction in frequency of cold bias data points. In the early morning and late evening (07–10 UTC and 16–19 UTC) the mean bias in shortwave radiation is $14.4 \pm 5.0 \text{ Wm}^{-2}$. Between 10 and 16 UTC the mean bias is 36.6 Wm^{-2} . During the afternoon, strong warm bias points (defined as those for which the model temperature is 1 K or more higher than observed) are associated with an over prediction of downwelling shortwave radiation, whilst strong cold bias points (bias $\leq -1 \text{ K}$) are typically associated with under predicted downwelling shortwave radiation, though in general cold bias points are evenly distributed between over and under predicted shortwave radiation. It seems likely, therefore that the cold bias in the daily minimum temperatures is the cause of the cold bias in daily maximum temperatures. It was hypothesised earlier that this is due to under prediction of cloud cover, and the observed over prediction of shortwave radiation also indicates too little cloud cover in the model at $\Delta 100\text{m}$.

4 Impact of the RH_{crit} profile

The above analysis suggests that a cloud cover bias affects the general ability of the model to accurately simulate the observed temperatures. This is now verified by performing an additional simulation using a modified version of the $\Delta 100m$ model setup.

As described earlier, although there are only a minimal set of differences between the $\Delta 100m$ and $\Delta 1.5km$ configurations one difference is the prescribed vertical profile of the relative humidity threshold, RH_{crit} , above which cloud forms. The values are higher in the $\Delta 100m$ configuration, based on the assumption that there will be less sub-grid scale variability in humidity at higher resolution, since at sufficiently high resolution variability in temperature and humidity would be resolved. In order to test the sensitivity of the simulation to cloud cover the prescribed vertical RH_{crit} profile in the $\Delta 100m$ simulation was replaced with a vertical profile equivalent to the $\Delta 1.5km$ profile, with values interpolated onto the $\Delta 100m$ vertical levels. The modified $\Delta 100m$ setup (hereafter referred to as $\Delta 100m_r$) is then tested in an additional simulation.

Figure 8 showed that a bias in the $\Delta 100m$ downward longwave radiation exists throughout the night. This bias is pronounced during 15-25 September and is absent at coarser resolution ($\Delta 1.5km$) over the same period, suggesting it is an internal feature of the $\Delta 100m$ simulation. This period also includes two IOPs (16-17 September; IOPs 6 and 7), allowing for a detailed comparison with observations, and so was chosen for a test of the $\Delta 100m_r$ configuration.

Figure 10 shows simulated mean vertical profiles of cloud fraction over the lowest 1 km for the $\Delta 1.5km$, $\Delta 100m$ and $\Delta 100m_r$ simulations. As expected, comparing figures 10a and 10b shows a large decrease in cloud cover between the $\Delta 1.5km$ and original $\Delta 100m$ simulation. This decrease in cloud cover is largely due to the change in RH_{crit} profile, since reverting the RH_{crit} profile at $\Delta 100m$ to the $\Delta 1.5km$ profile results in cloud cover fractions similar to $\Delta 1.5km$ (figure 10c).

Figure 11a shows the vertical cloud profile averaged over the re-run period and averaged spatially over the inner domain of the $\Delta 100m$ (i.e. the central domain of the simulation with constant 100 m grid spacing). Figures 11b and 11c show equivalent mean cloud profiles for day (07-19 UTC) and night (19-07 UTC) periods separately. In figure 11a the cloud cover over the lowest 100 m is similar in the $\Delta 100m$ and $\Delta 100m_r$ simulations but above this the $\Delta 100m_r$ cloud is close to that in the $\Delta 1.5km$ model. When the cloud profiles for day and night are separated, more substantial differences become apparent between $\Delta 1.5km$ and $\Delta 100m_r$. During the day the $\Delta 100m_r$ cloud fraction is close to that of $\Delta 1.5km$ whereas the original $\Delta 100m$ cloud cover is consistently smaller. During the night $\Delta 100m_r$ produces more cloud over the lowest 500 m than $\Delta 1.5km$ and $\Delta 100m$. Above 500 m the $\Delta 100m_r$ cloud is more similar to that of $\Delta 1.5km$ than $\Delta 100m$. The diurnal variability in near surface air conditions is larger in $\Delta 100m$ than $\Delta 1.5km$ during 15-25 September. The domain and time averaged $\Delta 100m$ near surface air has lower night temperatures, increased dew deposition and lower specific humidities than $\Delta 1.5km$. The cooler temperatures dominate, leading to the observed increased cloud cover at $\Delta 100m$ irrespective of the RH_{crit} profile used (figure 11c).




Figure 10: Hourly area averaged profiles of model cloud fraction (F) for a) $\Delta 1.5\text{km}$, b) $\Delta 100\text{m}$ and c) $\Delta 100\text{m}_r$. Vertical lines show 00 UTC 16 September and 00 UTC 19 September, and correspond to the period shown in figure 13. Area averaged are calculated over the 10km by 10km domain illustrated in figure 1.

Comparing the downward longwave radiation in the $\Delta 100\text{m}$ and $\Delta 100\text{m}_r$ simulations it is evident that the bias is reduced by reducing RH_{crit} , but not eliminated. The $\Delta 100\text{m}$ downward longwave radiation has a bias of $-59.4 \pm 17.3 \text{ Wm}^{-2}$ averaged over 15-25 September and at both sites. For $\Delta 100\text{m}_r$, this bias is $-43.1 \pm 18.1 \text{ Wm}^{-2}$: a reduction of 27 %. Both Duffryn and Spring Hill show similar biases. For the same period, the $\Delta 1.5\text{km}$ downward longwave radiation has a similar bias to the $\Delta 100\text{m}_r$ output, with a mean bias of $-44.2 \pm 18.1 \text{ Wm}^{-2}$, suggesting that there is a residual bias, perhaps connected with errors in the 4 km driving model over this period. Note that 15-25 September was chosen as a period during which there is substantial bias in the simulation. The model longwave bias of $\Delta 1.5\text{km}$ averaged over the whole of the climatology long term simulation period is -8.7 Wm^{-2} (see section 3.1).




Figure 11: The mean cloud cover profile for the lowest kilometre for a) all hours, b) 07-19 UTC and c) 19-07 UTC, averaged over all days in the re-run period (15-25 September).

During the re-run period downward shortwave radiation in the $\Delta 100\text{m}$ simulation is over predicted
 300 by $59.1 \pm 17.4 \text{ Wm}^{-2}$ (considering all data between 0700 and 1859 hours at both sites). In the $\Delta 100\text{m}_r$ simulation this bias is reduced to $11.2 \pm 15.8 \text{ Wm}^{-2}$, again suggesting improved cloud cover.

In general, screen-level temperatures agree more closely with observed temperatures in the $\Delta 100\text{m}_r$ simulation than the $\Delta 100\text{m}$ simulation. Over the re-run period the mean temperature bias is reduced
 305 from -1.1 K to -0.3 K at Duffryn and from -0.9 K to -0.5 K at Spring Hill. Daily maximum temperatures are within 0.5 K of observed maxima during this period (for both RH_{crit} profiles) and do not systematically improve with the updated profile (the bias at Duffryn changes from 0.3 K to 0.1 K whilst at Spring Hill the bias changes from -0.02 K to -0.3 K). Daily minimum temperatures do systematically improve, with bias at Duffryn changing from -2.3K to -0.2K and the bias at Spring
 310 Hill changing from -1.5K to -0.7K .

The impact of these changes on simulated cold pool strength is shown in figure 12, which illustrates how changing the RH_{crit} profile reduces the number of nights in which cold pools develop, more closely matching the observed cold pool variability over this period.




Figure 12: Impact of RH_{crit} profile change on the simulated cold pool strength, defined as the difference between screen temperature between Spring Hill and Duffryn (Spring Hill minus Duffryn). Positive values indicate the presence of cold air pools. a) $\Delta 100m$ cold pool strength, b) $\Delta 100m_r$ cold pool strength, c) observed cold pool strength.

Since the re-run period was chosen to include two IOP cases (16-17 September; IOP 6 and 7) it is also possible to compare simulated cloud profiles against radiosonde profile data. Figure 13 compares $\Delta 1.5km$ model cloud cover profiles with radiosonde profiles at Spring Hill. Radiosondes were released hourly from Spring Hill and Duffryn during the evenings of 16 and 17 September. Both nights show high relative humidity at heights between 500 m and 1 km, in particular during the night of the 17 September when the air is near saturated between 700 m and 1 km throughout the night. This is in reasonable agreement with cloud cover in the $\Delta 1.5km$ simulation and figures 10 and 13 demonstrate that not only does the updated RH_{crit} profile result in cloud cover at 100 m resolution which is similar to that in the $\Delta 1.5km$ simulation, it also produces cloud cover which is in better agreement with observations.

As discussed earlier, the model critical relative humidity threshold is designed to account for unresolved variability in the humidity and temperature fields. It is therefore reasonable to expect that as resolution increases, the fraction of resolved variability increases and the appropriate RH_{crit} increases towards 100% (variability fully resolved). In order to quantify this, variability was calculated as the spatial standard deviation of relative humidity at each model height at each hour during the 10 day re-run period. These profiles were then time-averaged. Figure 14 shows vertical profiles of relative humidity standard deviation. Also included in figure 14 are profiles for $\Delta 100m$ model




Figure 13: Radiosonde relative humidity profiles at Spring Hill during 16-18 September. Also shown are $\Delta 1.5\text{km}$ cloud fractions above 20% (diagonal hatching).

output which prior to analysis have been horizontally averaged to 1.5km resolution. Away from the land surface RH standard deviation is greatest in $\Delta 100\text{m}$ as expected, and similar in both $\Delta 1.5\text{km}$ and the reduced resolution $\Delta 100\text{m}$ data. However, figure 14 also shows that close to the land surface differences in the surface energy balance between the model resolutions dominate and result in higher RH standard deviation in $\Delta 1.5\text{km}$ than the higher resolution simulation. This is particularly pronounced during nights. As was noted earlier there are differences in the domain-averaged near surface conditions between model versions and $\Delta 100\text{m}$ is cooler and drier at night than $\Delta 1.5\text{km}$, resulting in lower RH variability in $\Delta 100\text{m}$ than $\Delta 1.5\text{km}$.

5 Conclusions

A high resolution, two month simulation of the flow in Clun valley has been produced, allowing an analysis of the base model climatology as well as producing a robust assessment of the ability of the MetUM to capture the variability and strength of cold pool formation. By comparing a two month




Figure 14: Standard deviations of relative humidity values simulated in, $\Delta 100\text{m}$ output (black), $\Delta 100\text{m}$ output averaged to 1.5km resolution (red) and $\Delta 1.5\text{km}$ output (green). Solid lines are simulations with RHcrit profiles from the operational 1.5km model, while dashed lines correspond to the RHcrit profile with higher thresholds described in ?. The subfigures show (a) total mean profiles , (b) average day profiles (07-19 UTC) and (c) night profiles (19-07 UTC).

simulation to a series of shorter, IOP, simulations it was demonstrated that short simulations provide accurate results without the requirement to spin up slowly evolving components of the model at the highest resolution [though this may not hold true for other regions](#). Within the context of COLPEX this is an important result as it validates the approach taken in previous studies which have only made use of shorter simulations. It is a useful result in general since it supports the scientific validity of future high resolution simulations, without the requirement for computationally expensive extended simulations.

In addition to testing the importance of extended spin up times, the longer simulation also allowed an assessment of the model climatology. Daily minimum temperatures were shown to have a cold

bias in the highest resolution simulations relative to observations. This bias was then demonstrated to be due to choices made within the cloud scheme about sub-grid humidity variability. Compared to the 1.5 km resolution simulation ($\Delta 1.5\text{km}$), the original 100 m simulation ($\Delta 100\text{m}$) predicted less cloud cover. $\Delta 1.5\text{km}$ also compared better with radiosonde observations made during an IOP.

As a sensitivity test a period in the ~~climatology~~ long term simulation was re-run, during which the bias in downward longwave radiation at $\Delta 100\text{m}$ was most pronounced. For this simulation ($\Delta 100\text{m}_r$) the vertical profile of RH_{crit} at 100 m resolution used the 1.5 km model profile, i.e. no change in sub-grid scale variability between resolutions was assumed. This reduced the differences in model cloud cover between the resolutions. The bias in the downward longwave radiative flux in the $\Delta 100\text{m}_r$ simulations was also reduced relative to the observations and this resulted in a reduction in the cold bias of daily minimum temperature and an improved simulation of cold pool variability. This demonstrates the importance of considerations of sub-grid-scale variability for accurate simulation at very high resolution. The conclusions presented here are based on simulations at a single location and require further investigation before drawing robust, general conclusions. The analysis has also focused on horizontal resolution, however ? present results from 70,140 and 178 vertical level configurations of the model and show that the COLPEX simulations do exhibit a sensitivity to vertical resolution. When the $\Delta 100\text{m}$ model was run with 70 vertical levels (without changing RH_{crit}) as an additional sensitivity test, there was a 16% reduction in cloud cover, suggesting the current conclusions are also sensitive to the vertical resolution.

Two clear sky case IOPs, 09 September (IOP 4) and 04 March (IOP 16) have been analysed in previous studies (?, ?) and it is important to note that when simulations of these IOPs were re-run with the $\Delta 100\text{m}_r$ setup there was negligible impact on the results, since both model and observations indicate clear sky conditions. The original $\Delta 100\text{m}$ model output accurately predicts the observed temperature in these IOPs, consistent with the analysis presented here which shows that the model is well formulated for clear-sky nights. Numerical weather prediction (NWP) models are increasingly being run at resolutions more traditionally associated with large eddy simulations. Moving into this regime presents a number of challenges since at horizontal grid spacing of 100-1000m models will only partially resolve eddies (??), a domain referred to as the grey zone (?). In the convective boundary layer, the dominant eddy length scales are typically around 1km in size, approaching the depth of the boundary layer and therefore most likely resolved by the current modelling setup (?). However stable boundary layer eddies have smaller dominant eddy scales (?) and will be poorly resolved at 100m resolution. ? for example, completed a range of simulations varying the model resolution, demonstrating this sensitivity to the atmospheric stability. Investigating the resolution sensitivity of large eddy simulations of the stable boundary layer, ? showed a convergence of model behaviour only as horizontal resolution approached 2m.

Interestingly, in contrast to the results of our study ? investigated stratocumulus simulations in the grey zone and found relatively little sensitivity to the choice of RH_{crit} . This discrepancy suggests

the application of a more sophisticated approach to representing unresolved variability in saturation,
390 in which the assumed sub-grid variability is not static but instead varies according to properties of
the flow. ? for example developed such a scheme with a dynamic representation of the unresolved
PDFs. Such an approach has the potential to circumvent the need to specify RH_{crit} values and may
be more appropriate.

Acknowledgements. We would like to thank everyone involved in the COLPEX field campaign. John Hughes'
395 contribution was funded by the UK Natural Environment Research Council and UK Met Office through the Joint
Weather and Climate Research Programme under grant NE/I007679/1. We acknowledge use of the MONSooN
system, a collaborative facility supplied under the Joint Weather and Climate Research Programme, a strategic
partnership between the Met Office and the Natural Environment Research Council.