

1 **~~Realized~~Realised ecological forecast through interactive Ecological Platform for**
2 **Assimilating Data into model (EcoPAD v1.0)**
3

4 Yuanyuan Huang^{1,2}, Mark Stacy³, Jiang Jiang^{1,4}, Nilutpal Sudi⁵, Shuang Ma^{1,6}, Volodymyr
5 Saruta^{1,6}, Chang Gyo Jung^{1,6}, Zheng Shi¹, Jianyang Xia^{7,8}, Paul J.Hanson⁹, Daniel Ricciuto⁹, Yiqi
6 Luo^{1,6,10}
7

8 1, Department of Microbiology and Plant Biology, University of Oklahoma, Norman, Oklahoma

9 2, Laboratoire des Sciences du Climat et de l'Environnement, 91191 Gif-sur-Yvette, France

10 3, University of Oklahoma Information Technology, Norman, Oklahoma, USA

11 4, Key Laboratory of Soil and Water Conservation and Ecological Restoration in Jiangsu Province,
12 Collaborative Innovation Center of Sustainable Forestry in Southern China of Jiangsu Province, Nanjing
13 Forestry University, Nanjing, Jiangsu, China

14 5, Department of Computer Science, University of Oklahoma, Norman, Oklahoma, USA

15 6, Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff, AZ, USA

16 7, Tiantong National Forest Ecosystem Observation and Research Station, School of Ecological and
17 Environmental Sciences, East China Normal University, Shanghai 200062, China.

18 8, Research Center for Global Change and Ecological Forecasting, East China Normal
19 University, Shanghai 200062, China

20 9, Environmental Sciences Division and Climate Change Science Institute, Oak Ridge National
21 Laboratory, Oak Ridge, Tennessee, USA

22 10, Department of Earth System Science, Tsinghua University, Beijing 100084 China
23

24 Correspondence: Yuanyuan Huang (yuanyuanhuang2011@gmail.com) and Yiqi Luo
25 (Yiqi.Luo@nau.edu)

26 **Abstract.** Predicting future changes in ecosystem services is not only highly desirable but also
27 becomes feasible as several forces (e.g., available big data, developed data assimilation (DA)
28 techniques, and advanced cyberinfrastructure) are converging to transform ecological research to
29 quantitative forecasting. To realize ecological forecasting, we have developed an Ecological
30 Platform for Assimilating Data (EcoPAD) into models. EcoPAD (v1.0) is a web-based software
31 system that automates data transfer and ~~processes~~processing from sensor networks to ecological
32 forecasting through data management, model simulation, data assimilation, forecasting and
33 visualization. It facilitates interactive data-model integration from which model is recursively
34 improved through updated data while data is systematically refined under the guidance of model.
35 EcoPAD (v1.0) relies on data from observations, process-oriented models, DA techniques, and
36 the web-based workflow.

37 We applied EcoPAD (v1.0) to the Spruce and Peatland Responses Under Climatic and
38 Environmental change (SPRUCE) experiment at North Minnesota. The EcoPAD-SPRUCE
39 realizes fully automated data transfer, feeds meteorological data to drive model simulations,
40 assimilates both manually measured and automated sensor data into Terrestrial ECOSystem
41 (TECO) model, and recursively forecast responses of various biophysical and biogeochemical
42 processes to five temperature and two CO₂ treatments in near real-time (weekly). ~~The near real-~~
43 ~~time forecasting~~Forecasting with EcoPAD-SPRUCE has revealed that ~~uncertainties or~~
44 mismatches in forecasting carbon pool dynamics are more related to model (e.g., model
45 structure, parameter, and initial value) than forcing variables, opposite to forecasting flux
46 variables. EcoPAD-SPRUCE quantified acclimations of methane production in response to
47 warming treatments through shifted posterior distributions of the CH₄:CO₂ ratio and temperature
48 sensitivity (Q₁₀) of methane production towards lower values. Different case studies indicated

49 that realistic forecasting of carbon dynamics relies on appropriate model structure, correct
50 parameterization and accurate external forcing. Moreover, EcoPAD-SPRUCE stimulated active
51 feedbacks between experimenters and ~~modelers so as~~ modellers to identify model components to
52 be improved and additional measurements to be made. It becomes the ~~first~~ interactive model-
53 experiment (ModEx) system and opens a novel avenue for interactive dialogue between
54 ~~modelers and experimenters~~ modellers and experimenters. Altogether, EcoPAD (v1.0) acts to
55 integrate multiple sources of information and knowledge to best inform ecological forecasting.

56 ~~EcoPAD also has the potential to become an interactive tool for resource management, to~~
57 ~~stimulate citizen science in ecology, and transform environmental education with its easily~~
58 ~~accessible web interface. —~~

59

60 **Key words:**

61 Data assimilation, SPRUCE, carbon, global change, real time, acclimation, forecast

62

63 **1. Introduction**

64 One ambitious goal of ecology as ~~one~~ science discipline is to forecast ~~future~~ states and
65 services of ecological systems. Forecasting ~~futures~~ in ecology is not only desirable for scientific
66 advances in this discipline but also has practical values to guide resource management and
67 decision-making toward a sustainable planet earth. The practical need for ecological forecasting
68 is particularly urgent in this rapidly changing world, which is experiencing unprecedented
69 natural resource depletion, increasing food demand, serious biodiversity crisis, accelerated
70 climate changes, and widespread pollutions in the air, waters, and soils [~~Clark et al., 2001;~~
71 ~~Mouquet et al., 2015~~][~~Clark et al., 2001; Mouquet et al., 2015~~]. As a result, a growing number of
72 studies have been reported in the last several decades on forecasting of, e.g., phenology [~~Diez et~~
73 ~~al., 2012~~][~~Diez et al., 2012~~], carbon dynamics [~~Gao et al., 2011; Luo et al., 2016; Thomas et al.,~~
74 ~~2017~~][~~Gao et al., 2011; Luo et al., 2016; Thomas et al., 2017~~], species dynamics [~~Clark et al.,~~
75 ~~2003; Kearney et al., 2010~~][~~Clark et al., 2003; Kearney et al., 2010~~], pollinator
76 performance [~~Corbet et al., 1995~~][~~Corbet et al., 1995~~], epidemics [~~Ong et al., 2010~~][~~Ong et al.,~~
77 ~~2010~~], fishery [~~Hare et al., 2010~~][~~Hare et al., 2010~~], algal bloom [~~Stumpf et al., 2009~~][~~Stumpf et~~
78 ~~al., 2009~~], crop yield [~~Bastiaanssen and Ali, 2003~~][~~Bastiaanssen and Ali, 2003~~], biodiversity
79 [~~Botkin et al., 2007~~][~~Botkin et al., 2007~~], plant extinction risk [~~Fordham et al., 2012~~][~~Fordham et~~
80 ~~al., 2012~~], and ecosystem service [~~Craft et al., 2009~~][~~Craft et al., 2009~~]. Despite its broad
81 applications, ecological forecasting is still sporadically practiced and lags far behind demand due
82 to the lack of infrastructure that enables timely integration of models with data. This paper
83 introduces the fully interactive infrastructure, the Ecological Platform for Assimilating Data
84 (EcoPAD) into models, to inform near-time ecological forecasting with iterative data-model
85 integration.

86 Ecological forecasting relies on both models and data. However, currently the ecology
87 research community has not yet adequately integrated observations with models to inform best
88 forecast. Forecasts generated from scenario approaches are qualitative and scenarios are often
89 not based on ecological knowledge [~~Coreau et al., 2009; Coreau et al., 2010~~][~~Coreau et al.,~~
90 ~~2009; Coreau et al., 2010~~]. Data-driven forecasts using statistical methods are generally limited
91 for extrapolation and sometimes contaminated by confounding factors [~~Schindler and Hilborn,~~
92 ~~2015~~][~~Schindler and Hilborn, 2015~~]. Recent emergent mechanism-free non-parametric approach,
93 which depends on the statistical pattern extracted from data, is reported to be promising for
94 short-term forecast [~~Sugihara et al., 2012; Perretti et al., 2013; Ward et al., 2014~~][~~Sugihara et~~
95 ~~al., 2012; Perretti et al., 2013; Ward et al., 2014~~], but has limited capability in long-term
96 prediction due to the lack of relevant ecological mechanisms. Process-based models provide the
97 capacity in long-term prediction and the flexibility in capturing short-term dynamics on the
98 basis of mechanistic understanding [~~Coreau et al., 2009; Purves et al., 2013~~][~~Coreau et al.,~~
99 ~~2009; Purves et al., 2013~~]. Wide applications ~~and tests~~ of process-based models are limited by
100 their often complicated numerical structure and sometimes unrealistic parameterization
101 [~~Moorcroft, 2006~~][~~Moorcroft, 2006~~]. The complex and uncertain nature of ecology precludes
102 practice of incorporating as many processes as possible into mechanistic models. Our current
103 incomplete knowledge about ecological systems or unrepresented processes under novel
104 conditions is partly reflected in model parameters which are associated with large
105 ~~uncertainty-uncertainties~~. Good forecasting therefore requires effective communication between
106 process-based models and data to estimate realistic model parameters and capture context-
107 dependent ecological phenomena.

108 Data-model fusion, or data-model integration, is an important step to ~~communicate~~
109 ~~model~~combine models with data. But previous data-model integration ~~activities~~activities have
110 mostly been done in an *ad hoc* manner instead of being interactive. For example, data from a
111 network of eddy covariance flux tower sites across United States and Canada was compared with
112 gross primary productivity (GPP) ~~estimates~~estimated from different models [~~Schaefer et al.,~~
113 ~~2012~~][~~Schaefer et al., 2012~~]. ~~Luo and Reynolds [1999]~~. Luo and Reynolds [1999] used a model
114 to examine ecosystem responses to gradual as in the real world vs. step increases in CO₂
115 concentration as in elevated CO₂ experiments. ~~Parton et al. [2007]~~Parton et al. [2007]
116 parameterized CO₂ impacts in an ecosystem model with data from a CO₂ experiment in
117 Colorado. Such model-experiment interactions encounter a few issues: 1) Models are not always
118 calibrated for individual sites and, therefore, not accurate; 2) It is not very effective because it is
119 usually one-time practice without many iterative processes between experimenters and
120 ~~modelers~~modellers [~~Dietze et al., 2013; Lebauer et al., 2013~~][~~Dietze et al., 2013; Lebauer et al.,~~
121 ~~2013~~]; 3) It is usually ~~one-directionary~~unidirectional as data is normally used to train models
122 while the guidance of model for efficient data collection is limited; and 4) It is not streamlined
123 and could not be disseminated with common practices among the research community [~~Dietze et~~
124 ~~al., 2013; Lebauer et al., 2013; Walker et al., 2014~~][~~Dietze et al., 2013; Lebauer et al., 2013;~~
125 ~~Walker et al., 2014~~].

126 A few research groups have developed data assimilation systems to ~~faciliate~~facilitate
127 data-model integration in a systematic way. For example, data-model integration systems, such
128 as the Data Assimilation Research Testbed - DART [~~Anderson et al., 2009~~], ~~the General~~
129 ~~Ensemble Biogeochemical Modeling System - GEMS~~ [~~Tan et al., 2005~~] and the Carbon Cycle
130 Data Assimilation Systems - CCDAS [~~Scholze et al., 2007; Peylin et al., 2016~~][~~Scholze et al.,~~

131 [2007; Peylin et al., 2016](#)], combine various data streams (e.g., FLUXNET data, satellite data and
132 inventory data) with process-based models through data assimilation algorithms such as the
133 Kalman filter [Anderson et al., 2009] and variational methods [[Peylin et al., 2016](#)][[Peylin et al.,
134 2016](#)]. These data ~~assimilation~~assimilation systems automate model parameterization and
135 provided an avenue to systematically improve models through combining as much data as
136 possible. ~~Model-Data-informed model~~ improvements normally happen after the ending of ~~ana~~
137 field experiment and the interactive data-model ~~intergration~~integration is limited as feedbacks
138 from models to ongoing ~~experimentale~~experimental studies are not adequately ~~realized~~realised. In
139 ~~addition~~addition, wide applications of these data assimilation systems in ecological forecasting
140 are constrained by limited user interactions with its steep learning curve to understand these
141 systems, especially for ~~exmperimenterse~~experimenters who have limited training in
142 ~~modeling~~modelling.

Formatted: English (United Kingdom)

143 ~~Realizing interactive ecological forecasting requires~~The web-based technology facilitates
144 interactions. Web-based modelling, which provides user-friendly interfaces to ~~faciliaterun~~
145 models in the background, is usually supported by the scientific workflow, the sequence of
146 processes through which a piece of work passes from initiation to completion. ~~Web-based~~
147 ~~modeling, which provides user friendly interfaces to run models in the background, is uually~~
148 ~~supported by scientific workflow.~~ For example, TreeWatch.Net has recently been developed to
149 make use of -high precision individual tree monitoring data to parameterize process-based tree
150 models in real-time and to assess instant tree hydraulics and carbon status with online result
151 visualization [[Steppe et al., 2016](#)][[Steppe et al., 2016](#)]. Although the web portal of
152 TreeWatch.Net is currently limited to the purpose of visualization ~~purposes~~, it largely broadens
153 the application of data-model integration and strengthens the interaction ~~of modeling results~~

Formatted: Font color: Auto, English (Canada)

154 ~~with~~between modelling researches and the general public. The Predictive Ecosystem Analyzer
155 (PEcAn) is a scientific workflow that wraps around different ecosystem models and manages the
156 flows of information coming in and out of the model [~~Lebauer et al., 2013~~][~~Lebauer et al.,~~
157 ~~2013~~]. PEcAn enables web-based model ~~simulations~~simulations. Such a workflow has
158 advantages, for ~~example~~example, making ecological ~~modeling~~modelling and analysis
159 convenient, transparent, reproducible and adaptable to new questions [~~Lebauer et al.,~~
160 ~~2013~~][~~Lebauer et al., 2013~~], and encouraging user-model interactions. PEcAn uses the Bayesian
161 meta-analysis to synthesize plant trait data to estimate model parameters and associated
162 ~~uncertainties~~uncertainties, i.e., the prior information for process-based models. Parameter
163 uncertainties are ~~propogated~~propagated to model ~~uncertainties~~uncertainties and displayed as
164 outputs. It is still not fully interactive in the way that states are not updated ~~iteratively~~iteratively,
165 according to observations and the web-based data assimilation and then ~~ecoloical~~ecological
166 forecasting have not yet been fully ~~realized~~realised.

Formatted: English (United Kingdom)

Formatted: English (Canada)

167 The iterative model-data ~~intergration~~integration provides an approach to constantly
168 improve ecological forecasting and is an important step ~~to realize real-ore~~especially for realizing
169 near real-time ecological forecasting. Instead of projecting into future ~~only one time~~ through
170 ~~assimulating available~~assimilating observations, ~~interactive only once, the iterative~~
171 constantly updates forecasting ~~as soon as~~along with ongoing new data ~~stream arrives~~streams
172 or/and ~~model is modified~~improved models. Forecasting is likely to be improved unidirectionally
173 in which either only models are ~~constantly~~ updated through observations, or only data
174 collections/field experimentations are regularly improved according to theoretical/model
175 information; but not both. Ecological forecasting can also be bidirectionally improved so that
176 both models and field ~~experimetations~~experimentations are optimized hand in hand over time.

177 Although the ~~bidirectional~~bidirectional case is rare in ecological forecasting, the unidirectional
178 iterative forecasting has been reported. One excellent example of forecasting through
179 dynamically and repeatedly integrating data with models is from infectious disease studies [~~Ong~~
180 ~~et al., 2010; Niu et al., 2014~~][Ong et al., 2010; Niu et al., 2014]. Dynamics of infectious diseases
181 are ~~traditionally~~traditionally captured by Susceptible-Infected-Removed (SIR) models. In the
182 forecasting of the Singapore H1N1-2009 infections, SIR model parameters and the number of
183 individuals in each state were updated daily, combining data renewed from local clinical reports.
184 The evolving of the epidemic related parameters and states were captured through iteratively
185 assimilating observations to inform forecasting. As a result, the model correctly forecasted the
186 timing of the peak and declining of the infection ahead of time. Iterative forecasting dynamically
187 integrates data with model and makes best use of both data and theoretical understandings of
188 ecological processes.

189 The aim of this paper is to present a fully interactive platform, a web-based Ecological
190 Platform for Assimilating Data into models (EcoPAD, v1.0), to best inform ecological
191 forecasting. The interactive feature of EcoPAD (v1.0) is reflected in the iterative model updating
192 and forecasting through dynamically integrating models with new observations, bidirectional
193 feedbacks between experimenters and ~~modelers~~modellers, and flexible user-model
194 communication through web-based simulation, data assimilation and forecasting. Such an
195 interactive platform provides the infrastructure to effectively integrate available resources, from
196 both models and data, ~~modelers~~modellers and experimenters, scientists and the general public, to
197 improve scientific understanding of ecological processes, to boost ecological forecasting practice
198 and transform ecology towards ~~qualitative~~quantitative forecasting.

199 In the following sections, we first describe the system design, major components and
200 functionality of EcoPAD ~~(v1.0)~~. We then use the Spruce and Peatland Responses Under
201 Climatic and Environmental change (SPRUCE) experiment ~~[Hanson et al., 2017]~~[Hanson et al.,
202 2017] as a testbed to elaborate new opportunities brought by the platform. We finally discuss
203 implications of EcoPAD ~~(v1.0)~~ for better ecological forecasting.
204

205 **2 EcoPAD: system design, components, and functionality**

206 **2.1 General description: web-based data assimilation and forecast**

207 EcoPAD ~~(v1.0, https://ecolab.nau.edu/ecopad_portal/)~~ focuses on linking ecological
208 experiments/data with models and allows easily accessible and reproducible data-model
209 integration with interactive web-based simulation, data assimilation and forecast capabilities.
210 Specially, EcoPAD ~~(v1.0)~~ enables the automated near time ecological forecasting which works
211 hand-in-hand between ~~modelers~~modellers and experimenters and updates periodically in a
212 manner similar to the weather forecasting. The system is designed to streamline web request-
213 response, data management, ~~modeling~~modelling, prediction and visualization to boost the overall
214 throughput of observational data, promote data-model communication, inform ecological
215 forecasting and improve scientific understanding of ecological processes.

216 To realize such data-informed ecological forecasting, the essential components of
217 EcoPAD ~~(v1.0)~~ include experiments/data, process-based models, data assimilation techniques
218 and the scientific workflow (Figures 1-3). The scientific workflow of EcoPAD ~~(v1.0)~~ that wraps
219 around ecological models and data assimilation algorithms acts to move datasets in and out of
220 structured and ~~cataloged~~catalogued data collections (metadata catalog) while leaving the logic of
221 the ecological models and data assimilation algorithms untouched (Figures 1, 3). Once a user

222 makes a request through the web browser or command line utilities, the scientific workflow takes
223 charge of triggering and executing corresponding tasks, be it pulling data from a remote server,
224 running a particular ecological model, automating forecasting or making the result easily
225 understandable to users (Figures 1, 3). With the workflow, the system is agnostic to operation
226 system, environment and programming language and is built to horizontally scale to meet the
227 demands of the model and the end user community.

228

229 **2.2 Components**

230 **2.2.1 Data**

231 Data is an important component of EcoPAD ([v1.0](#)) and EcoPAD ([v1.0](#)) offers systematic data
232 management to digest diverse data streams. The ‘big data’ ecology generates ~~plethora~~ large
233 volume of very different datasets across various scales [~~Hampton et al., 2013; Mouquet et al.,~~
234 ~~2015~~][[Hampton et al., 2013; Mouquet et al., 2015](#)]. These datasets might have high temporal
235 resolutions, such as those from real time ecological sensors, or the display of spatial information
236 from remote sensing sources and data stored in the geographic information system (GIS). These
237 datasets may also include, but are not limited to, inventory data, laboratory measurements,
238 FLUXNET databases or from long ~~term ecological networks;~~ term ecological networks
239 [[Baldocchi et al., 2001; Johnson et al., 2010; Robertson et al., 2012](#)]. Such data contain
240 information related to environmental forcing (e.g., precipitation, temperature and radiative
241 forcing), site characteristics (~~including e.g.,~~ soil texture; ~~and~~ species composition) and
242 biogeochemical information. Datasets in EcoPAD ([v1.0](#)) are derived from other research projects
243 in comma separated value files or other loosely structured data formats. These datasets are first
244 described and stored with appropriate metadata via either manual operation or scheduled

245 automation from sensors. Each project has a separate folder where data are stored. Data are
246 generally separated into two categories. One is used as boundary conditions for modelling and
247 the other category is related to observations that are used for data assimilation. Scheduled sensor
248 data are appended to existing data files with prescribed frequency. Attention is then spent on how
249 the particular dataset varies over space (x, y) and time (t). When the spatiotemporal variability is
250 understood, it is then placed in metadata records that allow for query through its scientific
251 workflow.

252 2.2.2 Ecological models

253 Process-based ecological model is another essential component of EcoPAD (Figure 1). In
254 this paper, the Terrestrial ECOsystem (TECO) model is applied as a general ecological model for
255 demonstration ~~purpose~~purposes since the workflow and data assimilation system of EcoPAD
256 (v1.0) are relatively independent on the specific ecological model. Linkages among the
257 workflow, data assimilation system and ecological model are based on messaging. For example,
258 the data assimilation system generates parameters that are passed to ecological models. The state
259 variables simulated from ecological models are passed back to the data assimilation system.
260 Models may have different formulations. As long as they take in the same parameters and
261 generate the same state variables, they are functionally identical from the “eye” of the data
262 assimilation system.

263 TECO simulates ecosystem carbon, nitrogen, water and energy dynamics [~~Weng and Luo,~~
264 ~~2008; Shi et al., 2016~~][~~Weng and Luo, 2008; Shi et al., 2016~~]. The original TECO model has 4
265 major submodules (canopy, soil water, vegetation dynamics and soil carbon/nitrogen) and is
266 further extended to incorporate methane biogeochemistry and snow dynamics [~~Huang et al.,~~
267 ~~2017; Ma et al., 2017~~][~~Huang et al., 2017; Ma et al., 2017~~]. As in the global land surface model

268 CABLE ~~[Wang and Leuning, 1998; Wang et al., 2010]~~[Wang and Leuning, 1998; Wang et al.,
269 2010], canopy photosynthesis that couples surface energy, water and carbon fluxes is based on -a
270 two-big-leaf model ~~[Wang and Leuning, 1998]~~[Wang and Leuning, 1998]. Leaf photosynthesis
271 and stomatal conductance are based on the common scheme from ~~Farquhar et al.~~
272 ~~[1980]~~Farquhar et al. [1980] and ~~Ball et al. [1987]~~Ball et al. [1987] respectively. Transpiration
273 and associated latent heat losses are controlled by stomatal conductance, soil water content and
274 the rooting profile. Evaporation losses of water are balanced between the soil water supply and
275 the atmospheric demand which is based on the difference between saturation vapor pressure at
276 the temperature of the soil and the actual atmospheric vapor pressure. Soil moisture in different
277 soil layers is regulated by water influxes (e.g., precipitation and percolation) and effluxes (e.g.,
278 transpiration and runoff). Vegetation dynamic tracks processes such as growth, allocation and
279 phenology. Soil carbon/nitrogen module tracks carbon and nitrogen through processes such as
280 litterfall, soil organic matter (SOM) decomposition and mineralization. SOM decomposition
281 ~~modeling~~modelling follows the general form of the Century model ~~[Parton et al., 1988]~~[Parton
282 et al., 1988] as in most ~~earth~~Earth system models ~~in which~~, SOM is divided into pools with
283 different turnover times (the inverse of decomposition rates) which are modified by
284 environmental factors such as the soil temperature and moisture.

285 **2.2.3 Data assimilation**

286 Data assimilation is a cutting-edge statistical approach that integrates data with model in
287 a ~~systematical~~systematic way (Figure 2). Data assimilation is growing in importance as the
288 process-based ecological models, despite largely simplifying the real systems, are in great need
289 to be complex enough to address sophisticate ecological issues ~~that~~. These ecological issues are
290 composed of an enormous number of biotic and abiotic factors interacting with each other. Data

291 assimilation techniques provide a framework to combine models with data to estimate model
292 parameters ~~[Shi et al., 2016]~~[Shi et al., 2016], test alternative ecological hypotheses through
293 different model structures ~~[Liang et al., 2015]~~[Liang et al., 2015], assess information content of
294 datasets ~~[Weng and Luo, 2011]~~[Weng and Luo, 2011], quantify uncertainties ~~[Weng et al., 2011;~~
295 ~~Keenan et al., 2012; Zhou et al., 2012]~~[Weng et al., 2011; Keenan et al., 2012; Zhou et al.,
296 2012], derive emergent ecological relationships [Bloom et al., 2016], identify model errors and
297 improve ecological predictions ~~[Luo et al., 2011b]~~[Luo et al., 2011b]. Under the Bayesian
298 paradigm, data assimilation techniques treat the model structure, initial and parameter values as
299 priors that represent our current understanding of the system. As new information from
300 observations or data becomes available, model parameters and state variables can be updated
301 accordingly. The posterior distributions of estimated parameters or state variables are imprinted
302 with information from both the model and the observation/data as the chosen parameters act to
303 reduce mismatches between observations and model simulations. Future predictions benefit from
304 such constrained posterior distributions through forward ~~modeling~~modelling (Figure A1). As a
305 result, the probability density function of predicted future states through data assimilation
306 normally has a narrower spread than that without data assimilation when everything else is equal
307 ~~[Luo et al., 2011b]~~Luo et al., 2011b; Weng and Luo, 2011; Weng and Luo, 2011; Niu et al., 2014].

Field Code Changed

308 EcoPAD (v1.0) is open to different data assimilation techniques depending on the
309 ecological questions under study since the scientific workflow of EcoPAD ~~is~~(v1.0) is relatively
310 independent on the specific data assimilation algorithm. For demonstration, the Markov chain
311 Monte Carlo (MCMC) ~~[Xu et al., 2006]~~[Xu et al., 2006] is described in this study.

312 MCMC is a class of sampling algorithms to draw samples from a probability distribution
313 obtained through constructed Markov Chain to approximate the equilibrium distribution, ~~which~~

314 ~~makes Bayesian inference, especially these with multi-dimensional integrals, workable.~~ The
315 Bayesian based MCMC method ~~is advantageous for better ecological forecasting as it~~ takes into
316 account various uncertainty sources which are crucial in interpreting and delivering forecasting
317 results ~~[Clark et al., 2004][Clark et al., 2001]~~. In the application of MCMC, the posterior
318 distribution of parameters for given observations is proportional to the prior distribution of
319 parameters and the likelihood function which is linked to the fit/match (or cost function) between
320 model simulations and observations. EcoPAD (v1.0) currently adopts a batch mode, that is, the
321 cost function is treated as a single function to be minimized and different observations are
322 standardized by their corresponding standard deviations ~~[Xu et al., 2006][Xu et al., 2006]~~. For
323 simplicity, we assume uniform distributions in priors, and Gaussian or multivariate Gaussian
324 distributions in observational errors, which can be ~~easily~~operationally expanded to other specific
325 distribution forms depending on the available information. Detailed description is available in ~~Xu~~
326 ~~et al. [2006]~~Xu et al. [2006].

327 **2.2.4 Scientific workflow**

328 EcoPAD (v1.0) relies on its scientific workflow to interface ecological models and data
329 assimilation algorithms, managing diverse data streams, automates iterative ecological
330 forecasting in response to various user requests. Workflow is a relatively new concept in the
331 ecology literature but essential to realize real or near-real time forecasting. Thus, we describe it
332 in ~~details~~detail below. The essential components of a scientific workflow of EcoPAD (v1.0)
333 include the metadata catalog, web application-programming interface (API), the asynchronous
334 task/job queue (Celery) and the container-based virtualization platform (Docker). The workflow
335 system of EcoPAD (v1.0) also provides structured result access and visualization.

336 **2.2.4.1 Metadata catalog and data management**

Formatted: Font: Times New Roman

337 Datasets can be placed and queried in EcoPAD (v1.0) via a common metadata catalog
338 which allows for effective management of diverse data streams. Calls are common for good
339 management of current large and heterogeneous ecological datasets [Ellison, 2010; Michener
340 and Jones, 2012; Vitolo et al., 2015][Ellison, 2010; Michener and Jones, 2012; Vitolo et al.,
341 2015]. Kepler [Ludascher et al., 2006][Ludascher et al., 2006] and the Analytic Web [Osterweil
342 et al., 2010][Osterweil et al., 2010] are two example systems that ~~endeavore~~endeavour to provide
343 efficient data management through storage of metadata including clear documentation of data
344 provenance. Similarly to these systems, EcoPAD (v1.0) takes advantage of modern information
345 technology, especially the metadata catalog, to manage diverse data streams. The EcoPAD (v1.0)
346 metadata schema includes description of the data product, security, access pattern, and
347 timestamp of last metadata update *etc.* We use ~~MongDB~~MongoDB (<https://www.mongodb.com/>
348), a NoSQL database technology, to manage heterogeneous datasets to make the documentation,
349 query and storage fast and convenient. Through ~~MongDB~~MongoDB, measured datasets can be
350 easily fed into ecological models for various purposes such as to initialize the model, calibrate
351 model parameters, evaluate model structure and drive model forecast. For datasets from real time
352 ecological sensors that are constantly updating, EcoPAD (v1.0) is set to automatically fetch new
353 data streams with adjustable frequency depending on research needs.

Formatted: Font: Times New Roman

354 2.2.4.2 Web API, asynchronous task queue and docker

355 The RESTful application-programming interface (API) which can deliver data to a wide
356 variety of applications is the gateway of EcoPAD (v1.0) and enables a wide array of user-
357 interfaces and data-dissemination activities. Once a user makes a request, such as through
358 clicking on relevant buttons from a web browser, the request is passed through the
359 Representational State Transfer (i.e., RESTful) API to trigger specific tasks. The RESTful API

360 bridges the talk between the client (e.g., a web browser or command line terminal) and the server
361 (Figure 3). The API exploits the full functionality and flexibility of the HyperText Transfer
362 Protocol (HTTP), such that data can be retrieved and ingested from the EcoPAD (v1.0) through
363 the use of simple HTTP headers and verbs (e.g., GET, PUT, POST, *etc.*). Hence, a user can
364 incorporate summary data from EcoPAD (v1.0) into a website with a single line of html code.
365 Users will also be able to access data directly through programming environments like R, Python
366 and Matlab. Simplicity, ease of use and interoperability are among the main advantages of this
367 API which enables web-based ~~modeling~~modelling.

368 Celery (<https://github.com/celery/celery>) is an asynchronous task/job queue that ~~run~~
369 ~~at~~runs in the background (Figure 3). The task queue (i.e., Celery) is a mechanism used to
370 distribute work across work units such as threads or machines. Celery communicates through
371 messages, and EcoPAD (v1.0) takes advantage of the RabbitMQ (<https://www.rabbitmq.com/>) to
372 manage messaging. After the user ~~submit~~submits a command, the request or message is passed to
373 Celery via the RESTful API. These messages may trigger different tasks, which include, but not
374 limited to, pull data from a remote server where original measurements are located, access data
375 through metadata catalog, run model simulation with user specified parameters, conduct data
376 assimilation which recursively updates model parameters, forecast future ecosystem status and
377 post-process of model results for visualization. The broker inside Celery receives task messages
378 and handles out tasks to available Celery workers which perform the actual tasks (Figure 3).
379 Celery workers are in charge of receiving messages from the broker, executing tasks and
380 returning task results. The worker can be a local or remote computation resource (e.g., the cloud)
381 that has connectivity to the metadata catalog. Workers can be distributed into different
382 ~~information technology~~ (IT) infrastructures, which makes EcoPAD (v1.0) workflow ~~easily~~

383 expandable. Each worker can perform different tasks depending on tools installed in each
384 worker. And one task can also be distributed into different workers. In such a way, EcoPAD
385 [\(v1.0\)](#) workflow enables parallelization and distributed computation of actual
386 ~~modeling~~[modelling](#) tasks across various IT infrastructures, and is flexible in implementing
387 additional computational resources by connecting additional workers.

388 Another key feature that makes EcoPAD [\(v1.0\)](#) easily portable and scalable among
389 different operation systems is the utilization of the container-based virtualization platform, the
390 docker- [\(https://www.docker.com/\)](https://www.docker.com/). Docker can run many applications which rely on different
391 libraries and environments on a single kernel with its lightweight containerization. Tasks that
392 execute TECO in different ways are wrapped inside different docker containers that can “talk”
393 with each other. Each docker container embeds the ecosystem model into a complete filesystem
394 that contains everything needed to run an ecosystem model: the source code, model input, run
395 time, system tools and libraries. Docker containers are both hardware-agnostic and platform-
396 agnostic, and they are not confined to a particular language, framework or packaging system.
397 Docker containers can be run from a laptop, workstation, virtual machine, or any cloud compute
398 instance. This is done to support the widely varied number of ecological models running in
399 various languages (e.g., Matlab, Python, Fortran, C and C++) and environments. In addition to
400 wrap the ecosystem model into a docker container, software applied in the workflow, such as the
401 Celery, Rabbitmq and MongoDB, are all lightweight and portable encapsulations through docker
402 containers. Therefore, the entire EcoPAD [\(v1.0\)](#) is readily portable and applicable in different
403 environments.

404 **2.2.4.3 Structured result access and visualization**

405 EcoPAD (v1.0) enables structured result storage, access and visualization to track and
406 ~~analyze~~analyse data-model fusion practice. Upon the completion of the model task ~~completion~~,
407 the model wrapper code calls a post processing ~~callback~~call-back function. This ~~callback~~call-
408 ~~back~~ function allows for model specific data requirements to be added to the model result
409 repository. Each task is associated with a unique task ID and model results are stored within the
410 local repository that can be queried by the unique task ID. The ~~easy~~store and query of model
411 results are ~~realized~~realised via the MongoDB and RESTful API (Figure 3). Researchers are
412 authorized to review and download model results and parameters submitted for each model run
413 through a web accessible URL (link). EcoPAD (v1.0) webpage also displays a list of historical
414 tasks (with URL) performed by each user. All current and historical model inputs and outputs are
415 available to download, including the aggregated results produced for the graphical web
416 applications. In addition, EcoPAD (v1.0) also provides a task report that contains all-inclusive
417 recap of parameters submitted, task status, and model outputs with links to all data and graphical
418 results for each task. Such structured result storage and access make sharing, tracking and
419 referring to ~~modeling~~modelling studies instant and clear.

420 **2.3 Scientific functionality**

421 Scientific functionality of EcoPAD (v1.0) includes web-based model simulation,
422 estimating model parameters or state variables, quantifying uncertainty of estimated parameters
423 and projected states of ecosystems, evaluating model structures, assessing sampling strategies,
424 ~~and~~ conducting ecological forecasting. ~~These~~These functions can be organized to answer various
425 scientific questions. In addition to the general description in this section, the scientific
426 functionality of EcoPAD (v1.0) is also illustrated through a few case studies in the following
427 sections.

428 EcoPAD ([v1.0](#)) is designed to perform web-based model simulation, which greatly
429 reduces the workload of traditional model simulation through manual code compilation and
430 execution. This functionality opens various new opportunities for ~~modelers~~modellers,
431 experimenters and the general public. Model simulation and result analysis are automatically
432 triggered after a ~~simple~~-click on the web-embedded button (Appendices Figures A2, A3 A6).
433 Users are freed from repeatedly compiling code, running code and writing programs to
434 ~~analyze~~analyse and display model results. Such ease of use has great potential to popularize
435 complex ~~modeling~~modelling studies that are difficult or inaccessible for experimenters and the
436 general public. As illustrated through the outreach activities from the TreeWatch.Net [*Steppe et*
437 *al., 2016*][*Steppe et al., 2016*], the potential functionality of such web-based model simulation
438 goes beyond its scientific value as its societal and educational impacts are critical in solving
439 ecological issues. The web-based model simulation also frees users from model running
440 environment, platform and software. Users can conduct model simulation and do analysis as long
441 as they have internet access. For example, ecologists can conduct model simulation and diagnose
442 the underlying reasons for a sudden increase in methane fluxes while they are making
443 measurements in the field. ~~Youngsters~~Non-ecologists, such as youngsters, can study ecological
444 dynamics through their phones or tablets while they are waiting for the bus. Resource managers
445 can make timely assessment of different resource utilization strategies on spot of a meeting.

446 EcoPAD ([v1.0](#)) is backed up by data assimilation techniques, which facilitate inference of
447 model parameters and states based on observations. Ecology have witnessed a growing number
448 of studies focusing on parameter estimation using inverse ~~modeling~~modelling or data
449 assimilation as large volumes of ecological measurements become available. To satisfy the
450 growing need of model parameterization through observations, EcoPAD ([v1.0](#)) streamlines

451 parameter estimations and updates. Researchers can ~~easily~~ review and download files that record
452 parameter values from EcoPAD [\(v1.0\)](#) result repository. Since these parameters may have
453 different ~~scientific values~~ [biological, physical or chemical meanings](#), the functionality of EcoPAD
454 [\(v1.0\)](#) related to parameter estimations can potentially embrace diverse subareas in ecology. For
455 example, soil scientists can study the acclimation of soil respiration to manipulative warming
456 through shifts in the distribution of the decomposition rate parameter from EcoPAD ~~(v1.0)~~. The
457 threshold parameter beyond which further harvesting of fish might cause a crash of fish stocks
458 can be ~~easily~~ extracted through fish stock assessment models and observations if mounted to
459 EcoPAD ~~(v1.0)~~.

460 EcoPAD [\(v1.0\)](#) promotes uncertainty analysis, model structure evaluation and error
461 identification. One of the advantages of the Bayesian statistics is its capacity in uncertainty
462 analysis compared to other optimization techniques ~~[Xu et al., 2006; Wang et al., 2009; Zhou et~~
463 ~~al., 2012]~~ [\[Xu et al., 2006; Wang et al., 2009; Zhou et al., 2012\]](#). Bayesian data assimilation (e.g.,
464 MCMC) takes into account observation uncertainties (errors), generates distributions of model
465 parameters and enables tracking of prediction uncertainties from different sources: ~~[Ellison,~~
466 ~~2004; Bloom et al., 2016; Jiang et al., 2018]~~. Uncertainty analysis through data assimilation
467 applied to areas such as ecosystem phenology, fish life cycle and species migration ~~[Clark et al.,~~
468 ~~2003; Cook et al., 2005; Crozier et al., 2008; Luo et al., 2011b]~~ [\[Clark et al., 2003; Cook et al.,](#)
469 [2005; Crozier et al., 2008; Luo et al., 2011b\]](#), can potentially take advantage of EcoPAD [\(v1.0\)](#)
470 platform to provide critical information for well informed decisions in face of pressing global
471 change challenges. In addition, the archive capacity of EcoPAD [\(v1.0\)](#) facilitates [future](#) inter-
472 comparisons among different models or different versions of the same model to evaluate model
473 structures and to disentangle structure uncertainties and errors.

474 The realization of both the near-time and long-term ecological forecast is one of the key
475 innovations of EcoPAD (v1.0). Forecasting capability of EcoPAD (v1.0) is supported by process
476 -based ecological models, multiple observational or experimental data, inverse parameter
477 estimation and uncertainty quantification through data assimilation, and forward simulation
478 under future external conditions. The systematically constrained forecast from EcoPAD (v1.0) is
479 accompanied by uncertainty/confidence estimates to quantify the amount of information that can
480 actually be utilized from a study. The automated near time forecast, which is constantly adjusted
481 once new observational data streams are available, provides experimenters advanced and timely
482 information to assess and adjust experimental plans. For example, with forecasted and displayed
483 biophysical and biochemical variables, experimenters could know in advance what the most
484 likely biophysical conditions are. Knowing if the water table may suddenly go aboveground in
485 response to a high rainfall forecast in the coming week, could allow researcher to emphasize
486 measurements associated with methane flux. In such a way, experimenters can not only rely on
487 historical ecosystem dynamics, but also refer to future predictions. Experimenters will benefit
488 especially from variables that are difficult to track in field due to situations such as harsh
489 environment, shortage in man power or on instrument limitation.

490 Equally important, EcoPAD (v1.0) creates new avenues to answer classic and novel
491 ecological questions, for example, the frequently reported acclimation phenomena in ecology.
492 While growing evidence points to altered ecological functions as organisms adjust to the rapidly
493 changing world [Medlyn et al., 1999; Luo et al., 2001; Wallenstein and Hall, 2012][Medlyn et
494 al., 1999; Luo et al., 2001; Wallenstein and Hall, 2012], traditional ecological models treat
495 ecological processes less dynamical, as the governing biological parameters or mechanisms fails
496 to explain such biological shifts. EcoPAD (v1.0) facilitates the shift of research paradigm from a

497 fixed process representation to a more dynamic description of ecological mechanisms with
498 constantly updated and archived parameters constrained by observations under different
499 conditions. Specifically to acclimation, EcoPAD (v1.0) promotes quantitative evaluations
500 while previous studies remain mostly qualitative [Wallenstein and Hall, 2012; Shi et al.,
501 2015][Wallenstein and Hall, 2012; Shi et al., 2015]. We will further illustrate how EcoPAD
502 (v1.0) can be used to address different ecological questions in the case studies of the SPRUCE
503 project.

504

505 **3 EcoPAD performance at testbed - SPRUCE**

506 **3.1 SPRUCE project overview**

507 EcoPAD (v1.0) is being applied to the Spruce and Peatland Responses Under Climatic
508 and Environmental change (SPRUCE) experiment located at the USDA Forest Service Marcell
509 Experimental Forest (MEF, 47°30.476' N, 93°27.162' W) in northern Minnesota [Kolka et al.,
510 2011][Kolka et al., 2011]. SPRUCE is an ongoing project focuses on long-term responses of
511 northern peatland to climate warming and increased atmospheric CO₂ concentration [Hanson et
512 al., 2017][Hanson et al., 2017]. At SPRUCE, ecologists measure various aspects of responses of
513 organisms (from microbes to trees) and ecological functions (carbon, nutrient and water cycles)
514 to a warming climate. One of the key features of the SPRUCE experiments is the manipulative
515 deep soil/peat heating (0-3 m) and whole ecosystem warming treatments (peat + air warmings)
516 which include tall trees (> 4 m) [Hanson et al., 2017][Hanson et al., 2017]. Together with
517 elevated atmospheric CO₂ treatments, SPRUCE provides a platform for exploring mechanisms
518 controlling the vulnerability of organisms, biogeochemical processes and ecosystems in response
519 to future novel climatic conditions. The SPRUCE peatland is especially sensitive to future

520 climate change and also plays an important role in feeding back to future climate change through
521 greenhouse gas emissions as it stores a large amount of soil organic carbon. Vegetation in the
522 SPRUCE site is dominated by *Picea mariana* (black spruce) and *Sphagnum spp* (peat moss). The
523 studied peatland also has an understory which include ericaceous and woody shrubs. There are
524 also a limited number of herbaceous species. The whole ecosystem warming treatments include a
525 large range of both aboveground and belowground temperature manipulations (ambient, control
526 plots of +0 °C, +2.25 °C, +4.5 °C, +6.75 °C and +9 °C) in large 115 m² open-topped enclosures
527 with elevated CO₂ manipulations (+0 or +500 ppm). The difference between ambient and +0
528 treatment plots is the open-topped and controlled-environment enclosure.

529 The SPRUCE project generates a large variety of observational datasets that reflect
530 ecosystem dynamics from different scales and are available from the project webpage
531 (<https://mnspruce.ornl.gov/>) and FTP site (<ftp://sprucedata.ornl.gov/>). These datasets come from
532 multiple sources: half hourly automated sensor records, species surveys, laboratory
533 measurements, laser scanning images *etc.* Involvements of both ~~modeling~~modelling and
534 experimental studies in the SPRUCE project create the opportunity for data-model
535 communication. Datasets are pulled from SPRUCE archives and stored in the EcoPAD ([v1.0](#))
536 metadata catalog for running the TECO model, conducting data-model fusion or forecasting. The
537 TECO model has been applied to simulate and forecast carbon dynamics with productions of
538 CO₂ and CH₄ from different carbon pools, soil temperature response, snow depth and freeze-
539 thaw cycles at the SRPUCE site [~~Huang et al., 2017; Ma et al., 2017; Jiang et al., 2018~~][Huang
540 et al., 2017; Ma et al., 2017; Jiang et al., 2018].

541

542 3.2 EcoPAD-SPRUCE web portal

543 We assimilate multiple streams of data from the SPRUCE experiment to the TECO
544 model using the MCMC algorithm, and forecast ecosystem dynamics in both near time and for
545 the next 10 years. Our forecasting system for SPRUCE is available at
546 https://ecolab.nau.edu/ecopad_portal/. From the web portal, users can check our current near-
547 and long-term forecasting results, conduct model simulation, data assimilation and forecasting
548 runs, and ~~analyze~~analyse/visualize model results. Detailed information about the interactive web
549 portal is provided in the Appendices.

550 3.3 Near time ecosystem forecasting and feedback to experimenters

551 As part of the forecasting functionality, EcoPAD-SPRUCE automates the near time
552 (weekly) forecasting with continuously updated observations from SPRUCE experiments (Figure
553 ~~54~~). We set up the system to automatically pull new data streams every Sunday from the
554 SPRUCE FTP site that holds observational data and update the forecasting results based on new
555 data streams. Updated forecasting results for the next week are customized for the SPRUCE
556 experiments with different manipulative treatments and displayed in the EcoPAD-SPRUCE
557 portal. At the same time, these results are sent back to SPRUCE communities and displayed
558 together with near-term observations for experimenter's reference.

559 3.4 New approaches to ecological studies towards better forecasting

560 3.4.1 Case 1: Interactive communications among ~~modelers~~modellers and experimenters

561 EcoPAD-SPRUCE provides a platform to stimulate interactive communications between
562 ~~modelers~~modellers and experimenters. Models require experimental data to constrain initial
563 conditions and parameters, and to verify model performance. A reasonable model is built upon
564 correct interpretation of information served by experimenters. Model simulations on the other
565 hand can expand ~~hypotheses~~hypothesis testing, and provide thorough or advanced information to

566 improve field experiments. Through recursively exchanging information between
567 ~~modelers~~modellers and experimenters, both models and field experiments can be improved. As
568 illustrated in Figure 54, through extensive communication between ~~modelers~~modellers and
569 experimenters, ~~modelers~~modellers generate model predictions. Model predictions provide
570 experimenters advanced information, help experimenters think, question and understand their
571 experiments. Questions raised by experimenters stimulate further discussion and communication.
572 Through communication, models or/and measurements are adjusted. With new measurements
573 or/and strengthened models, a second round of prediction is highly likely to be improved. As the
574 loop of prediction-question-discussion-adjustment-prediction goes on, forecasting is informed
575 with best understandings from both data and model.

576 We illustrate how the prediction-question-discussion-adjustment-prediction cycle and
577 stimulation of ~~modeler~~modeller-experimenter communication improves ecological predictions
578 through one episode during the study of the relative contribution of different pathways to
579 methane emissions. An initial methane model was built upon information (e.g., site
580 characteristics and environmental conditions) provided by SPRUCE field scientists, taking into
581 account important processes in methane dynamics, such as production, oxidation and emissions
582 through three pathways (i.e., diffusion, ebullition and plant-mediated transportation). The model
583 was used to predict relative contributions of different pathways to overall methane emissions
584 under different warming treatments after being constrained by measured surface methane fluxes.
585 Initial forecasting results which indicated a strong contribution from ebullition under high
586 warming treatments were sent back to the SPRUCE group. Experimenters doubted about such a
587 high contribution from the ebullition pathway and a discussion was stimulated. It is difficult to
588 accurately distinguish the three pathways from field measurements. Field experimenters

589 provided potential avenues to extract measurement information related to these pathways, while
590 ~~modelers~~ modellers examined model structure and parameters that may not be well constrained
591 by available field information. Detailed discussion is provided in Table 1. After extensive
592 discussion, several adjustments were adopted as a first step to move forward. For example, the
593 three-porosity model that was used to simulate the diffusion process was replaced by the
594 Millington-Quirk model to more realistically represent methane diffusions in peat soil; the
595 measured static chamber methane fluxes were also questioned and scrutinized more carefully to
596 clarify that they did not capture the episodic ebullition events. Measurements such as these
597 related to pore water gas data may provide additional inference related to ebullition. The updated
598 forecasting is more reasonable than the initial results although more studies are in need to
599 ultimately quantify methane fluxes from different pathways.

600 **3.4.2 Case 2: Acclimation of ecosystem carbon cycling to experimental manipulations**

601 As a first step, CH₄ static chamber flux measurements were assimilated into TECO to
602 assess potential acclimation phenomena during methane production under 5 warming treatments
603 (+0, +2.25, +4.5, +6.75, +9 °C). Initial results indicated a reduction in both the CH₄:CO₂ ratio and
604 the temperature sensitivity of methane production based on their posterior distributions (Figure
605 ~~65~~). The mean CH₄:CO₂ ratio decreased from 0.675 (~~control~~ (+0 °C treatment)) to 0.505 (+9 °C
606 ~~treatment~~), while the temperature sensitivity (Q₁₀) for CH₄ production decreased from 3.33
607 (~~control~~ (+0 °C)) to 1.22 (+9 °C treatment). Such shifts quantify potential acclimation of methane
608 production to warming and future climate warming is likely to have a smaller impact on emission
609 than most of current predictions that do not take into account of acclimation.

610 Despite these results are preliminary as more relevant datasets are under collection with
611 current ongoing warming manipulation and measurements, assimilating observations through

612 EcoPAD ([v1.0](#)) provides a quantitative approach to timely assess acclimation through time. [Melillo](#)
613 [et al. \[2017\]](#), [Melillo et al. \[2017\]](#) revealed that the thermal acclimation of the soil respiration in the
614 Harvard Forest is likely to be phase (time) dependent during their 26-year soil warming experiment.
615 EcoPAD ([v1.0](#)) provides the possibility in tracing the temporal path of acclimation with its
616 streamlined structure and archive capacity. [Shi et al. \[2015\]](#), [Shi et al. \[2015\]](#) assimilated carbon
617 related measurements in a tallgrass prairie into the TECO model to study acclimation after 9-years
618 warming treatments. They revealed a reduction in the allocation of GPP to shoot, the turnover rates
619 of the shoot and root carbon pools, and an increase in litter and fast carbon turnovers in response
620 to warming treatments. Similarly, as time goes on, the SPRUCE experiment will generate more
621 carbon cycling related datasets under different warming and CO₂ treatments, which can be
622 mounted to EcoPAD ([v1.0](#)) to systematically quantify acclimations in carbon cycling [through time](#)
623 [in the future](#).

624 3.4.3 Case 3: Partitioning of uncertainty sources

625 Uncertainties in ecological studies can come from observations (include forcing that
626 drives the model), different model structures to represent the real world and the specified model
627 parameters [\[Luo et al., 2016\]](#), [\[Luo et al., 2016\]](#). Previous studies tended to focus on one aspect of
628 the uncertainty sources instead of disentangling the contribution from different sources. For
629 example, the model intercomparison projects (MIPs), such as TRENDY, focus on uncertainty
630 caused by different model structures with prescribed external forcing [\[Sitch et al., 2008\]](#), [\[Sitch et](#)
631 [al., 2008\]](#), [Keenan et al. \[2012\]](#), [Keenan et al. \[2012\]](#) used data assimilation to constrain
632 parameter uncertainties in projecting Harvard forest carbon dynamics. [Ahlstrom et al. \[2012\]](#)
633 forced one particular vegetation model by 18 sets of forcings from climate models of the

634 Coupled Model Intercomparison Project Phase 5 (CMIP5), while the parameter or model
635 structure uncertainty is not taken into account.

636 EcoPAD (v1.0) is designed to provide a thorough picture of uncertainties from multiple
637 sources especially in carbon cycling studies. Through focusing on multiple instead of one source
638 of uncertainty, ecologists can allocate resources to areas that cause relative high uncertainty.

639 Attribution of uncertainties in EcoPAD ~~relies~~(v1.0) will rely on an ensemble of ecosystem
640 models, the data assimilation system and climate forcing with quantified uncertainty. ~~For~~
641 ~~example, Jiang et al. [2018]~~Jiang et al. [2018] focused specifically on the relative contribution
642 of parameter uncertainty vs. climate forcing uncertainty in forecasting carbon dynamics at the
643 SPRUCE site. Through assimilating the pre-treatment measurements (2011-2014) from the
644 SPRUCE experiment, Jiang et al. [2018]Jiang et al. [2018] estimated uncertainties of key
645 parameters that regulate the peatland carbon dynamics. Combined with the stochastically
646 generated climate forcing (e.g., precipitation and temperature), Jiang et al. [2018]Jiang et al.
647 [2018] found external forcing resulted in higher uncertainty than parameters in forecasting
648 carbon fluxes, but caused lower uncertainty than parameters in forecasting carbon pools.

649 Therefore, more efforts are required to improve forcing measurements for studies that focus on
650 carbon fluxes (e.g., GPP), while reductions in parameter uncertainties are more important for
651 studies in carbon pool dynamics. ~~Such kind of uncertainty assessment benefits from EcoPAD~~
652 ~~with its systematically archived model simulation, data assimilation and forecasting. Despite~~
653 Jiang et al. [2018] does not quantify model structure uncertainty, the project of incorporating
654 multiple models inside EcoPAD (v1.0) is in progress, and future uncertainty assessment will
655 benefit from EcoPAD (v1.0) with its systematically archived model simulation, data assimilation
656 and forecasting.

657 **3.4.4 Case 4: Improving biophysical estimation for better ecological prediction**

658 Carbon cycling studies can also benefit from EcoPAD (v1.0) through improvements in
659 ~~external forcing~~biophysical estimation. Soil environmental condition is an important regulator of
660 belowground biological activities and also feeds back to aboveground vegetation growth.

661 Biophysical variables such as soil temperature, soil moisture, ice content and snow depth, are
662 key predictors of ecosystem dynamics. After constraining the biophysical module by detailed
663 monitoring data from the SPRUCE experiment through the data assimilation component of
664 EcoPAD, ~~Huang et al. [2017]~~(v1.0), Huang et al. [2017] forecasted the soil thermal dynamics
665 under future conditions and studied the responses of soil temperature to hypothetical air
666 warming. This study emphasized the importance of accurate climate forcing in providing robust
667 thermal forecast. ~~In addition, Huang et al. [2017]~~In addition, Huang et al. [2017] revealed non-
668 uniform responses of soil temperature to air warming. Soil temperature responded stronger to air
669 warming during summer compared to winter. And soil temperature increased more in shallow
670 soil layers compared to deep soils in summer in response to air warming. Therefore,
671 extrapolating of manipulative experiments based on air warming alone may not reflect the real
672 temperature sensitivity of SOM if soil temperature is not monitored. As robust quantification of
673 environmental conditions is known to be a first step towards better understanding of ecological
674 process, improvement in soil thermal predictions through EcoPAD (v1.0) data assimilation
675 system is helpful in telling apart biogeochemical responses from environmental uncertainties and
676 also in providing field ecologists beforehand key environmental conditions.

677 **3.4.5 Case 5: How do updated model and data contribute to reliable forecasting?**

678 Through constantly adjusted model and external forcing according to observations and
679 weekly archived model parameter, model structure, external forcing and forecasting results, the

680 contribution of model and data updates can therefore be tracked through comparing forecasted
681 vs. ~~realized~~realised simulations. For example, Figure 76 illustrates how ~~realized~~updated external
682 forcing (compared to stochastically generated forcing) and shifts in ecosystem state variables
683 shape ecological predictions. Similarly as in other EcoPAD-SPURCE case studies, TECO is
684 trained through data assimilation with observations from 2011-2014 and is used to forecast GPP
685 and total soil organic carbon content at the beginning of 2015. For demonstrating purpose, Figure
686 76 only shows 3 series of forecasting results instead of updates from every week. Series 1 (S1)
687 records forecasted GPP and soil carbon with stochastically generated weather forcing from
688 January 2015-December 2024 (Figure 7a6a,b cyan). Series 2 (S2) records simulated GPP and
689 soil carbon with observed climate forcing from January 2015 to July 2016 and forecasted GPP
690 and soil carbon with stochastically generated forcing from August 2016 - December 2024
691 (Figure 7a6a,b red). Similarly, the stochastically generated forcing in Series 3 (S3) starts from
692 January 2017 (Figure 7a6a,b blue). For each series, predictions were conducted with randomly
693 sampled parameters from the posterior distributions and stochastically generated forcing. We
694 displayed 100 mean values (across an ensemble of forecasts with different parameters)
695 corresponding to 100 forecasts with stochastically generated forcing.

696 GPP is highly sensitive to climate forcing. The differences between the ~~realized~~updated
697 (S2, 3) and initial forecasts (S1) reach almost $800 \text{ gC m}^{-2} \text{ year}^{-1}$ (Figure 7e6c). The discrepancy
698 is strongly dampened in the following 1-2 years. The impact of ~~realized~~updated forecasts is close
699 to 0 after approximately 5 years. However, soil carbon pool shows a different pattern. Soil
700 carbon pool is increased by less than 150 gC m^{-2} , which is relative small compared to the carbon
701 pool size of *ca.* 62000 gC m^{-2} . The impact of ~~realized~~updated forecasts grows with time and
702 reaches the highest at the end of the simulation year 2024. GPP is sensitive to the immediate

703 change in climate forcing while the updated ecosystem status (or initial value) has minimum
704 impact in the long-term forecast of GPP. The impact of updated climate forcing is relatively
705 small for soil carbon forecasts during our study period. Soil carbon is less sensitive to the
706 immediate change of climate compared to GPP. However, the alteration of system status affects
707 soil carbon forecast especially in a longer time scale.

708 Since we are archiving ~~realized~~updated forecasts every week, we can track the relative
709 contribution of ecosystem status, forcing uncertainty and parameter distributions to the overall
710 forecasting patterns of different ecological variables and how these patterns evolve in time. In
711 addition, as growing observations of ecological variables (e.g., carbon fluxes and pool sizes)
712 become available, it is feasible to diagnose key factors that promote robust ecological forecasting
713 through comparing the archived forecasts vs. observation and analysing archives of model
714 parameters, initial values and climate forcing *etc.*

715

716 **4 Discussion**

717 **4.1 The necessity of interactive infrastructure to realize ecological forecasting**

718 Substantial increases in data availability from observational and experimental networks,
719 surges in computational capability, advancements in ecological models and sophisticated
720 statistical methodologies and pressing societal need for best management of natural resources
721 have shifted ecology to emphasis more on quantitative forecasts. However, quantitative
722 ecological forecast is still young and our knowledge about ecological forecasting is relatively
723 sparse, inconsistent and disconnected [~~Luo et al., 2011b; Petchey et al., 2015~~][Luo et al., 2011b;
724 Petchey et al., 2015]. Therefore, both optimistic and pessimistic viewpoints exist on the
725 predictability of ecology [~~Clark et al., 2001; Beckage et al., 2011; Purves et al., 2013; Petchey et~~

726 ~~al., 2015; Schindler and Hilborn, 2015]~~[Clark et al., 2001; Beckage et al., 2011; Purves et al.,
727 2013; Petchey et al., 2015; Schindler and Hilborn, 2015]. Ecological forecasting is complex and
728 advantages in one single direction, for example, observations alone or statistical methodology
729 alone, is less likely to lead to successful forecasting compared to approaches that effectively
730 integrate improvements from multiple sectors. Unfortunately, ~~realized~~realised ecological
731 forecasting that integrates available resources is relative rare due to lack of relevant
732 infrastructures.

733 EcoPAD (v1.0) provides such effective infrastructure with its interactive platform that
734 rigorously integrates merits from models, observations, statistical advance, information
735 technology and human resources from ~~experimenter, modeler as well as the general~~
736 ~~public~~experimenters and modellers to best inform ecological forecasting, boost forecasting
737 practice and delivery of forecasting results. Interactions enable exchanging and extending of
738 information so as to benefit from collective knowledge. For example, manipulative studies will
739 have a much broader impact if the implications of their results can be extended from the
740 regression between environmental variable and ecosystem response, such as be integrated into an
741 ecosystem model through model-data communication. Such an approach will allow gaining
742 information about the processes responsible for ecosystem's response, constraining models, and
743 making more reliable predictions. Going beyond common practice of model-data assimilation
744 from which model updating lags far behind observations, EcoPAD (v1.0) enables iterative model
745 updating and forecasting through dynamically integrating models with new observations in near
746 real-time. This near real-time interactive capacity relies on its scientific workflow that automates
747 data management, model simulation, data simulation and result visualization. The ~~open, timely,~~
748 ~~convenient, transparent, flexible, reproducible and traceable characteristics of this platform, also~~

749 ~~thanks to its scientific workflow, encouraged~~system design encourages thorough interactions
750 between experimenters and ~~modelers~~modellers. Forecasting results from SPRUCE were timely
751 shared among research groups with different background through the web interface. Expertise
752 from different research groups was integrated to improve a second round of forecasting. Again,
753 thanks to the workflow, new information or adjustment is relatively easy to
754 ~~incorporate~~incorporated into ~~future~~-forecasting efficiently, making the forecasting system fully
755 interactive and dynamical.

756 We also benefit from the interactive EcoPAD (v1.0) platform to broaden user-model
757 interactions and to broadcast forecasting results. Learning about the ecosystem models and data-
758 model fusion techniques may lag one's productivity and even discourage learning the
759 ~~modeling~~modelling techniques because of their complexity and long learning curve. Because
760 EcoPAD (v1.0) can be accessed from a web browser and does not require any coding from the
761 user's side, the time lag between learning the model structure and obtaining model-based results
762 for one's study is minimal, which opens the door for non-~~modeler~~modeller groups to "talk" with
763 models. The online storage of one's results lowers the risk of data loss. The results of each model
764 run can be easily tracked and shared with its unique ID and web address. In addition, the web-
765 based workflow also saves time for experts with automated model running, data assimilation,
766 forecasting, structured result access and instantaneous graphic outputs, bringing the possibility
767 for thorough exploration of more essence part of the system. The simplicity in use of EcoPAD
768 (v1.0) at the same time may limit their access to the code and lowers the flexibility. Flexibility
769 for users with higher demands, for example, those who wanted to test alternative data
770 assimilation methods, use a different carbon cycle model, change the number of calibrated
771 parameters, include the observations for other variables, is provided through the GitHub

772 repository (<https://github.com/ou-ecolab>). This GitHub repository contains code and instruction
773 for installing, configuring and controlling the whole system, users can easily adapt the workflow
774 to wrap their own model based on his or her needs.

775 ~~In addition to benefit from its workflow, the advantage of EcoPAD is also reflected in its~~
776 ~~data assimilation capacity especially for land carbon studies. One focus of EcoPAD is to~~
777 ~~constrain parameters of terrestrial carbon models to predict long-term carbon dynamics (e.g., 100~~
778 ~~years) which are determined more by parameters than initial values of state variables [Weng and~~
779 ~~Luo, 2011]. EcoPAD incorporates the Bayesian framework, especially the MCMC method, to~~
780 ~~constrain parameters. In comparison, DART uses the Ensemble Kalman Filter to adjust model~~
781 ~~state variables, instead of parameters, to match observations over time. In the past, complex~~
782 ~~models could not assimilate pool-related data to constrain their parameter estimation due to~~
783 ~~insurmountable computational demand in large-scale studies. For example, CCDAS normally~~
784 ~~only assimilates flux-based data [Peylin et al., 2016]. EcoPAD is flexible in assimilating both~~
785 ~~pool and flux-based data into complex models so that both fluxes and turnover rates of pools~~
786 ~~can be constrained with its matrix representation [Hararuk et al., 2014; Luo, 2017] and its~~
787 ~~capability to wrap different models.~~

788 **4.2 Implications for better ecological forecasting**

789 Specifically to reliable forecasting of carbon dynamics, our initial exploration from
790 EcoPAD-SPRUCE indicates that realistic model structure, correct parameterization and accurate
791 external environmental conditions are essential. Model structure captures important known
792 mechanisms that regulate ecosystem carbon dynamics. Adjustment in model structure is critical
793 in our improvement in methane forecasting. Model parameters may vary between observation
794 sites, change with time or environmental conditions [Medlyn et al., 1999; Luo et al.,

795 ~~2004~~[[Medlyn et al., 1999](#); [Luo et al., 2001](#)]. A static or wrong parameterization misses
796 important mechanisms (e.g., acclimation and adaptation) that regulate future carbon dynamics.
797 Not well constrained parameters, for example, caused by lack of information from observational
798 data, contribute to high forecasting uncertainty and low reliability of forecasting results. Correct
799 parameterization is especially important for long-term carbon pool predictions as parameter
800 uncertainty resulted in high forecasting uncertainty in our case study [[Jiang et al., 2018](#)][[Jiang et](#)
801 ~~al., 2018~~]. ~~Although the picture about how neglecting of parameter shift affects carbon~~
802 ~~predictions has not yet been fully revealed from EcoPAD-SPRUCE as field measurements are~~
803 ~~still ongoing, our initial exploration indicates non-negligible acclimation of ecosystem methane~~
804 ~~production in response to warming. Parameter values derived under the ambient condition was~~
805 ~~not applicable to the warming treatment in our methane case due to acclimation.~~ External
806 environmental condition is another important factor in carbon predictions. External
807 environmental condition includes both the external climatic forcing that is used to drive
808 ecosystem models and also the environmental condition that is simulated by ecosystem models.
809 As we showed that air warming may not proportionally transfer to soil warming, realistic soil
810 environmental information needs to be appropriately represented to predict soil carbon dynamics
811 [[Huang et al., 2017](#)][[Huang et al., 2017](#)]. The impact of external forcing is especially obvious in
812 short term carbon flux predictions. Forcing uncertainty resulted in higher forecasting uncertainty
813 in carbon flux compared to that from parameter uncertainty [[Jiang et al., 2018](#)][[Jiang et al.,](#)
814 [2018](#)]. Mismatches in forecasted vs. ~~realized~~realised forcing greatly increased simulated GPP
815 and the discrepancy diminished in the long run. Reliable external environmental condition, to
816 some extent, reduces the complexity in diagnosing ~~modeled~~modelled carbon dynamics.

817 Pool-based vs. flux-based predictions are regulated differently by external forcing and
818 initial states, which indicates that differentiated efforts are required to improve short vs. long-
819 term predictions. External forcing, which has not been well emphasized in previous carbon
820 studies, has strong impact on short term forecasting. The large response of GPP to forecasted vs.
821 ~~realized~~realised forcing as well the stronger forcing-caused uncertainty in GPP predictions
822 indicate correct forcing information is a key step in short term flux predictions. In this study, we
823 stochastically generated the climate forcing based on local climatic conditions (1961-2014),
824 which is not sufficient in capturing local short-term climate variability. As a result,
825 ~~realized~~updated GPP went outside our ensemble forecasting. On the other hand, parameters and
826 historical information about pool status are more important in long-term pool predictions.
827 Therefore, improvement in long-term pool size predictions cannot be reached by accurate
828 climatic information alone. Instead, it requires accumulation in knowledge related to site history
829 and processes that regulate pool dynamics.

830 Furthermore, reliable forecasting needs understanding of uncertainty sources in addition
831 to the future mean states. Uncertainty and complexity are major reasons that lead to the belief in
832 “computationally irreducible” and low intrinsic predictability of ecological systems ~~[Coreau et~~
833 ~~al., 2010; Beckage et al., 2011; Schindler and Hilborn, 2015]~~[Coreau et al., 2010; Beckage et
834 al., 2011; Schindler and Hilborn, 2015]. Recent advance in computational statistical methods
835 offers a way to formally accounting for various uncertainty sources in ecology ~~[Clark et al.,~~
836 ~~2001; Cressie et al., 2009]~~[Clark et al., 2001; Cressie et al., 2009]. And the Bayesian approach
837 embedded in EcoPAD (v1.0) brings the opportunity to understand and communicate forecasting
838 uncertainty. Our case study revealed that forcing uncertainty is more important in flux-based
839 predictions while parameter uncertainty is more critical in pool-based predictions. Actually, how

840 ~~forecasting~~ uncertainty ~~in carbon forecasting~~ changes with time, what are the dominate
841 ~~sources~~ contributor of ~~forecasting~~ uncertainty (e.g., parameter, initial condition, model structure,
842 observation errors, forcing ~~etc.~~ ~~under different conditions.~~), how uncertainty sources interact
843 among different components, ~~or to what extent unconstrained parameters affect forecasting~~
844 ~~uncertainty~~ are all valuable questions that can be explored through EcoPAD: (v1.0).

845 4.3 Applications of EcoPAD to manipulative experiments and observation sites

846 Broadly speaking, data-model integration stands to increase the overall precision and
847 accuracy of model-based experimentation [~~Luo et al., 2011b; Niu et al., 2014~~][~~Luo et al., 2011b;~~
848 ~~Niu et al., 2014~~]. Systems for which data have been collected in the field and which are well
849 represented by ecological models therefore have the capacity to receive the highest benefits from
850 EcoPAD (v1.0) to improve forecasts. In a global change context, experimental manipulations
851 including ecosystem responses to changes in precipitation regimes, carbon dioxide
852 concentrations, temperatures, season lengths, and species compositional shifts can now be
853 assimilated into ecosystem models [~~Xu et al., 2006; Gao et al., 2011; Lebauer et al., 2013; Shi et~~
854 ~~al., 2016~~][~~Xu et al., 2006; Gao et al., 2011; Lebauer et al., 2013; Shi et al., 2016~~]. Impacts of
855 these global change factors on carbon cycling and ecosystem functioning can now be measured
856 in a scientifically transparent and verifiable manner. This leads to ecosystem ~~modeling~~modelling
857 of systems and processes that can obtain levels of confidence that lend credibility with the public
858 to the science's forward progress toward forecasting and predicting [~~Clark et al., 2001~~][~~Clark et~~
859 ~~al., 2001~~]. These are the strengths of a widely-available interface devoted to data-model
860 integration towards better forecasting.

861 The data-model integration framework of EcoPAD (v1.0) creates a smart interactive
862 model-experiment (ModEx) system. ModEx has the capacity to form a feedback loop in which

863 field experiment guides [modelingmodelling](#) and [modelingmodelling](#) influences experimental
864 focus [[Luo et al., 2011a](#)][[Luo et al., 2011a](#)]. We demonstrated how EcoPAD ([v1.0](#)) works hand-
865 in-hand between [modelersmodellers](#) and experimenters in the life-cycle of the SPRUCE project.
866 Field experiment from SPRUCE community provides basic data to set up the ecosystem model
867 and update model parameters recursively, while the forecasting from ecosystem
868 [modelingmodelling](#) informs experimenters the potential key mechanisms that regulate ecosystem
869 dynamics and help experimenters to question and understand their measurements. The EcoPAD-
870 SPRUCE system operates while experimenters are making measurements or planning for future
871 researches. Information is constantly fed back between [modelersmodellers](#) and experimenters,
872 and simultaneous efforts from both parties illustrate how communications between model and
873 data advance and shape our understanding towards better forecasts during the lifecycle of a
874 scientific project. ModEx can be ~~easily~~ extended to other experimental systems to: 1, predict
875 what might be an ecosystem's response to treatments once experimenter selected a site and
876 decided the experimental plan; 2, assimilate data experimenters are collecting along the
877 experiment to constrain model predictions; 3, project what an ecosystem's responses may likely
878 be in the rest of the experiment; 4, tell experimenters what are those important datasets
879 experimenters may want to collect in order to understand the system; 5, periodically updates the
880 projections; and 6, improve the models, the data assimilation system, and field experiments
881 during the process.

882 In addition to the manipulative experimental, the data assimilation system of EcoPAD
883 ([v1.0](#)) can be used for automated model calibration for FLUXNET sites or other observation
884 networks, such as the NEON and LTER [~~[Johnson et al., 2010; Robertson et al., 2012](#)~~][[Johnson](#)
885 [et al., 2010; Robertson et al., 2012](#)]. The application of EcoPAD ([v1.0](#)) at FLUXNET, NEON or

Formatted: Font color: Auto

886 LTER sites includes three steps in general. First, build the climate forcing in the suitable formats
887 of EcoPAD (v1.0) from the database of each site; Second, collect the prior information (include
888 observations of state variables) in the data assimilation system from FLUXNET, NEON or
889 LTER sites; Third, incorporate the forcing and prior information into EcoPAD (v1.0), and then
890 run the EcoPAD (v1.0) with the dynamic data assimilation system. Furthermore, facing the
891 proposed continental scale ecology study [Schimel, 2014][Schimel, 2011], EcoPAD (v1.0) once
892 properly applied could also help evaluate and optimize field deployment of environmental
893 sensors and supporting cyberinfrastructure, that will be necessary for larger, more complex
894 environmental observing systems being planned in the US and across different continents.
895 Altogether, with its milestone concept, EcoPAD (v1.0) benefits from observation and
896 ~~modeling~~modelling and at the same time advances both observation and ~~modeling~~modelling of
897 ecological studies.

Formatted: Font color: Auto

898 4.4 Future developments

899 As we indicated, EcoPAD (v1.0) will expand as time goes on. The system is designed to
900 incorporate multiple ~~biogeochemical~~process-based models, diverse data assimilation techniques
901 and various ~~ecosystem~~ecological state variables-for different ecosystems. Case studies presented
902 in earlier sections are based primarily on one model. A multiple (or ensemble) model approach is
903 helpful in tracking uncertainty sources from our process understanding. With rapid evolving
904 ecological knowledge, emerging models with different hypotheses, such as the microbial-enzyme
905 model [Wieder et al., 2013][Wieder et al., 2013], enhance our capacity in ecological prediction
906 but can also benefit from rapid tests against data if incorporated into EcoPAD (v1.0). In addition
907 to MCMC [Braswell et al., 2005; Xu et al., 2006][Braswell et al., 2005; Xu et al., 2006], a
908 variety of data assimilation techniques have been recently applied to improve models for

909 ecological forecasting, such as the EnKF ~~[Gao et al., 2014]~~[\[Gao et al., 2011\]](#), Genetic Algorithm
910 ~~[Zhou and Luo, 2008]~~[\[Zhou and Luo, 2008\]](#) and 4-d variational assimilation ~~[Peylin et al.,~~
911 ~~2016]~~[\[Peylin et al., 2016\]](#). Future development will incorporate different optimization techniques
912 to offer users the option to search for the best model parameters by selecting and comparing the
913 possibly best method for their specific ~~studystudies~~. We focus mostly on carbon related state
914 variables in the SPRUCE example, and the data assimilation system in EcoPAD ([v1.0](#)) needs to
915 include more observed variables for constraining model parameters. For example, the NEON
916 sites not only provide measured ecosystem CO₂ fluxes and soil carbon stocks, but also resources
917 (e.g., GPP/Transpiration for water and GPP/intercepted PAR for light) use efficiency ~~[Johnson et~~
918 ~~al., 2010]~~[\[Johnson et al., 2010\]](#).

919 With these improvements, one goal of ~~the~~EcoPAD ([v1.0](#)) is to enable the research
920 community to ~~run models~~[understand and reduce forecasting uncertainties from different sources](#)
921 and forecast various aspects of future biogeochemical [and ecological](#) changes as data
922 ~~becomes~~[become](#) available. [The example of Jiang et al. \[2018\] partitioned forecasting uncertainty](#)
923 [from forcings and parameters. An exhaustive understanding of forecasting uncertainty in ecology](#)
924 [need to also consider model structures, data assimilation schemes as well as different ecological](#)
925 [state variables. Researchers interested in creating their own multiple model and/or multiple](#)
926 [assimilation scheme version of EcoPAD \(v1.0\) can start from the GitHub repository](#)
927 [\(https://github.com/ou-ecolab \)](https://github.com/ou-ecolab) where the source code of the EcoPAD (v1.0) workflow is
928 [archived. To add a new variable that is not forecasted in the EcoPAD-SPRUCE example, it](#)
929 [requires modellers and experimenters to work together to understand their process-based model,](#)
930 [their observations and how messaging works in the workflow of EcoPAD \(v1.0\) following the](#)
931 [example of EcoPAD-SPRUCE. To add a new model or a new data assimilation scheme for](#)

Formatted: Font color: Custom Color(RGB(26,26,26))

932 variables that are forecasted in EcoPAD-SPRUCE, researchers need to create additional dockers
933 and mount them to the existing workflow with the knowledge of how information are passed
934 within the workflow.

935 The power of EcoPAD (v1.0) not only lies in its scientific values, but also in the potential
936 service it can bring to the society. Forecasting with carefully quantified uncertainty is helpful in
937 providing support for natural resource manager and policy maker ~~[Clark et al., 2001]~~ [Clark et
938 al., 2001]. It is always difficult to bring the complex mathematical ecosystem models to the
939 general public, which creates a gap between current scientific advance and public awareness.

940 The web-based interface from EcoPAD (v1.0) makes ~~modeling~~ modelling as easy as possible
941 without losing the connection to the mathematics behind the models. It will greatly transform
942 environmental education and encourage citizen science ~~[Miller-Rushing et al., 2012; Kobori et~~
943 ~~al., 2016]~~ [Miller-Rushing et al., 2012; Kobori et al., 2016] in ecology and climate change with
944 future outreach activities to broadcast the EcoPAD (v1.0) platform.

945 **5 Conclusion**

946 The fully interactive web-based Ecological Platform for Assimilating Data (EcoPAD)
947 into models aims to promote data-model integration towards predictive ecology through bringing
948 the complex ecosystem model and data assimilation techniques ~~easily~~-accessible to different
949 audience. It is supported by meta-databases of biogeochemical variables, libraries of modules of
950 process models, toolbox of inversion techniques and ~~easily~~the scalable scientific workflow.

951 Through these components, it automates data management, model simulation, data assimilation,
952 ecological forecasting, and result visualization, providing an open, convenient, transparent,
953 flexible, scalable, traceable and readily portable platform to systematically conduct data-model
954 integration towards better ecological forecasting.

Formatted: Font color: Auto

955 We illustrated several of its functionalities through the Spruce and Peatland Responses
956 Under Climatic and Environmental change (SPRUCE) experiment. The iterative forecasting
957 approach from EcoPAD-SPRUCE through the prediction-question-discussion-adjustment-
958 prediction cycle and extensive communication between model and data creates a new paradigm
959 to best inform forecasting. In addition to forecasting, EcoPAD enables interactive web-based
960 approach to conduct model simulation, estimate model parameters or state variables, quantify
961 uncertainty of estimated parameters and projected states of ecosystems, evaluate model
962 structures, and assess sampling strategies. Altogether, EcoPAD-SPRUCE creates a smart
963 interactive model-experiment (ModEx) system from which experimenters can know what an
964 ecosystem's response might be at the beginning of their experiments, constrain models through
965 collected measurements, predict ecosystem's response in the rest of the experiments, adjust
966 measurements to better understand their system, periodically update projections and improve
967 models, the data assimilation system, and field experiments.

968 Specifically to forecasting carbon dynamics, EcoPAD-SPRUCE revealed that better
969 forecasting relies on improvements in model structure, parameterization and accurate external
970 forcing. Accurate external forcing is critical for short-term flux-based carbon predictions while
971 right process understanding, parameterization and historical information are essential for long-
972 term pool-based predictions. In addition, EcoPAD provides an avenue to disentangle different
973 sources of uncertainties in carbon cycling studies and to provide reliable forecasts with
974 accountable uncertainties.

975

976 **Code availability:**

977 EcoPAD portal is available at https://ecolab.nau.edu/ecopad_portal/ and code is provided at the
978 GitHub repository (<https://github.com/ou-ecolab>).

979 **Data availability:**

980 Relevant data for this manuscript is available at the SPRUCE project webpage
981 (<https://mnspruce.ornl.gov/>) and the EcoPAD web portal (https://ecolab.nau.edu/ecopad_portal/
982). Additional data can be requested from the corresponding author.

983 **Competing interests:**

984 The authors declare that they have no conflict of interest.

985 **Acknowledgement:**

986 SPRUCE components of this work (PJH, DMR) are based upon work supported by the U.S.
987 Department of Energy, Office of Science, Office of Biological and Environmental Research. Oak
988 Ridge National Laboratory is managed by UT-Battelle, LLC, for the U.S. Department of Energy
989 under contract DE-AC05-00OR22725.

990

991 **Literature Cited**

992 Ahlstrom, A., G. Schurgers, A. Arneth, and B. Smith (2012), Robustness and uncertainty in
993 terrestrial ecosystem carbon response to CMIP5 climate change projections,
994 Environmental Research Letters, 7(4), doi:10.1088/1748-9326/7/4/044008
995 Anderson, J., T. Hoar, K. Raeder, H. Liu, N. Collins, R. Torn, and A. Avellano (2009), The data
996 assimilation research testbed A Community Facility, Bulletin of the American
997 Meteorological Society, 90(9), 1283-1296, doi:10.1175/2009bams2618.1
998 [Baldocchi, D., E. Falge, L. H. Gu, R. Olson, D. Hollinger, S. Running, P. Anthoni, C. Bernhofer,](#)
999 [K. Davis, R. Evans, J. Fuentes, A. Goldstein, G. Katul, B. Law, X. H. Lee, Y. Malhi, T. Meyers, W.](#)
1000 [Munger, W. Oechel, K. T. P. U. K. Pilegaard, H. P. Schmid, R. Valentini, S. Verma, T. Vesala, K.](#)
1001 [Wilson, and S. Wofsy \(2001\). FLUXNET: A new tool to study the temporal and spatial](#)
1002 [variability of ecosystem-scale carbon dioxide, water vapor, and energy flux densities,](#)
1003 [Bulletin of the American Meteorological Society, 82\(11\), 2415-2434, doi:10.1175/1520-](#)
1004 [0477\(2001\)082<2415:fantts>2.3.co;2](#)
1005 Ball, J. T., I. E. Woodrow, and J. A. Berry (1987), A model predicting stomatal conductance
1006 and its contribution to the control of photosynthesis under different environmental

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

1007 conditions, in *Progress in Photosynthesis Research*, edited by J. Biggens, pp. 221– 224,
1008 Martinus Nijhoff, Zoetermeer, Netherlands.

1009 Bastiaanssen, W. G. M., and S. Ali (2003), A new crop yield forecasting model based on
1010 satellite measurements applied across the Indus Basin, Pakistan, *Agriculture Ecosystems &*
1011 *Environment*, 94(3), 321-340, doi:10.1016/s0167-8809(02)00034-8

1012 Beckage, B., L. J. Gross, and S. Kauffman (2011), The limits to prediction in ecological
1013 systems, *Ecosphere*, 2(11), doi:10.1890/es11-00211.1

1014 [Bloom, A. A., J. F. Exbrayat, I. R. van der Velde, L. Feng, and M. Williams \(2016\). The decadal](#)
1015 [state of the terrestrial carbon cycle: Global retrievals of terrestrial carbon allocation, pools,](#)
1016 [and residence times. *Proceedings of the National Academy of Sciences of the United States*](#)
1017 [of America, 113\(5\), 1285-1290, doi:10.1073/pnas.1515160113](#)

1018 Botkin, D. B., H. Saxe, M. B. Araujo, R. Betts, R. H. W. Bradshaw, T. Cedhagen, P. Chesson, T. P.
1019 Dawson, J. R. Etterson, D. P. Faith, S. Ferrier, A. Guisan, A. S. Hansen, D. W. Hilbert, C. Loehle,
1020 C. Margules, M. New, M. J. Sobel, and D. R. B. Stockwell (2007), Forecasting the effects of
1021 global warming on biodiversity, *Bioscience*, 57(3), 227-236, doi:10.1641/b570306

1022 Braswell, B. H., W. J. Sacks, E. Linder, and D. S. Schimel (2005), Estimating diurnal to annual
1023 ecosystem parameters by synthesis of a carbon flux model with eddy covariance net
1024 ecosystem exchange observations, *Global Change Biology*, 11(2), 335-355,
1025 doi:10.1111/j.1365-2486.2005.00897.x

1026 Clark, J. S., S. R. Carpenter, M. Barber, S. Collins, A. Dobson, J. A. Foley, D. M. Lodge, M.
1027 Pascual, R. Pielke, W. Pizer, C. Pringle, W. V. Reid, K. A. Rose, O. Sala, W. H. Schlesinger, D. H.
1028 Wall, and D. Wear (2001), Ecological forecasts: An emerging imperative, *Science*,
1029 293(5530), 657-660, doi:10.1126/science.293.5530.657

1030 Clark, J. S., M. Lewis, J. S. McLachlan, and J. HilleRisLambers (2003), Estimating population
1031 spread: What can we forecast and how well?, *Ecology*, 84(8), 1979-1988, doi:10.1890/01-
1032 0618

1033 Cook, B. I., T. M. Smith, and M. E. Mann (2005), The North Atlantic Oscillation and regional
1034 phenology prediction over Europe, *Global Change Biology*, 11(6), 919-926,
1035 doi:10.1111/j.1365-2486.2005.00960.x

1036 Corbet, S. A., N. M. Saville, M. Fussell, O. E. PrysJones, and D. M. Unwin (1995), The
1037 competition box: A graphical aid to forecasting pollinator performance, *Journal of Applied*
1038 *Ecology*, 32(4), 707-719, doi:10.2307/2404810

1039 Coreau, A., G. Pinay, J. D. Thompson, P. O. Cheptou, and L. Mermet (2009), The rise of
1040 research on futures in ecology: rebalancing scenarios and predictions, *Ecology Letters*,
1041 12(12), 1277-1286, doi:10.1111/j.1461-0248.2009.01392.x

1042 Coreau, A., S. Treyer, P. O. Cheptou, J. D. Thompson, and L. Mermet (2010), Exploring the
1043 difficulties of studying futures in ecology: what do ecological scientists think?, *Oikos*,
1044 119(8), 1364-1376, doi:10.1111/j.1600-0706.2010.18195.x

1045 Craft, C., J. Clough, J. Ehman, S. Joye, R. Park, S. Pennings, H. Y. Guo, and M. Machmuller
1046 (2009), Forecasting the effects of accelerated sea-level rise on tidal marsh ecosystem
1047 services, *Frontiers in Ecology and the Environment*, 7(2), 73-78, doi:10.1890/070219

1048 Cressie, N., C. A. Calder, J. S. Clark, J. M. V. Hoef, and C. K. Wikle (2009), Accounting for
1049 uncertainty in ecological analysis: the strengths and limitations of hierarchical statistical
1050 modeling, *Ecological Applications*, 19(3), 553-570, doi:10.1890/07-0744.1

Formatted: Space Before: 0 pt

1051 Crozier, L. G., R. W. Zabel, and A. F. Hamlett (2008), Predicting differential effects of climate
1052 change at the population level with life-cycle models of spring Chinook salmon, *Global*
1053 *Change Biology*, 14(2), 236-249, doi:10.1111/j.1365-2486.2007.01497.x
1054 Dietze, M. C., D. S. Lebauer, and R. Kooper (2013), On improving the communication
1055 between models and data, *Plant Cell and Environment*, 36(9), 1575-1585,
1056 doi:10.1111/pce.12043
1057 Diez, J. M., I. Ibanez, A. J. Miller-Rushing, S. J. Mazer, T. M. Crimmins, M. A. Crimmins, C. D.
1058 Bertelsen, and D. W. Inouye (2012), Forecasting phenology: from species variability to
1059 community patterns, *Ecology Letters*, 15(6), 545-553, doi:10.1111/j.1461-
1060 0248.2012.01765.x
1061 Ellison, A. M. (2004), [Bayesian inference in ecology](https://doi.org/10.1111/j.1461-0248.2004.00603.x), *Ecology Letters*, 7(6), 509-520,
1062 [doi:10.1111/j.1461-0248.2004.00603.x](https://doi.org/10.1111/j.1461-0248.2004.00603.x)
1063 [Ellison, A. M.](https://doi.org/10.1111/j.1461-0248.2004.00603.x) (2010), Repeatability and transparency in ecological research, *Ecology*, 91(9),
1064 2536-2539, doi:10.1890/09-0032.1
1065 Farquhar, G. D., S. V. Caemmerer, and J. A. Berry (1980), A biochemical-model of
1066 photosynthetic CO₂ assimilation in leaves of C₃ species, *Planta*, 149(1), 78-90,
1067 doi:10.1007/bf00386231
1068 Fordham, D. A., H. R. Akcakaya, M. B. Araujo, J. Elith, D. A. Keith, R. Pearson, T. D. Auld, C.
1069 Mellin, J. W. Morgan, T. J. Regan, M. Tozer, M. J. Watts, M. White, B. A. Wintle, C. Yates, and B.
1070 W. Brook (2012), Plant extinction risk under climate change: are forecast range shifts alone
1071 a good indicator of species vulnerability to global warming?, *Global Change Biology*, 18(4),
1072 1357-1371, doi:10.1111/j.1365-2486.2011.02614.x
1073 Gao, C., H. Wang, E. S. Weng, S. Lakshmvivaranan, Y. F. Zhang, and Y. Q. Luo (2011),
1074 Assimilation of multiple data sets with the ensemble Kalman filter to improve forecasts of
1075 forest carbon dynamics, *Ecological Applications*, 21(5), 1461-1473,
1076 Hampton, S. E., C. A. Strasser, J. J. Tewksbury, W. K. Gram, A. E. Budden, A. L. Batcheller, C. S.
1077 Duke, and J. H. Porter (2013), Big data and the future of ecology, *Frontiers in Ecology and*
1078 *the Environment*, 11(3), 156-162, doi:10.1890/120103
1079 Hanson, P. J., J. S. Riggs, W. R. Nettles, J. R. Phillips, M. B. Krassovski, L. A. Hook, L. Gu, A. D.
1080 Richardson, D. M. Aubrecht, D. M. Ricciuto, J. M. Warren, and C. Barbier (2017), Attaining
1081 whole-ecosystem warming using air and deep-soil heating methods with an elevated CO₂
1082 atmosphere, *Biogeosciences*, 14, 861-883, doi:10.5194/bg-14-861-2017
1083 Hararuk, O., J. Y. Xia, and Y. Q. Luo (2014), Evaluation and improvement of a global land
1084 model against soil carbon data using a Bayesian Markov chain Monte Carlo method, *Journal*
1085 *of Geophysical Research-Biogeosciences*, 119(3), 403-417, doi:10.1002/2013jg002535
1086 Hare, J. A., M. A. Alexander, M. J. Fogarty, E. H. Williams, and J. D. Scott (2010), Forecasting
1087 the dynamics of a coastal fishery species using a coupled climate-population model,
1088 *Ecological Applications*, 20(2), 452-464, doi:10.1890/08-1863.1
1089 Huang, Y., J. Jiang, S. Ma, D. Ricciuto, P. J. Hanson, and Y. Luo (2017), Soil thermal dynamics,
1090 snow cover and frozen depth under five temperature treatments in an ombrotrophic bog:
1091 Constrained forecast with data assimilation, *Journal of Geophysical Research:*
1092 *Biogeosciences*, doi:10.1002/2016JG003725
1093 Jiang, J., Y. Huang, S. Ma, M. Stacy, Z. Shi, D. M. Ricciuto, P. J. Hanson, and Y. Luo (2018),
1094 Forecasting responses of a northern peatland carbon cycle to elevated CO₂ and a gradient
1095 of experimental warming, *Journal of Geophysical Research: Biogeosciences*,
1096 doi:10.1002/2017jg004040

Formatted: Space Before: 0 pt

1097 Johnson, B. R., T. U. Kampe, and M. Kuester (2010), Development of airborne remote
 1098 sensing instruments for NEON, paper presented at SPIE Optical Engineering+
 1099 Applications, International Society for Optics and Photonics.
 1100 Kearney, M. R., B. A. Wintle, and W. P. Porter (2010), Correlative and mechanistic models of
 1101 species distribution provide congruent forecasts under climate change, *Conservation*
 1102 *Letters*, 3(3), 203-213, doi:10.1111/j.1755-263X.2010.00097.x
 1103 Keenan, T. F., E. Davidson, A. M. Moffat, W. Munger, and A. D. Richardson (2012), Using
 1104 model-data fusion to interpret past trends, and quantify uncertainties in future projections,
 1105 of terrestrial ecosystem carbon cycling, *Global Change Biology*, 18(8), 2555-2569,
 1106 doi:10.1111/j.1365-2486.2012.02684.x
 1107 Kobori, H., J. L. Dickinson, I. Washitani, R. Sakurai, T. Amano, N. Komatsu, W. Kitamura, S.
 1108 Takagawa, K. Koyama, T. Ogawara, and A. J. Miller-Rushing (2016), Citizen science: a new
 1109 approach to advance ecology, education, and conservation, *Ecological Research*, 31(1), 1-
 1110 19, doi:10.1007/s11284-015-1314-y
 1111 Kolka, R. K., S. D. Sebestyen, E. S. Verry, and K. N. Brooks (2011), *Peatland biogeochemistry*
 1112 *and watershed hydrology at the Marcell Experimental Forest*, 488 pp., CRC Press Boca Raton
 1113 Lebauer, D. S., D. Wang, K. T. Richter, C. C. Davidson, and M. C. Dietze (2013), Facilitating
 1114 feedbacks between field measurements and ecosystem models, *Ecological Monographs*,
 1115 83(2), 133-154, doi:10.1890/12-0137.1
 1116 Liang, J. Y., D. J. Li, Z. Shi, J. M. Tiedje, J. Z. Zhou, E. A. G. Schuur, K. T. Konstantinidis, and Y. Q.
 1117 Luo (2015), Methods for estimating temperature sensitivity of soil organic matter based on
 1118 incubation data: A comparative evaluation, *Soil Biology & Biochemistry*, 80, 127-135,
 1119 doi:10.1016/j.soilbio.2014.10.005
 1120 Ludascher, B., I. Altintas, C. Berkley, D. Higgins, E. Jaeger, M. Jones, E. A. Lee, J. Tao, and Y.
 1121 Zhao (2006), Scientific workflow management and the Kepler system, *Concurrency and*
 1122 *Computation-Practice & Experience*, 18(10), 1039-1065, doi:10.1002/cpe.994
 1123 Luo, Y. Q., and J. F. Reynolds (1999), Validity of extrapolating field CO₂ experiments to
 1124 predict carbon sequestration in natural ecosystems, *Ecology*, 80(5), 1568-1583,
 1125 doi:10.1890/0012-9658(1999)080[1568:voefce]2.0.co;2
 1126 Luo, Y. Q., S. Q. Wan, D. F. Hui, and L. L. Wallace (2001), Acclimatization of soil respiration to
 1127 warming in a tall grass prairie, *Nature*, 413(6856), 622-625, doi:10.1038/35098065
 1128 Luo, Y. Q., J. Melillo, S. L. Niu, C. Beier, J. S. Clark, A. T. Classen, E. Davidson, J. S. Dukes, R. D.
 1129 Evans, C. B. Field, C. I. Czimczik, M. Keller, B. A. Kimball, L. M. Kueppers, R. J. Norby, S. L.
 1130 Pelini, E. Pendall, E. Rastetter, J. Six, M. Smith, M. G. Tjoelker, and M. S. Torn (2011a),
 1131 Coordinated approaches to quantify long-term ecosystem dynamics in response to global
 1132 change, *Global Change Biology*, 17(2), 843-854, doi:10.1111/j.1365-2486.2010.02265.x
 1133 Luo, Y. Q., K. Ogle, C. Tucker, S. F. Fei, C. Gao, S. LaDeau, J. S. Clark, and D. S. Schimel (2011b),
 1134 Ecological forecasting and data assimilation in a data-rich era, *Ecological Applications*,
 1135 21(5), 1429-1442,
 1136 Luo, Y. Q., A. Ahlstrom, S. D. Allison, N. H. Batjes, V. Brovkin, N. Carvalhais, A. Chappell, P.
 1137 Ciais, E. A. Davidson, A. C. Finzi, K. Georgiou, B. Guenet, O. Hararuk, J. W. Harden, Y. J. He, F.
 1138 Hopkins, L. F. Jiang, C. Koven, R. B. Jackson, C. D. Jones, M. J. Lara, J. Y. Liang, A. D. McGuire,
 1139 W. Parton, C. H. Peng, J. T. Randerson, A. Salazar, C. A. Sierra, M. J. Smith, H. Q. Tian, K. E. O.
 1140 Todd-Brown, M. Torn, K. J. van Groenigen, Y. P. Wang, T. O. West, Y. X. Wei, W. R. Wieder, J.
 1141 Y. Xia, X. Xu, X. F. Xu, and T. Zhou (2016), Toward more realistic projections of soil carbon

1142 dynamics by Earth system models, *Global Biogeochemical Cycles*, 30(1), 40-56,
 1143 doi:10.1002/2015gb005239
 1144 Luo, Y. Q. (2017), Transient dynamics of terrestrial carbon storage: mathematical
 1145 foundation and its applications,
 1146 Ma, S., J. Jiang, Y. Huang, D. Ricciuto, P. J. Hanson, and Y. Luo (2017), Data-constrained
 1147 projections of methane fluxes in a Northern Minnesota Peatland in response to elevated
 1148 CO₂ and warming (Accepted), *Journal of Geophysical Research: Biogeosciences*,
 1149 Medlyn, B. E., F. W. Badeck, D. G. G. De Pury, C. V. M. Barton, M. Broadmeadow, R.
 1150 Ceulemans, P. De Angelis, M. Forstreuter, M. E. Jach, S. Kellomaki, E. Laitat, M. Marek, S.
 1151 Philippot, A. Rey, J. Strassmeyer, K. Laitinen, R. Liozon, B. Portier, P. Roberntz, K. Wang,
 1152 and P. G. Jarvis (1999), Effects of elevated CO₂ on photosynthesis in European forest
 1153 species: a meta-analysis of model parameters, *Plant Cell and Environment*, 22(12), 1475-
 1154 1495, doi:10.1046/j.1365-3040.1999.00523.x
 1155 Melillo, J. M., S. D. Frey, K. M. DeAngelis, W. J. Werner, M. J. Bernard, F. P. Bowles, G. Pold, M.
 1156 A. Knorr, and A. S. Grandy (2017), Long-term pattern and magnitude of soil carbon
 1157 feedback to the climate system in a warming world, *Science*, 358(6359), 101-105,
 1158 doi:10.1126/science.aan2874
 1159 Michener, W. K., and M. B. Jones (2012), Ecoinformatics: supporting ecology as a data-
 1160 intensive science, *Trends in Ecology & Evolution*, 27(2), 85-93,
 1161 doi:10.1016/j.tree.2011.11.016
 1162 Miller-Rushing, A., R. Primack, and R. Bonney (2012), The history of public participation in
 1163 ecological research, *Frontiers in Ecology and the Environment*, 10(6), 285-290,
 1164 doi:10.1890/110278
 1165 Moorcroft, P. R. (2006), How close are we to a predictive science of the biosphere?, *Trends*
 1166 *in Ecology & Evolution*, 21(7), 400-407, doi:10.1016/j.tree.2006.04.009
 1167 Mouquet, N., Y. Lagadeuc, V. Devictor, L. Doyen, A. Duputie, D. Eveillard, D. Faure, E. Garnier,
 1168 O. Gimenez, P. Huneman, F. Jabot, P. Jarne, D. Joly, R. Julliard, S. Kefi, G. J. Kergoat, S. Lavorel,
 1169 L. Le Gall, L. Meslin, S. Morand, X. Morin, H. Morlon, G. Pinay, R. Pradel, F. M. Schurr, W.
 1170 Thuiller, and M. Loreau (2015), REVIEW: Predictive ecology in a changing world, *Journal of*
 1171 *Applied Ecology*, 52(5), 1293-1310, doi:10.1111/1365-2664.12482
 1172 Niu, S. L., Y. Q. Luo, M. C. Dietze, T. F. Keenan, Z. Shi, J. W. Li, and F. S. Chapin (2014), The
 1173 role of data assimilation in predictive ecology, *Ecosphere*, 5(5), doi:10.1890/es13-00273.1
 1174 Ong, J. B. S., M. I. C. Chen, A. R. Cook, H. C. Lee, V. J. Lee, R. T. P. Lin, P. A. Tambyah, and L. G.
 1175 Goh (2010), Real-Time Epidemic Monitoring and Forecasting of H1N1-2009 Using
 1176 Influenza-Like Illness from General Practice and Family Doctor Clinics in Singapore, *Plos*
 1177 *One*, 5(4), doi:10.1371/journal.pone.0010036
 1178 Osterweil, L. J., L. A. Clarke, A. M. Ellison, E. Boose, R. Podorozhny, and A. Wise (2010), Clear
 1179 and Precise Specification of Ecological Data Management Processes and Dataset
 1180 Provenance, *Ieee Transactions on Automation Science and Engineering*, 7(1), 189-195,
 1181 doi:10.1109/tase.2009.2021774
 1182 Parton, W. J., J. W. B. Stewart, and C. V. Cole (1988), Dynamics of c, n, p and s in grassland
 1183 soils - a model, *Biogeochemistry*, 5(1), 109-131, doi:10.1007/bf02180320
 1184 Parton, W. J., J. A. Morgan, G. M. Wang, and S. Del Grosso (2007), Projected ecosystem
 1185 impact of the Prairie Heating and CO₂ Enrichment experiment, *New Phytologist*, 174(4),
 1186 823-834, doi:10.1111/j.1469-8137.2007.02052.x

1187 Perretti, C. T., S. B. Munch, and G. Sugihara (2013), Model-free forecasting outperforms the
1188 correct mechanistic model for simulated and experimental data, *Proceedings of the*
1189 *National Academy of Sciences of the United States of America*, 110(13), 5253-5257,
1190 doi:10.1073/pnas.1216076110
1191 Petchey, O. L., M. Pontarp, T. M. Massie, S. Kefi, A. Ozgul, M. Weilenmann, G. M. Palamara, F.
1192 Altermatt, B. Matthews, J. M. Levine, D. Z. Childs, B. J. McGill, M. E. Schaepman, B. Schmid, P.
1193 Spaak, A. P. Beckerman, F. Pennekamp, and I. S. Pearse (2015), The ecological forecast
1194 horizon, and examples of its uses and determinants, *Ecology Letters*, 18(7), 597-611,
1195 doi:10.1111/ele.12443
1196 Peylin, P., C. Bacour, N. MacBean, S. Leonard, P. Rayner, S. Kuppel, E. Koffi, A. Kane, F.
1197 Maignan, F. Chevallier, P. Ciais, and P. Prunet (2016), A new stepwise carbon cycle data
1198 assimilation system using multiple data streams to constrain the simulated land surface
1199 carbon cycle, *Geoscientific Model Development*, 9(9), 3321-3346, doi:10.5194/gmd-9-
1200 3321-2016
1201 Purves, D., J. Scharlemann, M. Harfoot, T. Newbold, D. P. Tittensor, J. Hutton, and S. Emmott
1202 (2013), Time to model all life on Earth, *Nature*, 493(7432), 295-297,
1203 Robertson, G. P., S. L. Collins, D. R. Foster, N. Brokaw, H. W. Ducklow, T. L. Gragson, C. Gries,
1204 S. K. Hamilton, A. D. McGuire, and J. C. Moore (2012), Long-term ecological research in a
1205 human-dominated world, *BioScience*, 62(4), 342-353,
1206 Schaefer, K., C. R. Schwalm, C. Williams, M. A. Arain, A. Barr, J. M. Chen, K. J. Davis, D.
1207 Dimitrov, T. W. Hilton, D. Y. Hollinger, E. Humphreys, B. Poulter, B. M. Raczka, A. D.
1208 Richardson, A. Sahoo, P. Thornton, R. Vargas, H. Verbeeck, R. Anderson, I. Baker, T. A. Black,
1209 P. Bolstad, J. Q. Chen, P. S. Curtis, A. R. Desai, M. Dietze, D. Dragoni, C. Gough, R. F. Grant, L. H.
1210 Gu, A. Jain, C. Kucharik, B. Law, S. G. Liu, E. Lokipitiya, H. A. Margolis, R. Matamala, J. H.
1211 McCaughey, R. Monson, J. W. Munger, W. Oechel, C. H. Peng, D. T. Price, D. Ricciuto, W. J.
1212 Riley, N. Roulet, H. Q. Tian, C. Tonitto, M. Torn, E. S. Weng, and X. L. Zhou (2012), A model-
1213 data comparison of gross primary productivity: Results from the North American Carbon
1214 Program site synthesis, *Journal of Geophysical Research-Biogeosciences*, 117,
1215 doi:10.1029/2012jg001960
1216 Schimel, D. (2011), The era of continental-scale ecology, *Frontiers in Ecology and the*
1217 *Environment*, 9(6), 311-311,
1218 Schindler, D. E., and R. Hilborn (2015), Prediction, precaution, and policy under global
1219 change, *Science*, 347(6225), 953-954, doi:10.1126/science.1261824
1220 Scholze, M., T. Kaminski, P. Rayner, W. Knorr, and R. Giering (2007), Propagating
1221 uncertainty through prognostic carbon cycle data assimilation system simulations, *Journal*
1222 *of Geophysical Research-Atmospheres*, 112(D17), doi:10.1029/2007jd008642
1223 Shi, Z., X. Xu, O. Hararuk, L. F. Jiang, J. Y. Xia, J. Y. Liang, D. J. Li, and Y. Q. Luo (2015),
1224 Experimental warming altered rates of carbon processes, allocation, and carbon storage in
1225 a tallgrass prairie, *Ecosphere*, 6(11), doi:10.1890/es14-00335.1
1226 Shi, Z., Y. H. Yang, X. H. Zhou, E. S. Weng, A. C. Finzi, and Y. Q. Luo (2016), Inverse analysis of
1227 coupled carbon-nitrogen cycles against multiple datasets at ambient and elevated CO₂,
1228 *Journal of Plant Ecology*, 9(3), 285-295, doi:10.1093/jpe/rtv059
1229 Sitch, S., C. Huntingford, N. Gedney, P. E. Levy, M. Lomas, S. L. Piao, R. Betts, P. Ciais, P. Cox,
1230 P. Friedlingstein, C. D. Jones, I. C. Prentice, and F. I. Woodward (2008), Evaluation of the
1231 terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using

1232 five Dynamic Global Vegetation Models (DGVMs), *Global Change Biology*, 14(9), 2015-
1233 2039, doi:10.1111/j.1365-2486.2008.01626.x
1234 Steppe, K., J. S. von der Crone, and D. J. W. Pauw (2016), TreeWatch.net: A Water and
1235 Carbon Monitoring and Modeling Network to Assess Instant Tree Hydraulics and Carbon
1236 Status, *Frontiers in Plant Science*, 7, doi:10.3389/fpls.2016.00993
1237 Stumpf, R. P., M. C. Tomlinson, J. A. Calkins, B. Kirkpatrick, K. Fisher, K. Nierenberg, R.
1238 Currier, and T. T. Wynne (2009), Skill assessment for an operational algal bloom forecast
1239 system, *Journal of Marine Systems*, 76(1-2), 151-161, doi:10.1016/j.jmarsys.2008.05.016
1240 Sugihara, G., R. May, H. Ye, C. H. Hsieh, E. Deyle, M. Fogarty, and S. Munch (2012), Detecting
1241 Causality in Complex Ecosystems, *Science*, 338(6106), 496-500,
1242 doi:10.1126/science.1227079
1243 ~~Tan, Z. X., S. G. Liu, C. A. Johnston, T. R. Loveland, L. L. Tieszen, J. X. Liu, and R. Kurtz (2005),~~
1244 ~~Soil organic carbon dynamics as related to land use history in the northwestern Great~~
1245 ~~Plains, *Global Biogeochemical Cycles*, 19(3), doi:10.1029/2005gb002536~~
1246 Thomas, R. Q., E. B. Brooks, A. L. Jersild, E. Ward, R. H. Wynne, T. J. Albaugh, H. Dinon-
1247 Aldridge, H. E. Burkhart, J. Domec, T. R. Fox, C. A. Gonzalez-Benecke, T. A. Martin, A.
1248 Noormets, D. A. Sampson, and R. O. Teskey (2017), Leveraging 35 years of *Pinus taeda*
1249 research in the southeastern US to constrain forest carbon cycle predictions: regional data
1250 assimilation using ecosystem experiments, *Biogeosciences*, 14, 3525-3547,
1251 Vitolo, C., Y. Elkhatib, D. Reusser, C. J. A. Macleod, and W. Buytaert (2015), Web technologies
1252 for environmental Big Data, *Environmental Modelling & Software*, 63, 185-198,
1253 doi:10.1016/j.envsoft.2014.10.007
1254 Walker, A. P., P. J. Hanson, M. G. De Kauwe, B. E. Medlyn, S. Zaehle, S. Asao, M. Dietze, T.
1255 Hickler, C. Huntingford, C. M. Iversen, A. Jain, M. Lomas, Y. Q. Luo, H. McCarthy, W. J. Parton,
1256 I. C. Prentice, P. E. Thornton, S. S. Wang, Y. P. Wang, D. Warlind, E. S. Weng, J. M. Warren, F. I.
1257 Woodward, R. Oren, and R. J. Norby (2014), Comprehensive ecosystem model-data
1258 synthesis using multiple data sets at two temperate forest free-air CO₂ enrichment
1259 experiments: Model performance at ambient CO₂ concentration, *Journal of Geophysical*
1260 *Research-Biogeosciences*, 119(5), 937-964, doi:10.1002/2013jg002553
1261 Wallenstein, M. D., and E. K. Hall (2012), A trait-based framework for predicting when and
1262 where microbial adaptation to climate change will affect ecosystem functioning,
1263 *Biogeochemistry*, 109(1-3), 35-47, doi:10.1007/s10533-011-9641-8
1264 Wang, Y. P., and R. Leuning (1998), A two-leaf model for canopy conductance,
1265 photosynthesis and partitioning of available energy I: Model description and comparison
1266 with a multi-layered model, *Agricultural and Forest Meteorology*, 91(1-2), 89-111,
1267 doi:10.1016/s0168-1923(98)00061-6
1268 Wang, Y. P., C. M. Trudinger, and I. G. Enting (2009), A review of applications of model-data
1269 fusion to studies of terrestrial carbon fluxes at different scales, *Agricultural and Forest*
1270 *Meteorology*, 149(11), 1829-1842, doi:10.1016/j.agrformet.2009.07.009
1271 Wang, Y. P., R. M. Law, and B. Pak (2010), A global model of carbon, nitrogen and
1272 phosphorus cycles for the terrestrial biosphere, *Biogeosciences*, 7(7), 2261-2282,
1273 doi:10.5194/bg-7-2261-2010
1274 Ward, E. J., E. E. Holmes, J. T. Thorson, and B. Collen (2014), Complexity is costly: a meta-
1275 analysis of parametric and non-parametric methods for short-term population forecasting,
1276 *Oikos*, 123(6), 652-661, doi:10.1111/j.1600-0706.2014.00916.x

Formatted: Space Before: 0 pt

1277 Weng, E. S., and Y. Q. Luo (2008), Soil hydrological properties regulate grassland ecosystem
1278 responses to multifactor global change: A modeling analysis, *Journal of Geophysical*
1279 *Research-Biogeosciences*, 113(G3), doi:10.1029/2007jg000539
1280 Weng, E. S., and Y. Q. Luo (2011), Relative information contributions of model vs. data to
1281 short- and long-term forecasts of forest carbon dynamics, *Ecological Applications*, 21(5),
1282 1490-1505,
1283 Weng, E. S., Y. Q. Luo, C. Gao, and R. Oren (2011), Uncertainty analysis of forest carbon sink
1284 forecast with varying measurement errors: a data assimilation approach, *Journal of Plant*
1285 *Ecology*, 4(3), 178-191, doi:10.1093/jpe/rtr018
1286 Wieder, W. R., G. B. Bonan, and S. D. Allison (2013), Global soil carbon projections are
1287 improved by modelling microbial processes, *Nature Climate Change*, 3(10), 909-912,
1288 doi:10.1038/nclimate1951
1289 Xu, T., L. White, D. F. Hui, and Y. Q. Luo (2006), Probabilistic inversion of a terrestrial
1290 ecosystem model: Analysis of uncertainty in parameter estimation and model prediction,
1291 *Global Biogeochemical Cycles*, 20(2), doi:10.1029/2005gb002468
1292 Zhou, T., and Y. Q. Luo (2008), Spatial patterns of ecosystem carbon residence time and
1293 NPP-driven carbon uptake in the conterminous United States, *Global Biogeochemical*
1294 *Cycles*, 22(3), doi:10.1029/2007gb002939
1295 Zhou, X. H., T. Zhou, and Y. Q. Luo (2012), Uncertainties in carbon residence time and NPP-
1296 driven carbon uptake in terrestrial ecosystems of the conterminous USA: a Bayesian
1297 approach, *Tellus Series B-Chemical and Physical Meteorology*, 64,
1298 doi:10.3402/tellusb.v64i0.17223
1299

1300 **Tables**

1301 Table 1. Discussion stimulated by EcoPAD-SPRUCE forecasting among ~~modelers~~ modellers and
1302 experimenters on how to improve predictions of the relative contribution of different pathways
1303 of methane emissions

	Discussion
1	No strong bubbles are noted at field and a non-observation constrained modeling <u>modelling</u> study at a similar site from another project concluded minor ebullition contribution, which are at odds with TECO result.
2	CH ₄ :CO ₂ ratio might explain the discrepancy. The other modeling <u>modelling</u> study assumed that decomposed C is mainly turned into CO ₂ and a smaller fraction is turned into CH ₄ . The large CH ₄ :CO ₂ ratio at this site may result in higher CH ₄ flux. It seems that the most “flexible” term is ebullition because any “excess” (above saturation) CH ₄ is immediately released to ebullition, while the plant transport term is constrained by vegetation data.
3	Experimental researches on the relative contribution to methane emission from three different pathways are rare.
4	Current available observations include net surface flux of methane from the large collars, incubation data that should represent methane sources within the profile, and gas/DOC profile data that can indicate active zones within the peat profile. What are additional data needed to constrain relative contribution of different pathways?
5	I had always thought that peatlands don’t bubble much, but the super-sensitive GPS measurements found movements of the surface of the GLAP peatlands consistent with degassing events, and subsurface radar images did show layers that were interpreted as bubble-layers.
6	Pore water gas data, perhaps N ₂ or Ar may shed some light on the relative importance of ebullition.
7	It is really hard to accurately distinguish the three pathways. It has to rely on multiple approaches. Particularly for the SPRUCE site, the vegetation cover varies, vegetation species varies. How many channels each species has affect the transport? Meanwhile, the presence of plant (even not vascular plant) will lead to more gas transport, but as bubbles, rather than plant-mediated transport.
8	It depends on model structure and algorithm to simulate diffusion, vascular, and ebullition. Most models assume a threshold to allow ebullition. Diffusion is treated in similar ways as ebullition in some models (most one layer or two layers models). For the multiple layers models, the diffusion occurs from bottom to top mm by mm, layer by layer, therefore, the gas diffusion from top layer to atmosphere is considered the diffusion flux. If that is the case, the time step and wind speed and pressure matter (most models do not consider wind and pressure impacts). Plant transport is really dependent on the parameter for plant species, aerenchyma, etc. The gas transportability of plant is associated with biomass, NPP, or root biomass, seasonality of plant growth, etc. in models. All these differences might cause biases in the final flux.
9	With only the CH ₄ emission data cannot constrain the relative contribution of three pathways. Concentration data in different soil layers may help constrain.
10	Diffusion coefficient calculation in TECO adopts the “three-porosity-model” which is ideal for mineral soil, but may not fit the organic soil. “Millington-Quirk model” for should be a better choice for peat soil.
11	The boundary condition should be taken care of, but it brings in more uncertainties including the wind speed and piston velocity, etc.,
12	CH ₄ emissions captured in static chambers does not include the episodic ebullition events. So (1) the static chambers underestimate the total methane emission and (2) might need to exclude the ebullition pathway when using the observation data to constrain the CH ₄ emission. But this point seems haven't been paid attention to in other models.

1304

1305 **Figure Legends**

1306 **Figure 1** Schema of approaches to forecast future ecological responses from common practice
1307 (the upper panel) and the Ecological Platform for Assimilation of Data (EcoPAD) (bottom
1308 panel). The common practice makes use of observations to develop or calibrate models to make
1309 predictions while the EcoPAD approach advances the common practice through its fully
1310 interactive platform. EcoPAD consists of four major components: experiment/data, model, data
1311 assimilation and the scientific workflow- (green arrows or lines). Data and model are iteratively
1312 integrated through its data assimilation systems to improve forecasting. And its near-real time
1313 forecasting results are shared among research groups through its web interface to guide new data
1314 collections. The scientific workflow enables web-based data transfer from sensors, model
1315 simulation, data assimilation, forecasting, result analysis, visualization and reporting,
1316 encouraging broad user-model interactions especially for the experimenters and the general
1317 public with limited background in ~~modeling-modelling~~. Images from the SPRUCE field
1318 experiments (<https://mnspruce.ornl.gov/>) are used to represent data collection and the flowchart
1319 of TECO model is used to delegate ecological models.

1320 **Figure 2** The data assimilation system inside the Ecological Platform for Assimilation of Data
1321 (EcoPAD) towards better forecasting of terrestrial carbon dynamics

1322 **Figure 3** The scientific workflow of EcoPAD. The workflow wraps ecological models and data
1323 assimilation algorithms with the docker containerization platform. Users trigger different tasks
1324 through the Representational State Transfer (i.e., RESTful) application-programming interface
1325 (API). Tasks are managed through the asynchronous task queue, Celery. Tasks can be executed
1326 concurrently on a single or more worker servers across different scalable IT infrastructures.

1327 MongoDB is a database software that takes charge of data management in EcoPAD and

1328 RabbitMQ is a message broker.

1329

1330 **Figure 4.** ~~Near time forecasting of EcoPAD-SPRUCE. EcoPAD automatically synchronizes real~~
1331 ~~time observations from environmental sensors managed by the SPRUCE experimental~~
1332 ~~communities. Data from observations are assimilated and used to update forecasting. Weekly~~
1333 ~~forecasting results are displayed in the EcoPAD-SPRUCE web portal~~
1334 ~~(http://ecolab.cybercommons.org/ecopad_portal/) as well as sent back to the experimental groups~~
1335 ~~to guide future experimental design and sampling.~~

1336 **Figure 5.** Schema of interactive communication between ~~modelers~~modellers and experimenters
1337 through the prediction-question-discussion-adjustment-prediction cycle to improve ecological
1338 forecasting. The schema is inspired by an episode of experimenter-~~modeler~~modeller
1339 communication stimulated by the EcoPAD-SPRUCE platform. The initial methane model
1340 constrained by static chamber methane measurements was used to predict relative contributions
1341 of three methane emission pathways (i.e., ebullition, plant mediated transportation (PMT) and
1342 diffusion) to the overall methane fluxes under different warming treatments (+ 0 °C, +2.25 °C,
1343 +4.5 °C, +6.75 °C and +9 °C). The initial results indicated a dominant contribution from
1344 ebullition especially under +9 °C which was doubted by experimenters. The discrepancy
1345 stimulated communications between ~~modelers~~modellers and experimenters with detailed
1346 information listed in Table 1. After extensive discussion, the model structure was adjusted and
1347 field observations were ~~reevaluated~~re-evaluated. And a second round of forecasting yielded more
1348 reliable predictions.

1349 **Figure 65.** Posterior distribution of the ratio of CH₄:CO₂ (panel a) and the temperature
1350 sensitivity of methane production (Q_{10_CH4}, panel b) under 5 warming treatments.


1351 **Figure 7.** ~~Realized6. Updated~~ vs. ~~unrealizedun-updated~~ forecasting of gross primary production
1352 (GPP, panels a,c) and soil organic C content (SoilC, panels b,d). The upper panels show 3 series
1353 of forecasting with ~~differentupdated vs. stochastically generated~~ weather forcing. Cyan indicates
1354 forecasting with 100 stochastically generated weather forcing from January 2015 to December
1355 2024 (S1); red corresponds to ~~realizedupdated~~ forecasting with two stages, that is, updating with
1356 measured weather forcing from January 2015 to July 2016 followed by forecasting with 100
1357 stochastically generated weather forcing from August 2016 to December 2024 (S2); and blue
1358 shows ~~realizedupdated~~ forecasting with measured weather forcing from January 2015 to
1359 December 2016 followed by forecasting with 100 stochastically generated weather forcing from
1360 January 2017 to December 2024 (S3). The bottom panels display mismatches between
1361 ~~realizedupdated~~ forecasting (S2,3) and the original ~~unrealizedun-updated~~ forecasting (S1). Red
1362 displays the difference between S2 and S1 (S2-S1) and blue shows discrepancy between S3 and
1363 S1 (S3-S1). Dashed green lines ~~indicatesindicate~~ the start of forecasting with stochastically
1364 generated weather forcing. Note that the left 2 panels are plotted on yearly time-scale and the
1365 right 2 panels show results on monthly time-scale.

1366

1367



1368

1369 **Figure 1**



1370

1371




1372

1373

1374 **Figure 2**

1375



1376

1377


1378

1379


1380

1381 Figure 3

1382



1383




1384

1385


1386

1387 **Figure 4**



1388

1389




1390

1391

1392 **Figure 5**

1393




1394

1395

1396

1397

1398




1399

1400

1401

1402 **Figure 6**

1403



1404


1405

1406

1407

1408

1409 **Figure 7**



1410

1411

Formatted: Font: Times New Roman