

First I sincerely apologize for not having been able to process in due time your manuscript and your detailed responses to the reviewers comments. After a careful reading of your responses, I believe that you have answered for a large part to the main reviewer's comments. Although one reviewer do not recommend the publication of your paper in GMD, I suggest that it is considered for publication after few last revisions. Indeed, you can still improve the manuscript by taking into account more significantly some of the reviewers critics:

**Responses: Thank you for considering our manuscript for publication. Please check below for our further improvements.**

- Shortening the manuscript: As pointed by one reviewer, the manuscript has « quite a few repetitive elements (e.g. the list of elements included in the workflow appears at multiple places) ». I find that these redundancies are still present and that shortening the manuscript would greatly help. For example information on the potential of EcoPad are presented in the results and then re-state in the discussion section. Also some of the concept brought in the introduction are mentioned also again with similar phrasing in the results or discussion (Ex. 1<sup>st</sup> paragraph of the discussion section ; 1st paragraph of section 3.4.1 (case 1) is also redundant with the introduction, etc...).

Please consider decreasing all repetitions that occur in the manuscript in order to make it more concise and thus easier to read. Try to focus the manuscript on what is new by maybe shortening the summary of past experiences described elsewhere. You may also consider grouping the discussion with the results as for some parts the “discussion section” resumes what has been presented in the “result sections”.

**Responses: We carefully considered the redundancy issue of this manuscript. We put one entire section (2.3 Scientific functionality) into the appendix and removed the 1st paragraph of the discussion section as well as the 1st paragraph of section 3.4.1 (case 1). We had a second thought on merging the ‘discussion’ and ‘result’ sections. We still keep separate result and discussion sections as we thought the merged section would be quite long. Alternatively, we condensed the discussion sections by removing the redundancy parts. For example, we removed lines 686-702, 810-814, 838-401, 860-866 (line numbers correspond to tracked version).**

- Technical developments of EcoPAD: As stated by reviewer 1, I also find that an important message of the paper is linked to the “technical implementation of EcoPAD”: how generic EcoPAD is, in order to facilitate the inclusion of other models, other experiments and other data assimilation systems. It is not straightforward to relate model state variables to observations for a meaningful data assimilation. I thus agree that more details on the technical engineering could be provided (how it will facilitate the inclusion of other model, data stream, DA,...), possibly in an appendix in order not to overload the core of the manuscript and even if this is slightly in contradiction to reviewer2 suggestions (i.e. that these technical aspects are not the core of GMD). In your response to reviewer1's comment you insist on the scientific messages of the paper; I do agree but these can be more concise and focused on the most novel parts linked to ecological forecasting. To my mind some statements are relatively general and well recognized by the scientific community while the description of how EcoPAD may become a widely used platform is less clear.

**Responses: We added “Details on adding a new model or data assimilation approach” into the appendix section.**

- Promotion to non-specialist of EcoPAD: Although your response to reviewer 1 comment is solid, I believe that the discussion section do not emphasize on the limits/risks of web-based tools. No need for large changes but few general warnings/self criticisms could be beneficial.

**Responses: We agree with the viewpoint that promotion of the non-specialist use of web-based ecological models should be careful. We added “On one hand the web-based system with open source broadens the user community. On the other hand, it increases the risk of misuse and misinterpretation. We encourage users to be critical and consult system developers to avoid inappropriate application of the system.” (lines 737-740).**

- Note that reviewer1 concern about your expression ““help experimentaters think” is an interesting expression”, is that such expression is quite negative and may imply that experimentaters do not think enough on average!

**Responses: Thanks a lot for the careful thought to help us improve the manuscript. We have removed this kind of expression from previous revision.**

- Figure 7 (now 6) about updated vs forecasted meteorological impact: Although the new caption and text is more clear, I still find that few more details are needed for a non specialist to understand clearly what is done: what is the updated meteorology and how the stochastically generated forcing is done. Please consider providing few additional information so that the set up of the simulations become clearer.

**Responses: We added “ ‘updated’ means the real meteorological forcing monitored from the weather station. We use stochastically generated forcing to represent future meteorological conditions. Future precipitation and air temperature were generated by vector autoregression using historical dataset (1961–2014) monitored by the weather station. PAR, relative humidity and wind speed were randomly sampled from the joint frequency distribution at a given hour each month. Detailed information on weather forcing is available in Jiang et al. [2018]”. (lines 645-650 ) to provide more information about the source of updated meteorology and how the stochastically generated forcing is created. We also added one sentence “ “updated” means the real meteorology forcing monitored from field weather station.” (line 1294-1295) to the figure legend.**

- Grammar and Typo correction (as pointed by reviewer 2) : although you have clear most of them, I still find some typos or grammatical issues that could be cleared with a thorough reading (ex. P22: “SPRUCE is an ongoing project focuses....”)

**Responses: We carefully went through the manuscript and corrected some typos or grammatical issues, such as line 39,65, 69,122, 149, 196, 228, 291,292, 301, 302, 304, 305, 311,312, 387,477, 577 etc.**

- Else I do agree that it is difficult to account for some of reviewer2 comments on the need to discuss more in details why some parameters are not well constrained and in the same time to focus the paper on the concept of EcoPAD. Maybe few more self-critical views on the limits of EcoPAD would help.

**Responses: Thanks for the suggestion of self-critical views. We added “EcoPAD acts as a tool to link model and data, not as a substitution for neither model nor data. Ecological forecasting through EcoPAD relies strongly on theoretical (model) and empirical (data) ecological studies. Questions such as what are major factors regulating temporal variability of methane emissions cannot be directly answered by EcoPAD. How to make use of EcoPAD to inspire breakthroughs in both theoretical and empirical ecological studies worth future exploration.” (Lines 876-884) into the future development section.**

1 **Realised ecological forecast through interactive Ecological Platform for Assimilating Data**  
2 **into model (EcoPAD (v1.0))**  
3

4 Yuanyuan Huang<sup>1,2</sup>, Mark Stacy<sup>3</sup>, Jiang Jiang<sup>1,4</sup>, Nilutpal Sudi<sup>5</sup>, Shuang Ma<sup>1,6</sup>, Volodymyr  
5 Saruta<sup>1,6</sup>, Chang Gyo Jung<sup>1,6</sup>, Zheng Shi<sup>1</sup>, Jianyang Xia<sup>7,8</sup>, Paul J.Hanson<sup>9</sup>, Daniel Ricciuto<sup>9</sup>, Yiqi  
6 Luo<sup>1,6,10</sup>  
7

8 1, Department of Microbiology and Plant Biology, University of Oklahoma, Norman, Oklahoma

9 2, Laboratoire des Sciences du Climat et de l'Environnement, 91191 Gif-sur-Yvette, France

10 3, University of Oklahoma Information Technology, Norman, Oklahoma, USA

11 4, Key Laboratory of Soil and Water Conservation and Ecological Restoration in Jiangsu Province,  
12 Collaborative Innovation Center of Sustainable Forestry in Southern China of Jiangsu Province, Nanjing  
13 Forestry University, Nanjing, Jiangsu, China

14 5, Department of Computer Science, University of Oklahoma, Norman, Oklahoma, USA

15 6, Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff, AZ, USA

16 7, Tiantong National Forest Ecosystem Observation and Research Station, School of Ecological and  
17 Environmental Sciences, East China Normal University, Shanghai 200062, China.

18 8, Research Center for Global Change and Ecological Forecasting, East China Normal  
19 University, Shanghai 200062, China

20 9, Environmental Sciences Division and Climate Change Science Institute, Oak Ridge National  
21 Laboratory, Oak Ridge, Tennessee, USA

22 10, Department of Earth System Science, Tsinghua University, Beijing 100084 China  
23

24 Correspondence: Yuanyuan Huang ([yuanyuanhuang2011@gmail.com](mailto:yuanyuanhuang2011@gmail.com)) and Yiqi Luo  
25 ([Yiqi.Luo@nau.edu](mailto:Yiqi.Luo@nau.edu))

26 **Abstract.** Predicting future changes in ecosystem services is not only highly desirable but also  
27 becomes feasible as several forces (e.g., available big data, developed data assimilation (DA)  
28 techniques, and advanced cyberinfrastructure) are converging to transform ecological research to  
29 quantitative forecasting. To ~~realize~~realise ecological forecasting, we have developed an  
30 Ecological Platform for Assimilating Data (EcoPAD) (v1.0) into models. EcoPAD (v1.0) is a  
31 web-based software system that automates data transfer and processing from sensor networks to  
32 ecological forecasting through data management, model simulation, data assimilation,  
33 forecasting and visualization. It facilitates interactive data-model integration from which model  
34 is recursively improved through updated data while data is systematically refined under the  
35 guidance of model. EcoPAD (v1.0) relies on data from observations, process-oriented models,  
36 DA techniques, and the web-based workflow.

37 We applied EcoPAD (v1.0) to the Spruce and Peatland Responses Under Climatic and  
38 Environmental change (SPRUCE) experiment at North Minnesota. The EcoPAD-SPRUCE  
39 ~~realizes~~realises fully automated data transfer, feeds meteorological data to drive model  
40 simulations, assimilates both manually measured and automated sensor data into Terrestrial  
41 ECOSystem (TECO) model, and recursively forecast responses of various biophysical and  
42 biogeochemical processes to five temperature and two CO<sub>2</sub> treatments in near real-time  
43 (weekly). Forecasting with EcoPAD-SPRUCE has revealed that mismatches in forecasting  
44 carbon pool dynamics are more related to model (e.g., model structure, parameter, and initial  
45 value) than forcing variables, opposite to forecasting flux variables. EcoPAD-SPRUCE  
46 quantified acclimations of methane production in response to warming treatments through  
47 shifted posterior distributions of the CH<sub>4</sub>:CO<sub>2</sub> ratio and temperature sensitivity (Q<sub>10</sub>) of methane  
48 production towards lower values. Different case studies indicated that realistic forecasting of

49 carbon dynamics relies on appropriate model structure, correct parameterization and accurate  
50 external forcing. Moreover, EcoPAD-SPRUCE stimulated active feedbacks between  
51 experimenters and modellers to identify model components to be improved and additional  
52 measurements to be made. It becomes the interactive model-experiment (ModEx) system and  
53 opens a novel avenue for interactive dialogue between modellers and experimenters. Altogether,  
54 EcoPAD (v1.0) acts to integrate multiple sources of information and knowledge to best inform  
55 ecological forecasting.

56

57

58 **Key words:**

59 Data assimilation, SPRUCE, carbon, global change, real time, acclimation, forecast

60

## 61 1. Introduction

62 One ambitious goal of ecology as a science discipline is to forecast states and services of  
63 ecological systems. Forecasting in ecology is not only desirable for scientific advances in this  
64 discipline but also has practical values to guide resource management and decision-making  
65 ~~toward~~towards a sustainable planet ~~earth~~Earth. The practical need for ecological forecasting is  
66 particularly urgent in this rapidly changing world, which is experiencing unprecedented natural  
67 resource depletion, increasing food demand, serious biodiversity crisis, accelerated climate  
68 changes, and widespread pollutions in the air, waters, and soils [Clark *et al.*, 2001; Mouquet *et*  
69 *al.*, 2015]. As a result, a growing number of studies have ~~been~~-reported ~~in the last several~~  
70 ~~decades on~~-forecasting of, e.g., phenology [Diez *et al.*, 2012], carbon dynamics [Gao *et al.*, 2011;  
71 Luo *et al.*, 2016; Thomas *et al.*, 2017], species dynamics [Clark *et al.*, 2003; Kearney *et al.*,  
72 2010], pollinator performance [Corbet *et al.*, 1995], epidemics [Ong *et al.*, 2010], fishery [Hare  
73 *et al.*, 2010], algal bloom [Stumpf *et al.*, 2009], crop yield [Bastiaanssen and Ali, 2003],  
74 biodiversity [Botkin *et al.*, 2007], plant extinction risk [Fordham *et al.*, 2012], and ecosystem  
75 service [Craft *et al.*, 2009]-~~in the last several decades~~. Despite its broad applications, ecological  
76 forecasting is still sporadically practiced and lags far behind demand due to the lack of  
77 infrastructure that enables timely integration of models with data. This paper introduces the fully  
78 interactive infrastructure, the Ecological Platform for Assimilating Data (EcoPAD) (v1.0) into  
79 models, to inform near-time ecological forecasting with iterative data-model integration.

80 Ecological forecasting relies on both models and data. However, currently the ecology  
81 research community has not yet adequately integrated observations with models to inform best  
82 forecast. Forecasts generated from scenario approaches are qualitative and scenarios are often  
83 not based on ecological knowledge [Coreau *et al.*, 2009; Coreau *et al.*, 2010]. Data-driven

84 forecasts using statistical methods are generally limited for extrapolation and sometimes  
85 contaminated by confounding factors [Schindler and Hilborn, 2015]. Recent emergent  
86 mechanism-free non-parametric approach, which depends on the statistical pattern extracted  
87 from data, is reported to be promising for short-term forecast [Sugihara et al., 2012; Perretti et  
88 al., 2013; Ward et al., 2014], but has limited capability in long-term prediction due to the lack of  
89 relevant ecological mechanisms. Process-based models provide the capacity in long-term  
90 prediction and the flexibility in capturing short-term dynamics on the basis of mechanistic  
91 understanding [Coreau et al., 2009; Purves et al., 2013]. Wide applications of process-based  
92 models are limited by their often complicated numerical structure and sometimes unrealistic  
93 parameterization [Moorcroft, 2006]. The complex and uncertain nature of ecology precludes  
94 practice of incorporating as many processes as possible into mechanistic models. Our current  
95 incomplete knowledge about ecological systems or unrepresented processes under novel  
96 conditions is partly reflected in model parameters which are associated with large uncertainties.  
97 Good forecasting therefore requires effective communication between process-based models and  
98 data to estimate realistic model parameters and capture context-dependent ecological  
99 phenomena.

100         Data-model fusion, or data-model integration, is an important step to combine models  
101 with data. But previous data-model integration activities have mostly been done in an *ad hoc*  
102 manner instead of being interactive. For example, data from a network of eddy covariance flux  
103 tower sites across United States and Canada was compared with gross primary productivity  
104 (GPP) estimated from different models [Schaefer et al., 2012]. Luo and Reynolds [1999] used a  
105 model to examine ecosystem responses to gradual as in the real world vs. step increases in CO<sub>2</sub>  
106 concentration as in elevated CO<sub>2</sub> experiments. Parton et al. [2007] parameterized CO<sub>2</sub> impacts in

107 an ecosystem model with data from a CO<sub>2</sub> experiment in Colorado. Such model-experiment  
108 interactions encounter a few issues: 1) Models are not always calibrated for individual sites and,  
109 therefore, not accurate; 2) It is not very effective because it is usually one-time practice without  
110 many iterative processes between experimenters and modellers [Dietze *et al.*, 2013; Lebauer *et*  
111 *al.*, 2013]; 3) It is usually unidirectional as data is normally used to train models while the  
112 guidance of model for efficient data collection is limited; and 4) It is not streamlined and could  
113 not be disseminated with common practices among the research community [Dietze *et al.*, 2013;  
114 Lebauer *et al.*, 2013; Walker *et al.*, 2014].

115 A few research groups have developed data assimilation systems to facilitate data-model  
116 integration in a systematic way. For example, data-model integration systems, such as the Data  
117 Assimilation Research Testbed - DART [Anderson *et al.*, 2009] and the Carbon Cycle Data  
118 Assimilation Systems - CCDAS [Scholze *et al.*, 2007; Peylin *et al.*, 2016], combine various data  
119 streams (e.g., FLUXNET data, satellite data and inventory data) with process-based models  
120 through data assimilation algorithms such as the Kalman filter [Anderson *et al.*, 2009] and  
121 variational methods [Peylin *et al.*, 2016]. These data assimilation systems automate model  
122 parameterization and ~~provided~~provide an avenue to systematically improve models through  
123 combining as much data as possible. Data-informed model improvements normally happen after  
124 the ending of a field experiment and the interactive data-model integration is limited as  
125 feedbacks from models to ongoing experimental studies are not adequately realised. In addition,  
126 wide applications of these data assimilation systems in ecological forecasting are constrained by  
127 limited user interactions with its steep learning curve to understand these systems, especially for  
128 experimenters who have limited training in modelling.

129           The web-based technology facilitates interactions. Web-based modelling, which provides  
130 user-friendly interfaces to run models in the background, is usually supported by the scientific  
131 workflow, the sequence of processes through which a piece of work passes from initiation to  
132 completion. For example, TreeWatch.Net has recently been developed to make use of high  
133 precision individual tree monitoring data to parameterize process-based tree models in real-time  
134 and to assess instant tree hydraulics and carbon status with online result visualization [*Steppe et*  
135 *al.*, 2016]. Although the web portal of TreeWatch.Net is currently limited to the purpose of  
136 visualization, it largely broadens the application of data-model integration and strengthens the  
137 interaction between modelling researches and the general public. The Predictive Ecosystem  
138 Analyzer (PEcAn) is a scientific workflow that wraps around different ecosystem models and  
139 manages the flows of information coming in and out of the model [*Lebauer et al.*, 2013]. PEcAn  
140 enables web-based model simulations. Such a workflow has advantages, for example, making  
141 ecological modelling and analysis convenient, transparent, reproducible and adaptable to new  
142 questions [*Lebauer et al.*, 2013], and encouraging user-model interactions. PEcAn uses the  
143 Bayesian meta-analysis to synthesize plant trait data to estimate model parameters and associated  
144 uncertainties, i.e., the prior information for process-based models. Parameter uncertainties are  
145 propagated to model uncertainties and displayed as outputs. It is still not fully interactive in the  
146 way that states are not updated iteratively according to observations and the web-based data  
147 assimilation and then ecological forecasting have not yet been fully realised.

148           The iterative model-data integration provides an approach to constantly improve  
149 ecological forecasting and is an important step especially ~~for realizing~~in realising the near real-  
150 time ecological forecasting. Instead of projecting into future through assimilating observations  
151 only once, the iterative forecasting constantly updates forecasting along with ongoing new data

152 streams or/and improved models. Forecasting is likely to be improved unidirectionally in which  
153 either only models are updated through observations, or only data collections/field  
154 experimentations are improved according to theoretical/model information, but not both.  
155 Ecological forecasting can also be bidirectionally improved so that both models and field  
156 experimentations are optimized hand in hand over time. Although the bidirectional case is rare in  
157 ecological forecasting, the unidirectional iterative forecasting has been reported. One excellent  
158 example of forecasting through dynamically and repeatedly integrating data with models is from  
159 infectious disease studies [Ong *et al.*, 2010; Niu *et al.*, 2014]. Dynamics of infectious diseases  
160 are traditionally captured by Susceptible-Infected-Removed (SIR) models. In the forecasting of  
161 the Singapore H1N1-2009 infections, SIR model parameters and the number of individuals in  
162 each state were updated daily, combining data renewed from local clinical reports. The evolving  
163 of the epidemic related parameters and states were captured through iteratively assimilating  
164 observations to inform forecasting. As a result, the model correctly forecasted the timing of the  
165 peak and declining of the infection ahead of time. Iterative forecasting dynamically integrates  
166 data with model and makes best use of both data and theoretical understandings of ecological  
167 processes.

168 The aim of this paper is to present a fully interactive platform, a web-based Ecological  
169 Platform for Assimilating Data into models (EcoPAD<sub>v1.0</sub>) to best inform ecological  
170 forecasting. The interactive feature of EcoPAD (v1.0) is reflected in the iterative model updating  
171 and forecasting through dynamically integrating models with new observations, bidirectional  
172 feedbacks between experimenters and modellers, and flexible user-model communication  
173 through web-based simulation, data assimilation and forecasting. Such an interactive platform  
174 provides the infrastructure to effectively integrate available resources, from both models and

175 data, modellers and experimenters, scientists and the general public, to improve scientific  
176 understanding of ecological processes, to boost ecological forecasting practice and transform  
177 ecology towards quantitative forecasting.

178 In the following sections, we first describe the system design, ~~and~~ major components ~~and~~  
179 ~~functionality~~ of EcoPAD (v1.0). We then use the Spruce and Peatland Responses Under Climatic  
180 and Environmental change (SPRUCE) experiment [Hanson *et al.*, 2017] as a testbed to elaborate  
181 ~~the functionality and~~ new opportunities brought by the platform. We finally discuss implications  
182 of EcoPAD (v1.0) for better ecological forecasting.

183

## 184 ~~2 EcoPAD: (v1.0): system design, and components, and functionality~~

### 185 ~~2.1 General description: web-based data assimilation and forecast~~

186 EcoPAD (v1.0) ([https://ecolab.nau.edu/ecopad\\_portal/](https://ecolab.nau.edu/ecopad_portal/)) focuses on linking ecological  
187 experiments/data with models and allows easily accessible and reproducible data-model  
188 integration with interactive web-based simulation, data assimilation and forecast capabilities.  
189 Specially, EcoPAD (v1.0) enables the automated near time ecological forecasting which works  
190 hand-in-hand between modellers and experimenters and updates periodically in a manner similar  
191 to the weather forecasting. The system is designed to streamline web request-response, data  
192 management, modelling, prediction and visualization to boost the overall throughput of  
193 observational data, promote data-model communication, inform ecological forecasting and  
194 improve scientific understanding of ecological processes- ([see Appendix for detailed](#)  
195 [functionalities of EcoPAD \(v1.0\)](#)).

196 To ~~realize~~~~realise~~ such data-informed ecological forecasting, the essential components of  
197 EcoPAD (v1.0) include experiments/data, process-based models, data assimilation techniques

198 and the scientific workflow (Figures 1-3). The scientific workflow of EcoPAD (v1.0) that wraps  
199 around ecological models and data assimilation algorithms acts to move datasets in and out of  
200 structured and catalogued data collections (metadata catalog) while leaving the logic of the  
201 ecological models and data assimilation algorithms untouched (Figures 1, 3). Once a user makes  
202 a request through the web browser or command line utilities, the scientific workflow takes  
203 charge of triggering and executing corresponding tasks, be it pulling data from a remote server,  
204 running a particular ecological model, automating forecasting or making the result easily  
205 understandable to users (Figures 1, 3). With the workflow, the system is agnostic to operation  
206 system, environment and programming language and is built to horizontally scale to meet the  
207 demands of the model and the end user community.

208

## 209 **2.2 Components**

### 210 **2.2.1 Data**

211 Data is an important component of EcoPAD (v1.0) and EcoPAD (v1.0) offers systematic data  
212 management to digest diverse data streams. The ‘big data’ ecology generates a large volume of  
213 very different datasets across various scales [Hampton *et al.*, 2013; Mouquet *et al.*, 2015]. These  
214 datasets might have high temporal resolutions, such as those from real time ecological sensors, or  
215 the display of spatial information from remote sensing sources and data stored in the geographic  
216 information system (GIS). These datasets may also include, but are not limited to, inventory data,  
217 laboratory measurements, FLUXNET databases or from long-term ecological networks  
218 [Baldocchi *et al.*, 2001; Johnson *et al.*, 2010; Robertson *et al.*, 2012] . Such data contain  
219 information related to environmental forcing (e.g., precipitation, temperature and radiative  
220 forcing), site characteristics (e.g., soil texture and species composition) and biogeochemical

221 information. Datasets in EcoPAD (v1.0) are derived from other research projects in comma  
222 separated value files or other loosely structured data formats. These datasets are first described  
223 and stored with appropriate metadata via either manual operation or scheduled automation from  
224 sensors. Each project has a separate folder where data are stored. Data are generally separated  
225 into two categories. One is used as boundary conditions for modelling and the other category is  
226 related to observations that are used for data assimilation. Scheduled sensor data are appended to  
227 existing data files with prescribed frequency. Attention is then spent on how the particular  
228 dataset varies over space (x, y) and time (t). When the spatiotemporal variability is understood, it  
229 is then placed in metadata records that allow for query through its scientific workflow.

### 230 **2.2.2 Ecological models**

231 Process-based ecological model is another essential component of EcoPAD (Figure 1). In  
232 this paper, the Terrestrial ECOsystem (TECO) model is applied as a general ecological model for  
233 demonstration purposes since the workflow and data assimilation system of EcoPAD (v1.0) are  
234 relatively independent on the specific ecological model. Linkages among the workflow, data  
235 assimilation system and ecological model are based on messaging. For example, the data  
236 assimilation system generates parameters that are passed to ecological models. The state  
237 variables simulated from ecological models are passed back to the data assimilation system.  
238 Models may have different formulations. As long as they take in the same parameters and  
239 generate the same state variables, they are functionally identical from the “eye” of the data  
240 assimilation system.

241 TECO simulates ecosystem carbon, nitrogen, water and energy dynamics [*Weng and Luo,*  
242 2008; *Shi et al.*, 2016]. The original TECO model has 4 major submodules (canopy, soil water,  
243 vegetation dynamics and soil carbon/nitrogen) and is further extended to incorporate methane

244 biogeochemistry and snow dynamics [Huang et al., 2017; Ma et al., 2017]. As in the global land  
245 surface model CABLE [Wang and Leuning, 1998; Wang et al., 2010], canopy photosynthesis  
246 that couples surface energy, water and carbon fluxes is based on a two-big-leaf model [Wang and  
247 Leuning, 1998]. Leaf photosynthesis and stomatal conductance are based on the common scheme  
248 from Farquhar et al. [1980] and Ball et al. [1987] respectively. Transpiration and associated  
249 latent heat losses are controlled by stomatal conductance, soil water content and the rooting  
250 profile. Evaporation losses of water are balanced between the soil water supply and the  
251 atmospheric demand which is based on the difference between saturation vapor pressure ~~at the~~  
252 ~~temperature of the soil~~ and the actual atmospheric vapor pressure. Soil moisture in different soil  
253 layers is regulated by water influxes (e.g., precipitation and percolation) and effluxes (e.g.,  
254 transpiration and runoff). Vegetation dynamic tracks processes such as growth, allocation and  
255 phenology. Soil carbon/nitrogen module tracks carbon and nitrogen through processes such as  
256 litterfall, soil organic matter (SOM) decomposition and mineralization. SOM decomposition  
257 modelling follows the general form of the Century model [Parton et al., 1988] as in most Earth  
258 system models. SOM is divided into pools with different turnover times (the inverse of  
259 decomposition rates) which are modified by environmental factors such as the soil temperature  
260 and moisture.

### 261 **2.2.3 Data assimilation**

262 ~~Data assimilation is a cutting-edge statistical approach that integrates data with model in~~  
263 ~~a systematic way (Figure 2).~~ Data assimilation is growing in importance as the process-based  
264 ecological models, despite largely simplifying the real systems, are in great need to be complex  
265 enough to address sophisticated ecological issues. These ecological issues are composed of an  
266 enormous number of biotic and abiotic factors interacting with each other. Data assimilation

267 techniques provide a framework to combine models with data to estimate model parameters [*Shi*  
268 *et al.*, 2016], test alternative ecological hypotheses through different model structures [*Liang et*  
269 *al.*, 2015], assess information content of datasets [*Weng and Luo*, 2011], quantify uncertainties  
270 [*Weng et al.*, 2011; *Keenan et al.*, 2012; *Zhou et al.*, 2012], derive emergent ecological  
271 relationships [*Bloom et al.*, 2016], identify model errors and improve ecological predictions [*Luo*  
272 *et al.*, 2011b]- ([Figure 2](#)). Under the Bayesian paradigm, data assimilation techniques treat the  
273 model structure, initial and parameter values as priors that represent our current understanding of  
274 the system. As new information from observations or data becomes available, model parameters  
275 and state variables can be updated accordingly. The posterior distributions of estimated  
276 parameters or state variables are imprinted with information from both the model and the  
277 observation/data as the chosen parameters act to reduce mismatches between observations and  
278 model simulations. Future predictions benefit from such constrained posterior distributions  
279 through forward modelling (Figure A1). As a result, the probability density function of predicted  
280 future states through data assimilation normally has a narrower spread than that without data  
281 assimilation when everything else is equal [*Luo et al.*, 2011b; *Weng and Luo*, 2011; *Niu et al.*,  
282 2014].

283 EcoPAD (v1.0) is open to different data assimilation techniques ~~depending on the~~  
284 ~~ecological questions under study~~ since the scientific workflow of EcoPAD (v1.0) is ~~relatively~~  
285 independent on the specific data assimilation algorithm. For demonstration, the Markov chain  
286 Monte Carlo (MCMC) [*Xu et al.*, 2006] is described in this study.

287 MCMC is a class of sampling algorithms to draw samples from a probability distribution  
288 obtained through constructed Markov Chain to approximate the equilibrium distribution. The  
289 Bayesian based MCMC method takes into account various uncertainty sources which are crucial

290 in interpreting and delivering forecasting results [Clark *et al.*, 2001]. In the application of  
291 MCMC, the posterior distribution of ~~parameters~~ a parameter for given observations is  
292 proportional to the prior distribution of ~~parameters~~ that parameter and the likelihood function  
293 which is linked to the fit/match (or cost function) between model simulations and observations.  
294 EcoPAD (v1.0) currently adopts a batch mode, that is, the cost function is treated as a single  
295 function to be minimized and different observations are standardized by their corresponding  
296 standard deviations [Xu *et al.*, 2006]. For simplicity, we assume uniform distributions in priors,  
297 and Gaussian or multivariate Gaussian distributions in observational errors, which can be  
298 operationally expanded to other specific distribution forms depending on the available  
299 information. Detailed description is available in Xu *et al.* [2006].

#### 300 **2.2.4 Scientific workflow**

301 EcoPAD (v1.0) relies on its scientific workflow to interface with ecological models and  
302 data assimilation algorithms, ~~managing~~ manage diverse data streams, automates iterative  
303 ecological forecasting in response to various user requests. Workflow is a relatively new concept  
304 in the ecology literature but essential to ~~realize~~ realise real or near-real time forecasting. Thus, we  
305 describe it in detail below. The essential components of ~~the~~ a scientific workflow of EcoPAD  
306 (v1.0) include the metadata catalog, web application-programming interface (API), the  
307 asynchronous task/job queue (Celery) and the container-based virtualization platform (Docker).  
308 The workflow system of EcoPAD (v1.0) also provides structured result access and visualization.

##### 309 **2.2.4.1 Metadata catalog and data management**

310 Datasets can be placed and queried in EcoPAD (v1.0) via a common metadata catalog  
311 which allows for effective management of diverse data streams. Calls ~~are common~~ for good  
312 management of current large and heterogeneous ecological datasets are common [Ellison, 2010;

313 *Michener and Jones, 2012; Vitolo et al., 2015*]. Kepler [*Ludascher et al., 2006*] and the Analytic  
314 Web [*Osterweil et al., 2010*] are two example systems that endeavour to provide efficient data  
315 management through the storage of metadata including clear documentation of data provenance.  
316 Similarly to these systems, EcoPAD (v1.0) takes advantage of modern information technology,  
317 especially the metadata catalog, to manage diverse data streams. The EcoPAD (v1.0) metadata  
318 schema includes description of the data product, security, access pattern, and timestamp of last  
319 metadata update *etc.* We use MongoDB (<https://www.mongodb.com/>), a NoSQL database  
320 technology, to manage heterogeneous datasets to make the documentation, query and storage fast  
321 and convenient. Through MongoDB, measured datasets can be easily fed into ecological models  
322 for various purposes such as to initialize the model, calibrate model parameters, evaluate model  
323 structure and drive model forecast. For datasets from real time ecological sensors that are  
324 constantly updating, EcoPAD (v1.0) is set to automatically fetch new data streams with  
325 adjustable frequency ~~depending on~~according to research needs.

#### 326 **2.2.4.2 Web API, asynchronous task queue and docker**

327 The RESTful application-programming interface (API) which can deliver data to a wide  
328 variety of applications is the gateway of EcoPAD (v1.0) and enables a wide array of user-  
329 interfaces and data-dissemination activities. Once a user makes a request, such as through  
330 clicking on relevant buttons from a web browser, the request is passed through the  
331 Representational State Transfer (i.e., RESTful) API to trigger specific tasks. The RESTful API  
332 bridges the talk between the client (e.g., a web browser or command line terminal) and the server  
333 (Figure 3). The API exploits the full functionality and flexibility of the HyperText Transfer  
334 Protocol (HTTP), such that data can be retrieved and ingested from the EcoPAD (v1.0) through  
335 the use of simple HTTP headers and verbs (e.g., GET, PUT, POST, *etc.*). Hence, a user can

336 incorporate summary data from EcoPAD (v1.0) into a website with a single line of html code.  
337 Users will also be able to access data directly through programming environments like R, Python  
338 and Matlab. Simplicity, ease of use and interoperability are among the main advantages of this  
339 API which enables web-based modelling.

340 Celery (<https://github.com/celery/celery>) is an asynchronous task/job queue that runs in  
341 the background (Figure 3). The task queue (i.e., Celery) is a mechanism used to distribute work  
342 across work units such as threads or machines. Celery communicates through messages, and  
343 EcoPAD (v1.0) takes advantage of the RabbitMQ (<https://www.rabbitmq.com/>) to manage  
344 messaging. After the user submits a command, the request or message is passed to Celery via the  
345 RESTful API. These messages may trigger different tasks, which include, but not limited to, pull  
346 data from a remote server where original measurements are located, access data through  
347 metadata catalog, run model simulation with user specified parameters, conduct data assimilation  
348 which recursively updates model parameters, forecast future ecosystem status and post-process  
349 of model results for visualization. The broker inside Celery receives task messages and handles  
350 out tasks to available Celery workers which perform the actual tasks (Figure 3). Celery workers  
351 are in charge of receiving messages from the broker, executing tasks and returning task results.  
352 The worker can be a local or remote computation resource (e.g., the cloud) that has connectivity  
353 to the metadata catalog. Workers can be distributed into different information technology (IT)  
354 infrastructures, which makes EcoPAD (v1.0) workflow expandable. Each worker can perform  
355 different tasks depending on tools installed in each worker. And one task can also be distributed  
356 into different workers. In such a way, EcoPAD (v1.0) workflow enables parallelization and  
357 distributed computation of actual modelling tasks across various IT infrastructures, and is  
358 flexible in implementing additional computational resources by connecting additional workers.

359 Another key feature that makes EcoPAD (v1.0) easily portable and scalable among  
360 different operation systems is the utilization of the container-based virtualization platform, the  
361 docker (<https://www.docker.com/>). Docker can run many applications which rely on different  
362 libraries and environments on a single kernel with its lightweight containerization. Tasks that  
363 execute TECO in different ways are wrapped inside different docker containers that can “talk”  
364 with each other. Each docker container embeds the ecosystem model into a complete filesystem  
365 that contains everything needed to run an ecosystem model: the source code, model input, run  
366 time, system tools and libraries. Docker containers are both hardware-agnostic and platform-  
367 agnostic, and they are not confined to a particular language, framework or packaging system.  
368 Docker containers can be run from a laptop, workstation, virtual machine, or any cloud compute  
369 instance. This is done to support the widely varied number of ecological models running in  
370 various languages (e.g., Matlab, Python, Fortran, C and C++) and environments. In addition to  
371 wrap the ecosystem model into a docker container, software applied in the workflow, such as the  
372 Celery, Rabbitmq and MongoDB, are all lightweight and portable encapsulations through docker  
373 containers. Therefore, the entire EcoPAD (v1.0) is readily portable and applicable in different  
374 environments.

#### 375 **2.2.4.3 Structured result access and visualization**

376 EcoPAD (v1.0) enables structured result storage, access and visualization to track and  
377 analyse data-model fusion practice. Upon the completion of the model task, the model wrapper  
378 code calls a post processing call-back function. This call-back function allows for model specific  
379 data requirements to be added to the model result repository. Each task is associated with a  
380 unique task ID and model results are stored within the local repository that can be queried by the  
381 unique task ID. The store and query of model results are realised via the MongoDB and RESTful

382 API (Figure 3). Researchers are authorized to review and download model results and parameters  
383 submitted for each model run through a web accessible URL (link). EcoPAD (v1.0) webpage  
384 also displays a list of historical tasks (with URL) performed by each user. All current and  
385 historical model inputs and outputs are available to download, including the aggregated results  
386 produced for the graphical web applications. In addition, EcoPAD (v1.0) also provides a task  
387 report that contains all-inclusive recap of ~~submitted~~ parameters-~~submitted~~, task status, and model  
388 outputs with links to all data and graphical results for each task. Such structured result storage  
389 and access make sharing, tracking and referring to modelling studies instant and clear.

390 ~~2.~~

### 391 ~~3 Scientific functionality~~

392 ~~Scientific functionality of EcoPAD (v1.0) includes web-based model simulation,~~  
393 ~~estimating model parameters or state variables, quantifying uncertainty of estimated parameters~~  
394 ~~and projected states of ecosystems, evaluating model structures, assessing sampling strategies~~  
395 ~~and conducting ecological forecasting. These functions can be organized to answer various~~  
396 ~~scientific questions. In addition to the general description in this section, the scientific~~  
397 ~~functionality of EcoPAD (v1.0) is also illustrated through a few case studies in the following~~  
398 ~~sections.~~

399 ~~EcoPAD (v1.0) is designed to perform web-based model simulation, which greatly~~  
400 ~~reduces the workload of traditional model simulation through manual code compilation and~~  
401 ~~execution. This functionality opens various new opportunities for modellers, experimenters and~~  
402 ~~the general public. Model simulation and result analysis are automatically triggered after a click~~  
403 ~~on the web-embedded button (Appendices Figures A2, A3-A6). Users are freed from repeatedly~~  
404 ~~compiling code, running code and writing programs to analyse and display model results. Such~~

Formatted: Font: Bold

405 ease of use has great potential to popularize complex modelling studies that are difficult or  
406 inaccessible for experimenters and the general public. As illustrated through the outreach  
407 activities from the TreeWatch.Net [Steppe *et al.*, 2016], the potential functionality of such web-  
408 based model simulation goes beyond its scientific value as its societal and educational impacts  
409 are critical in solving ecological issues. The web-based model simulation also frees users from  
410 model running environment, platform and software. Users can conduct model simulation and do  
411 analysis as long as they have internet access. For example, ecologists can conduct model  
412 simulation and diagnose the underlying reasons for a sudden increase in methane fluxes while  
413 they are making measurements in the field. Non-ecologists, such as youngsters, can study  
414 ecological dynamics through their phones or tablets while they are waiting for the bus. Resource  
415 managers can make timely assessment of different resource utilization strategies on spot of a  
416 meeting.

417 EcoPAD (v1.0) is backed up by data assimilation techniques, which facilitate inference of  
418 model parameters and states based on observations. Ecology have witnessed a growing number  
419 of studies focusing on parameter estimation using inverse modelling or data assimilation as large  
420 volumes of ecological measurements become available. To satisfy the growing need of model  
421 parameterization through observations, EcoPAD (v1.0) streamlines parameter estimations and  
422 updates. Researchers can review and download files that record parameter values from EcoPAD  
423 (v1.0) result repository. Since these parameters may have different biological, physical or  
424 chemical meanings, the functionality of EcoPAD (v1.0) related to parameter estimations can  
425 potentially embrace diverse subareas in ecology. For example, soil scientists can study the  
426 acclimation of soil respiration to manipulative warming through shifts in the distribution of the  
427 decomposition rate parameter from EcoPAD (v1.0). The threshold parameter beyond which

428 further harvesting of fish might cause a crash of fish stocks can be extracted through fish stock  
429 assessment models and observations if mounted to EcoPAD (v1.0).

430 EcoPAD (v1.0) promotes uncertainty analysis, model structure evaluation and error  
431 identification. One of the advantages of the Bayesian statistics is its capacity in uncertainty  
432 analysis compared to other optimization techniques [Xu *et al.*, 2006; Wang *et al.*, 2009; Zhou *et*  
433 *al.*, 2012]. Bayesian data assimilation (e.g., MCMC) takes into account observation uncertainties  
434 (errors), generates distributions of model parameters and enables tracking of prediction  
435 uncertainties from different sources [Ellison, 2004; Bloom *et al.*, 2016; Jiang *et al.*, 2018].  
436 Uncertainty analysis through data assimilation applied to areas such as ecosystem phenology,  
437 fish life cycle and species migration [Clark *et al.*, 2003; Cook *et al.*, 2005; Crozier *et al.*, 2008;  
438 Luo *et al.*, 2011b], can potentially take advantage of EcoPAD (v1.0) platform to provide critical  
439 information for well-informed decisions in face of pressing global change challenges. In  
440 addition, the archive capacity of EcoPAD (v1.0) facilitates future inter-comparisons among  
441 different models or different versions of the same model to evaluate model structures and to  
442 disentangle structure uncertainties and errors.

443 The realization of both the near time and long-term ecological forecast is one of the key  
444 innovations of EcoPAD (v1.0). Forecasting capability of EcoPAD (v1.0) is supported by  
445 process-based ecological models, multiple observational or experimental data, inverse parameter  
446 estimation and uncertainty quantification through data assimilation, and forward simulation  
447 under future external conditions. The systematically constrained forecast from EcoPAD (v1.0) is  
448 accompanied by uncertainty/confidence estimates to quantify the amount of information that can  
449 actually be utilized from a study. The automated near-time forecast, which is constantly adjusted  
450 once new observational data streams are available, provides experimenters advanced and timely

451 information to assess and adjust experimental plans. For example, with forecasted and displayed  
452 biophysical and biochemical variables, experimenters could know in advance what the most  
453 likely biophysical conditions are. Knowing if the water table may suddenly go aboveground in  
454 response to a high rainfall forecast in the coming week, could allow researcher to emphasize  
455 measurements associated with methane flux. In such a way, experimenters can not only rely on  
456 historical ecosystem dynamics, but also refer to future predictions. Experimenters will benefit  
457 especially from variables that are difficult to track in field due to situations such as harsh  
458 environment, shortage in man power or on instrument limitation.

459 Equally important, EcoPAD (v1.0) creates new avenues to answer classic and novel  
460 ecological questions, for example, the frequently reported acclimation phenomena in ecology.  
461 While growing evidence points to altered ecological functions as organisms adjust to the rapidly  
462 changing world [Medlyn *et al.*, 1999; Luo *et al.*, 2001; Wallenstein and Hall, 2012], traditional  
463 ecological models treat ecological processes less dynamical, as the governing biological  
464 parameters or mechanisms fails to explain such biological shifts. EcoPAD (v1.0) facilitates the  
465 shift of research paradigm from a fixed process representation to a more dynamic description of  
466 ecological mechanisms with constantly updated and archived parameters constrained by  
467 observations under different conditions. Specifically to acclimation, EcoPAD (v1.0) promotes  
468 quantitative evaluations while previous studies remain mostly qualitative [Wallenstein and  
469 Hall, 2012; Shi *et al.*, 2015]. We will further illustrate how EcoPAD (v1.0) can be used to  
470 address different ecological questions in the case studies of the SPRUCE project.

## 472 **3 EcoPAD performance at testbed - SPRUCE**

### 473 **3.1 SPRUCE project overview**

474 EcoPAD (v1.0)- is being applied to the Spruce and Peatland Responses Under Climatic  
475 and Environmental change (SPRUCE) experiment located at the USDA Forest Service Marcell  
476 Experimental Forest (MEF, 47°30.476' N, 93°27.162' W) in northern Minnesota [Kolka *et al.*,  
477 2011]. SPRUCE is an ongoing project that focuses on long-term responses of northern peatland  
478 to climate warming and increased atmospheric CO<sub>2</sub> concentration [Hanson *et al.*, 2017]. At  
479 SPRUCE, ecologists measure various aspects of responses of organisms (from microbes to trees)  
480 and ecological functions (carbon, nutrient and water cycles) to a warming climate. One of the  
481 key features of the SPRUCE experiments is the manipulative deep soil/peat heating (0-3 m) and  
482 whole ecosystem warming treatments (peat + air warmings) which include tall trees (> 4 m)  
483 [Hanson *et al.*, 2017]. Together with elevated atmospheric CO<sub>2</sub> treatments, SPRUCE provides a  
484 platform for exploring mechanisms controlling the vulnerability of organisms, biogeochemical  
485 processes and ecosystems in response to future novel climatic conditions. The SPRUCE peatland  
486 is especially sensitive to future climate change and also plays an important role in feeding back  
487 to future climate change through greenhouse gas emissions as it stores a large amount of soil  
488 organic carbon. Vegetation in the SPRUCE site is dominated by *Picea mariana* (black spruce)  
489 and *Sphagnum spp* (peat moss). The studied peatland also has an understory which include  
490 ericaceous and woody shrubs. There are also a limited number of herbaceous species. The whole  
491 ecosystem warming treatments include a large range of both aboveground and belowground  
492 temperature manipulations (ambient, control plots of +0 °C, +2.25 °C, +4.5 °C, +6.75 °C and +9  
493 °C) in large 115 m<sup>2</sup> open-topped enclosures with elevated CO<sub>2</sub> manipulations (+0 or +500 ppm).  
494 The difference between ambient and +0 °C treatment plots is the open-topped and controlled-  
495 environment enclosure.

496 The SPRUCE project generates a large variety of observational datasets that reflect  
497 ecosystem dynamics from different scales and are available from the project webpage  
498 (<https://mnspruce.ornl.gov/>) and FTP site (<ftp://sprucedata.ornl.gov/>). These datasets come from  
499 multiple sources: half hourly automated sensor records, species surveys, laboratory  
500 measurements, laser scanning images *etc.* Involvements of both modelling and experimental  
501 studies in the SPRUCE project create the opportunity for data-model communication. Datasets  
502 are pulled from SPRUCE archives and stored in the EcoPAD (v1.0) metadata catalog for running  
503 the TECO model, conducting data-model fusion or forecasting. The TECO model has been  
504 applied to simulate and forecast carbon dynamics with productions of CO<sub>2</sub> and CH<sub>4</sub> from  
505 different carbon pools, soil temperature response, snow depth and freeze-thaw cycles at the  
506 SPRUCE site [*Huang et al.*, 2017; *Ma et al.*, 2017; *Jiang et al.*, 2018].

507

### 508 **3.2 EcoPAD-SPRUCE web portal**

509 We assimilate multiple streams of data from the SPRUCE experiment to the TECO  
510 model using the MCMC algorithm, and forecast ecosystem dynamics in both near time and for  
511 the next 10 years. Our forecasting system for SPRUCE is available at  
512 [https://ecolab.nau.edu/ecopad\\_portal/](https://ecolab.nau.edu/ecopad_portal/). From the web portal, users can check our current near-  
513 and long-term forecasting results, conduct model simulation, data assimilation and forecasting  
514 runs, and analyse/visualize model results. Detailed information about the interactive web portal  
515 is provided in the Appendices.

### 516 **3.3 Near time ecosystem forecasting and feedback to experimenters**

517 As part of the forecasting functionality, EcoPAD-SPRUCE automates the near time  
518 (weekly) forecasting with continuously updated observations from SPRUCE experiments (Figure

519 4). We set up the system to automatically pull new data streams every Sunday from the SPRUCE  
520 FTP site that holds observational data and update the forecasting results based on new data  
521 streams. Updated forecasting results for the next week are customized for the SPRUCE  
522 experiments with different manipulative treatments and displayed in the EcoPAD-SPRUCE  
523 portal. At the same time, these results are sent back to SPRUCE communities and displayed  
524 together with near-term observations for experimenter's reference.

### 525 **3.4 New approaches to ecological studies towards better forecasting**

#### 526 **3.4.1 Case 1: Interactive communications among modellers and experimenters**

527 EcoPAD-SPRUCE provides a platform to stimulate interactive communications between  
528 modellers and experimenters. ~~Models require experimental data to constrain initial conditions  
529 and parameters, and to verify model performance. A reasonable model is built upon correct  
530 interpretation of information served by experimenters. Model simulations on the other hand can  
531 expand hypothesis testing, and provide thorough or advanced information to improve field  
532 experiments. Through recursively exchanging information between modellers and experimenters,  
533 both models and field experiments can be improved. As illustrated in Figure 4, through extensive  
534 communication between modellers and experimenters, modellers generate model predictions.  
535 Model predictions provide experimenters advanced information, help experimenters think,  
536 question and understand their experiments. Questions raised by experimenters stimulate further  
537 discussion and communication. Through communication, models or/and measurements are  
538 adjusted. With new measurements or/and strengthened models, a second round of prediction is  
539 highly likely to be improved. As the loop of prediction-question-discussion-adjustment-  
540 prediction goes on, forecasting is informed with best understandings from both data and model.~~

541 through the loop of prediction-question-discussion-adjustment-prediction (Figure 4). We  
542 illustrate how the prediction-question-discussion-adjustment-prediction cycle and stimulation of  
543 modeller-experimenter communication improves ecological predictions through one episode  
544 during the study of the relative contribution of different pathways to methane emissions. An  
545 initial methane model was built upon information (e.g., site characteristics and environmental  
546 conditions) provided by SPRUCE field scientists, taking into account important processes in  
547 methane dynamics, such as production, oxidation and emissions through three pathways (i.e.,  
548 diffusion, ebullition and plant-mediated transportation). The model was used to predict relative  
549 contributions of different pathways to overall methane emissions under different warming  
550 treatments after being constrained by measured surface methane fluxes. Initial forecasting results  
551 which indicated a strong contribution from ebullition under high warming treatments were sent  
552 back to the SPRUCE group. Experimenters doubted about such a high contribution from the  
553 ebullition pathway and a discussion was stimulated. It is difficult to accurately distinguish the  
554 three pathways from field measurements. Field experimenters provided potential avenues to  
555 extract measurement information related to these pathways, while modellers examined model  
556 structure and parameters that may not be well constrained by available field information.  
557 Detailed discussion is provided in Table 1. After extensive discussion, several adjustments were  
558 adopted as a first step to move forward. For example, the three-porosity model that was used to  
559 simulate the diffusion process was replaced by the Millington-Quirk model to more realistically  
560 represent methane diffusions in peat soil; the measured static chamber methane fluxes were also  
561 questioned and scrutinized more carefully to clarify that they did not capture the episodic  
562 ebullition events. Measurements such as these related to pore water gas data may provide  
563 additional inference related to ebullition. The updated forecasting is more reasonable than the

564 initial results although more studies are in need to ultimately quantify methane fluxes from  
565 different pathways.

### 566 **3.4.2 Case 2: Acclimation of ecosystem carbon cycling to experimental manipulations**

567 As a first step, CH<sub>4</sub> static chamber flux measurements were assimilated into TECO to  
568 assess potential acclimation phenomena during methane production under 5 warming treatments  
569 (+0, +2.25, +4.5, +6.75, +9 °C). Initial results indicated a reduction in both the CH<sub>4</sub>:CO<sub>2</sub> ratio and  
570 the temperature sensitivity of methane production based on their posterior distributions (Figure 5).  
571 The mean CH<sub>4</sub>:CO<sub>2</sub> ratio decreased from 0.675 (+0 °C treatment) to 0.505 (+9 °C), while the  
572 temperature sensitivity (Q<sub>10</sub>) for CH<sub>4</sub> production decreased from 3.33 (+0 °C) to 1.22 (+9 °C  
573 treatment). Such shifts quantify potential acclimation of methane production to warming and future  
574 climate warming is likely to have a smaller impact on emission than most of current predictions  
575 that do not take into account of acclimation.

576 Despite these results are preliminary as more relevant datasets are under collection with  
577 current ongoing warming ~~manipulation~~manipulations and measurements, assimilating  
578 observations through EcoPAD (v1.0) provides a quantitative approach to timely assess acclimation  
579 through time. *Melillo et al.* [2017] revealed that the thermal acclimation of the soil respiration in  
580 the Harvard Forest is likely to be phase (time) dependent during their 26-year soil warming  
581 experiment. EcoPAD (v1.0) provides the possibility in tracing the temporal path of acclimation  
582 with its streamlined structure and archive capacity. *Shi et al.* [2015] assimilated carbon related  
583 measurements in a tallgrass prairie into the TECO model to study acclimation after 9-years  
584 warming treatments. They revealed a reduction in the allocation of GPP to shoot, the turnover rates  
585 of the shoot and root carbon pools, and an increase in litter and fast carbon turnovers in response  
586 to warming treatments. Similarly, as time goes on, the SPRUCE experiment will generate more

587 carbon cycling related datasets under different warming and CO<sub>2</sub> treatments, which can be  
588 mounted to EcoPAD (v1.0) to systematically quantify acclimations in carbon cycling through time  
589 in the future.

### 590 **3.4.3 Case 3: Partitioning of uncertainty sources**

591         Uncertainties in ecological studies can come from observations (include forcing that  
592 drives the model), different model structures to represent the real world and the specified model  
593 parameters [*Luo et al.*, 2016]. Previous studies tended to focus on one aspect of the uncertainty  
594 sources instead of disentangling the contribution from different sources. For example, the model  
595 intercomparison projects (MIPs), such as TRENDY, focus on uncertainty caused by different  
596 model structures with prescribed external forcing [*Sitch et al.*, 2008]. *Keenan et al.* [2012] used  
597 data assimilation to constrain parameter uncertainties in projecting Harvard forest carbon  
598 dynamics. *Ahlstrom et al.* [2012] forced one particular vegetation model by 18 sets of forcings  
599 from climate models of the Coupled Model Intercomparison Project Phase 5 (CMIP5), while the  
600 parameter or model structure uncertainty is not taken into account.

601         EcoPAD (v1.0) is designed to provide a thorough picture of uncertainties from multiple  
602 sources especially in carbon cycling studies. Through focusing on multiple instead of one source  
603 of uncertainty, ecologists can allocate resources to areas that cause relative high uncertainty.  
604 Attribution of uncertainties in EcoPAD (v1.0) will rely on an ensemble of ecosystem models, the  
605 data assimilation system and climate forcing with quantified uncertainty. *Jiang et al.* [2018]  
606 focused specifically on the relative contribution of parameter uncertainty vs. climate forcing  
607 uncertainty in forecasting carbon dynamics at the SPRUCE site. Through assimilating the pre-  
608 treatment measurements (2011-2014) from the SPRUCE experiment, *Jiang et al.* [2018]  
609 estimated uncertainties of key parameters that regulate the peatland carbon dynamics. Combined

610 with the stochastically generated climate forcing (e.g., precipitation and temperature), *Jiang et*  
611 *al.* [2018] found external forcing resulted in higher uncertainty than parameters in forecasting  
612 carbon fluxes, but caused lower uncertainty than parameters in forecasting carbon pools.  
613 Therefore, more efforts are required to improve forcing measurements for studies that focus on  
614 carbon fluxes (e.g., GPP), while reductions in parameter uncertainties are more important for  
615 studies in carbon pool dynamics. Despite *Jiang et al.* [2018] does not quantify model structure  
616 uncertainty, the project of incorporating multiple models inside EcoPAD (v1.0) is in progress,  
617 and future uncertainty assessment will benefit from EcoPAD (v1.0) with its systematically  
618 archived model simulation, data assimilation and forecasting.

#### 619 **3.4.4 Case 4: Improving biophysical estimation for better ecological prediction**

620 Carbon cycling studies can also benefit from EcoPAD (v1.0) through improvements in  
621 biophysical estimation. Soil environmental condition is an important regulator of belowground  
622 biological activities and also feeds back to aboveground vegetation growth. Biophysical  
623 variables such as soil temperature, soil moisture, ice content and snow depth, are key predictors  
624 of ecosystem dynamics. After constraining the biophysical module by detailed monitoring data  
625 from the SPRUCE experiment through the data assimilation component of EcoPAD (v1.0),  
626 *Huang et al.* [2017] forecasted the soil thermal dynamics under future conditions and studied the  
627 responses of soil temperature to hypothetical air warming. This study emphasized the importance  
628 of accurate climate forcing in providing robust thermal forecast. In addition, *Huang et al.* [2017]  
629 revealed non-uniform responses of soil temperature to air warming. Soil temperature responded  
630 stronger to air warming during summer compared to winter. And soil temperature increased  
631 more in shallow soil layers compared to deep soils in summer in response to air warming.  
632 Therefore, extrapolating of manipulative experiments based on air warming alone may not

633 reflect the real temperature sensitivity of SOM if soil temperature is not monitored. As robust  
634 quantification of environmental conditions is known to be a first step towards better  
635 understanding of ecological process, improvement in soil thermal predictions through EcoPAD  
636 (v1.0) data assimilation system is helpful in telling apart biogeochemical responses from  
637 environmental uncertainties and also in providing field ecologists beforehand key environmental  
638 conditions.

### 639 **3.4.5 Case 5: How do updated model and data contribute to reliable forecasting?**

640 Through constantly adjusted model and external forcing according to observations and  
641 weekly archived model parameter, model structure, external forcing and forecasting results, the  
642 contribution of model and data updates can therefore be tracked through comparing forecasted  
643 vs. realised simulations. For example, Figure 6 illustrates how updated external forcing  
644 (compared to stochastically generated forcing) and shifts in ecosystem state variables shape  
645 ecological predictions. “updated” means the real meteorological forcing monitored from the  
646 weather station. We use stochastically generated forcing to represent future meteorological  
647 conditions. Future precipitation and air temperature were generated by vector autoregression  
648 using historical dataset (1961–2014) monitored by the weather station. PAR, relative humidity  
649 and wind speed were randomly sampled from the joint frequency distribution at a given hour  
650 each month. Detailed information on weather forcing is available in Jiang et al. [2018]. Similarly  
651 as in other EcoPAD-SPURCE case studies, TECO is trained through data assimilation with  
652 observations from 2011-2014 and is used to forecast GPP and total soil organic carbon content at  
653 the beginning of 2015. For demonstrating purpose, Figure 6 only shows 3 series of forecasting  
654 results instead of updates from every week. Series 1 (S1) records forecasted GPP and soil carbon  
655 with stochastically generated weather forcing from January 2015-December 2024 (Figure 6a,b

656 cyan). Series 2 (S2) records simulated GPP and soil carbon with observed ([updated](#)) climate  
657 forcing\_ from January 2015 to July 2016 and forecasted GPP and soil carbon with stochastically  
658 generated forcing from August 2016 - December 2024 (Figure 6a,b red). Similarly, the  
659 stochastically generated forcing in Series 3 (S3) starts from January 2017 (Figure 6a,b blue). For  
660 each series, predictions were conducted with randomly sampled parameters from the posterior  
661 distributions and stochastically generated forcing. We displayed 100 mean values (across an  
662 ensemble of forecasts with different parameters) corresponding to 100 forecasts with  
663 stochastically generated forcing.

664 GPP is highly sensitive to climate forcing. The differences between the updated (S2, 3)  
665 and initial forecasts (S1) reach almost  $800 \text{ gC m}^{-2} \text{ year}^{-1}$  (Figure 6c). The discrepancy is strongly  
666 dampened in the following 1-2 years. The impact of updated forecasts is close to 0 after  
667 approximately 5 years. However, soil carbon pool shows a different pattern. Soil carbon pool is  
668 increased by less than  $150 \text{ gC m}^{-2}$ , which is relative small compared to the carbon pool size of  
669 *ca.*  $62000 \text{ gC m}^{-2}$ . The impact of updated forecasts grows with time and reaches the highest at  
670 the end of the simulation year 2024. GPP is sensitive to the immediate change in climate forcing  
671 while the updated ecosystem status (or initial value) has minimum impact in the long-term  
672 forecast of GPP. The impact of updated climate forcing is relatively small for soil carbon  
673 forecasts during our study period. Soil carbon is less sensitive to the immediate change of  
674 climate compared to GPP. However, the alteration of system status affects soil carbon forecast  
675 especially in a longer time scale.

676 Since we are archiving updated forecasts every week, we can track the relative  
677 contribution of ecosystem status, forcing uncertainty and parameter distributions to the overall  
678 forecasting patterns of different ecological variables and how these patterns evolve in time. In

679 addition, as growing observations of ecological variables (e.g., carbon fluxes and pool sizes)  
680 become available, it is feasible to diagnose key factors that promote robust ecological forecasting  
681 through comparing the archived forecasts vs. observation and analysing archives of model  
682 parameters, initial values and climate forcing *etc.*

683

#### 684 **4 Discussion**

##### 685 **4.1 The necessity of interactive infrastructure to ~~realizer~~realise ecological forecasting**

686 ~~Substantial increases in data availability from observational and experimental networks,~~  
687 ~~surges in computational capability, advancements in ecological models and sophisticated~~  
688 ~~statistical methodologies and pressing societal need for best management of natural resources~~  
689 ~~have shifted ecology to emphasis more on quantitative forecasts. However, quantitative~~  
690 ~~ecological forecast is still young and our knowledge about ecological forecasting is relatively~~  
691 ~~sparse, inconsistent and disconnected [Luo *et al.*, 2011b; Petchey *et al.*, 2015]. Therefore, both~~  
692 ~~optimistic and pessimistic viewpoints exist on the predictability of ecology [Clark *et al.*, 2001;~~  
693 ~~Beckage *et al.*, 2011; Purves *et al.*, 2013; Petchey *et al.*, 2015; Schindler and Hilborn, 2015].~~  
694 ~~Ecological forecasting is complex and advantages in one single direction, for example,~~  
695 ~~observations alone or statistical methodology alone, is less likely to lead to successful forecasting~~  
696 ~~compared to approaches that effectively integrate improvements from multiple sectors.~~  
697 ~~Unfortunately, realised ecological forecasting that integrates available resources is relative rare~~  
698 ~~due to lack of relevant infrastructures.~~

699 ~~EcoPAD (v1.0) provides such effective infrastructure with its interactive platform that~~  
700 ~~rigorously integrates merits from models, observations, statistical advance, information~~  
701 ~~technology and human resources from experimenters and modellers to best inform ecological~~

702 ~~forecasting, boost forecasting practice and delivery of forecasting results.~~ Interactions enable  
703 exchanging and extending of information so as to benefit from collective knowledge. For  
704 example, manipulative studies will have a much broader impact if the implications of their  
705 results can be extended from the regression between environmental variable and ecosystem  
706 response, such as be integrated into an ecosystem model through model-data communication.  
707 Such an approach will allow gaining information about the processes responsible for ecosystem's  
708 response, constraining models, and making more reliable predictions. Going beyond common  
709 practice of model-data assimilation from which model updating lags far behind observations,  
710 EcoPAD (v1.0) enables iterative model updating and forecasting through dynamically  
711 integrating models with new observations in near real-time. This near real-time interactive  
712 capacity relies on its scientific workflow that automates data management, model simulation,  
713 data simulation and result visualization. The system design encourages thorough interactions  
714 between experimenters and modellers. Forecasting results from SPRUCE were timely shared  
715 among research groups with different background through the web interface. Expertise from  
716 different research groups was integrated to improve a second round of forecasting. Again, thanks  
717 to the workflow, new information or adjustment is incorporated into forecasting efficiently,  
718 making the forecasting system fully interactive ~~and dynamical.~~

719 We also benefit from the interactive EcoPAD (v1.0) platform to broaden user-model  
720 interactions and to broadcast forecasting results. Learning about the ecosystem models and data-  
721 model fusion techniques may lag one's productivity and even discourage learning the modelling  
722 techniques because of their complexity and long learning curve. Because EcoPAD (v1.0) can be  
723 accessed from a web browser and does not require any coding from the user's side, the time lag  
724 between learning the model structure and obtaining model-based results for one's study is

725 minimal, which opens the door for non-modeller groups to “talk” with models. The online  
726 storage of one’s results lowers the risk of data loss. The results of each model run can be easily  
727 tracked and shared with its unique ID and web address. In addition, the web-based workflow also  
728 saves time for experts with automated model running, data assimilation, forecasting, structured  
729 result access and instantaneous graphic outputs, bringing the possibility for thorough exploration  
730 of more essence part of the system. The simplicity in use of EcoPAD (v1.0) at the same time  
731 may limit their access to the code and lowers the flexibility. Flexibility for users with higher  
732 demands, for example, those who wanted to test alternative data assimilation methods, use a  
733 different carbon cycle model, change the number of calibrated parameters, include the  
734 observations for other variables, is provided through the GitHub repository  
735 (<https://github.com/ou-ecolab> ). This GitHub repository contains code and instruction for  
736 installing, configuring and controlling the whole system, users can ~~easily~~ adapt the workflow to  
737 wrap their own model based on his or her needs. On one hand the web-based system with open  
738 source broadens the user community. On the other hand, it increases the risk of misuse and  
739 misinterpretation. We encourage users to be critical and consult system developers to avoid  
740 inappropriate application of the system.

#### 741 **4.2 Implications for better ecological forecasting**

742 Specifically to reliable forecasting of carbon dynamics, our initial exploration from  
743 EcoPAD-SPRUCE indicates that realistic model structure, correct parameterization and accurate  
744 external environmental conditions are essential. Model structure captures important ~~known~~  
745 mechanisms that regulate ecosystem carbon dynamics. Adjustment in model structure is critical  
746 in our improvement in methane forecasting. Model parameters may vary between observation  
747 sites, change with time or environmental conditions [Medlyn *et al.*, 1999; Luo *et al.*, 2001]. A

748 static or wrong parameterization misses important mechanisms (e.g., acclimation and adaptation)  
749 that regulate future carbon dynamics. Not well constrained parameters, for example, caused by  
750 lack of information from observational data, contribute to high forecasting uncertainty and low  
751 reliability of forecasting results. Correct parameterization is especially important for long-term  
752 carbon pool predictions as parameter uncertainty resulted in high forecasting uncertainty in our  
753 case study [*Jiang et al.*, 2018]. Parameter values derived under the ambient condition was not  
754 applicable to the warming treatment in our methane case due to acclimation. External  
755 environmental condition is another important factor in carbon predictions. External  
756 environmental condition includes both the external climatic forcing that is used to drive  
757 ecosystem models and also the environmental condition that is simulated by ecosystem models.  
758 As we showed that air warming may not proportionally transfer to soil warming, realistic soil  
759 environmental information needs to be appropriately represented to predict soil carbon dynamics  
760 [*Huang et al.*, 2017]. The impact of external forcing is especially obvious in short-term carbon  
761 flux predictions. Forcing uncertainty resulted in higher forecasting uncertainty in carbon flux  
762 compared to that from parameter uncertainty [*Jiang et al.*, 2018]. Mismatches in forecasted vs.  
763 realised forcing greatly increased simulated GPP and the discrepancy diminished in the long run.  
764 Reliable external environmental condition, to some extent, reduces the complexity in diagnosing  
765 modelled carbon dynamics.

766 Pool-based vs. flux-based predictions are regulated differently by external forcing and  
767 initial states, which indicates that differentiated efforts are required to improve short- vs. long-  
768 term predictions. External forcing, which has not been well emphasized in previous carbon  
769 studies, has strong impact on short-term forecasting. The large response of GPP to forecasted vs.  
770 realised forcing as well the stronger forcing-caused uncertainty in GPP predictions indicate

771 correct forcing information is a key step in short-term flux predictions. In this study, we  
772 stochastically generated the climate forcing based on local climatic conditions (1961-2014),  
773 which is not sufficient in capturing local short-term climate variability. As a result, updated GPP  
774 went outside our ensemble forecasting. On the other hand, parameters and historical information  
775 about pool status are more important in long-term pool predictions. Therefore, improvement in  
776 long-term pool size predictions cannot be reached by accurate climatic information alone.  
777 Instead, it requires accumulation in knowledge related to site history and processes that regulate  
778 pool dynamics.

779         Furthermore, reliable forecasting needs understanding of uncertainty sources in addition  
780 to the future mean states. Uncertainty and complexity are major reasons that lead to the belief in  
781 “computationally irreducible” and low intrinsic predictability of ecological systems [Coreau *et*  
782 *al.*, 2010; Beckage *et al.*, 2011; Schindler and Hilborn, 2015]. Recent advance in computational  
783 statistical methods offers a way to formally accounting for various uncertainty sources in  
784 ecology [Clark *et al.*, 2001; Cressie *et al.*, 2009]. And the Bayesian approach embedded in  
785 EcoPAD (v1.0) brings the opportunity to understand and communicate forecasting uncertainty.  
786 Our case study revealed that forcing uncertainty is more important in flux-based predictions  
787 while parameter uncertainty is more critical in pool-based predictions. Actually, how forecasting  
788 uncertainty changes with time, what are the dominate contributor of forecasting uncertainty (e.g.,  
789 parameter, initial condition, model structure, observation errors, forcing *etc.*), how uncertainty  
790 sources interact among different components, or to what extent unconstrained parameters affect  
791 forecasting uncertainty are all valuable questions that can be explored through EcoPAD (v1.0).

#### 792 **4.3 Applications of EcoPAD (v1.0) to manipulative experiments and observation sites**

793 Broadly speaking, data-model integration stands to increase the overall precision and  
794 accuracy of model-based experimentation [Luo *et al.*, 2011b; Niu *et al.*, 2014]. Systems for  
795 which data have been collected in the field and which are well represented by ecological models  
796 therefore have the capacity to receive the highest benefits from EcoPAD (v1.0) to improve  
797 forecasts. In a global change context, experimental manipulations including ecosystem responses  
798 to changes in precipitation regimes, carbon dioxide concentrations, temperatures, season lengths,  
799 and species compositional shifts can now be assimilated into ecosystem models [Xu *et al.*, 2006;  
800 Gao *et al.*, 2011; Lebauer *et al.*, 2013; Shi *et al.*, 2016]. Impacts of these global change factors  
801 on carbon cycling and ecosystem functioning can now be measured in a scientifically transparent  
802 and verifiable manner. This leads to ecosystem modelling of systems and processes that can  
803 obtain levels of confidence that lend credibility with the public to the science's forward progress  
804 toward forecasting and predicting [Clark *et al.*, 2001]. These are the strengths of a widely-  
805 available interface devoted to data-model integration towards better forecasting.

806 The data-model integration framework of EcoPAD (v1.0) creates a smart interactive  
807 model-experiment (ModEx) system. ModEx has the capacity to form a feedback loop in which  
808 field experiment guides modelling and modelling influences experimental focus [Luo *et al.*,  
809 2011a]. We demonstrated how EcoPAD (v1.0) works hand-in-hand between modellers and  
810 experimenters in the life-cycle of the SPRUCE project. ~~Field experiment from SPRUCE~~  
811 ~~community provides basic data to set up the ecosystem model and update model parameters~~  
812 ~~recursively, while the forecasting from ecosystem modelling informs experimenters the potential~~  
813 ~~key mechanisms that regulate ecosystem dynamics and help experimenters to question and~~  
814 ~~understand their measurements.~~ The EcoPAD-SPRUCE system operates while experimenters are  
815 making measurements or planning for future researches. Information is constantly fed back

816 between modellers and experimenters, and simultaneous efforts from both parties illustrate how  
817 communications between model and data advance and shape our understanding towards better  
818 forecasts during the lifecycle of a scientific project. ModEx can be extended to other  
819 experimental systems to: 1, predict what might be an ecosystem's response to treatments once  
820 experimenter selected a site and decided the experimental plan; 2, assimilate data experimenters  
821 are collecting along the experiment to constrain model predictions; 3, project what an  
822 ecosystem's responses may likely be in the rest of the experiment; 4, tell experimenters what are  
823 those important datasets experimenters may want to collect in order to understand the system; 5,  
824 periodically updates the projections; and 6, improve the models, the data assimilation system,  
825 and field experiments during the process.

826 In addition to the manipulative ~~experimental~~experiments, the data assimilation system of  
827 EcoPAD (v1.0)- can be used for automated model calibration for FLUXNET sites or other  
828 observation networks, such as the NEON and LTER [Johnson et al., 2010; Robertson et al.,  
829 2012]. The application of EcoPAD (v1.0) at FLUXNET, NEON or LTER sites includes three  
830 steps in general. First, build the climate forcing in the suitable formats of EcoPAD (v1.0) from  
831 the database of each site; Second, collect the prior information (include observations of state  
832 variables) in the data assimilation system from FLUXNET, NEON or LTER sites; Third,  
833 incorporate the forcing and prior information into EcoPAD (v1.0), and then run the EcoPAD  
834 (v1.0) with the dynamic data assimilation system. Furthermore, facing the proposed continental  
835 scale ecology study [Schimel, 2011], EcoPAD (v1.0) once properly applied could also help  
836 evaluate and optimize field deployment of environmental sensors and supporting  
837 cyberinfrastructure, that will be necessary for larger, more complex environmental observing  
838 systems being planned in the US and across different continents. ~~Altogether, with its milestone~~

Formatted: Font color: Custom Color(26,26,26)

~~concept, EcoPAD (v1.0) benefits from observation and modelling and at the same time advances both observation and modelling of ecological studies.~~

#### 4.4 Future developments

~~As we indicated,~~ EcoPAD (v1.0) will expand as time goes on. The system is designed to incorporate multiple process-based models, diverse data assimilation techniques and various ecological state variables for different ecosystems. Case studies presented in earlier sections are based primarily on one model. A multiple (or ensemble) model approach is helpful in tracking uncertainty sources from our process understanding. With rapid evolving ecological knowledge, emerging models with different hypotheses, such as the microbial-enzyme model [Wieder *et al.*, 2013], enhance our capacity in ecological prediction but can also benefit from rapid tests against data if incorporated into EcoPAD (v1.0). In addition to MCMC [Braswell *et al.*, 2005; Xu *et al.*, 2006], a variety of data assimilation techniques have been recently applied to improve models for ecological forecasting, such as the EnKF [Gao *et al.*, 2011], Genetic Algorithm [Zhou and Luo, 2008] and 4-d variational assimilation [Peylin *et al.*, 2016]. Future development will incorporate different optimization techniques to offer users the option to search for the best model parameters by selecting and comparing the possibly best method for their specific studies. We focus mostly on carbon related state variables in the SPRUCE example, and the data assimilation system in EcoPAD (v1.0) needs to include more observed variables for constraining model parameters. For example, the NEON sites not only provide measured ecosystem CO<sub>2</sub> fluxes and soil carbon stocks, but also resources (e.g., GPP/Transpiration for water and GPP/intercepted PAR for light) use efficiency [Johnson *et al.*, 2010].

~~With these improvements, one goal of EcoPAD (v1.0) is to enable the research community to understand and reduce forecasting uncertainties from different sources and~~

Formatted: Font color: Custom Color(26,26,26)

862 ~~forecast various aspects of future biogeochemical and ecological changes as data become~~  
863 ~~available. The example of Jiang et al. [2018] partitioned forecasting uncertainty from forcings~~  
864 ~~and parameters. An exhaustive understanding of forecasting uncertainty in ecology need to also~~  
865 ~~consider model structures, data assimilation schemes as well as different ecological state~~  
866 ~~variables.~~ Researchers interested in creating their own multiple model and/or multiple  
867 assimilation scheme version of EcoPAD (v1.0) can start from the GitHub repository  
868 (<https://github.com/ou-ecolab> ) where the source code of the EcoPAD (v1.0) workflow is  
869 archived. To add a new variable that is not forecasted in the EcoPAD-SPRUCE example, it  
870 requires modellers and experimenters to work together to understand their process-based model,  
871 ~~their~~ observations and how messaging works in the workflow of EcoPAD (v1.0) following the  
872 example of EcoPAD-SPRUCE. To add a new model or a new data assimilation scheme for  
873 variables that are forecasted in EcoPAD-SPRUCE, researchers need to create additional dockers  
874 and mount them to the existing workflow with the knowledge of how information are passed  
875 within the workflow: ~~—————~~ (see Appendix for detailed information).

876 With these improvements, one goal of the EcoPAD (v1.0) is to enable the research  
877 community to understand and reduce forecasting uncertainties from different sources and  
878 forecast various aspects of future biogeochemical and ecological changes as data becomes  
879 available. EcoPAD (v1.0) acts as a tool to link model and data, not as a substitution for neither  
880 model nor data. Ecological forecasting through EcoPAD (v1.0) relies strongly on theoretical  
881 (model) and empirical (data) ecological studies. Questions such as what are major factors  
882 regulating temporal variability of methane emissions cannot be directly answered by EcoPAD  
883 (v1.0). How to make use of EcoPAD (v1.0) to inspire breakthroughs in both theoretical and  
884 empirical ecological studies worth future exploration.

Formatted: Font color: Auto

Formatted: Font color: Custom Color(26,26,26)

885 The power of EcoPAD (v1.0) ~~not only~~ lies in ~~its scientific values, but also in the~~  
886 potential service it can bring to the society. Forecasting with carefully quantified uncertainty is  
887 helpful in providing support for natural resource manager and policy maker [Clark et al., 2001].  
888 It is always difficult to bring the complex mathematical ecosystem models to the general public,  
889 which creates a gap between current scientific advance and public awareness. The web-based  
890 interface from EcoPAD (v1.0) makes modelling as easy as possible without losing the  
891 connection to the mathematics behind the models. It will greatly transform environmental  
892 education and encourage citizen science [Miller-Rushing et al., 2012; Kobori et al., 2016] in  
893 ecology and climate change with future outreach activities to broadcast the EcoPAD (v1.0)  
894 platform.

Formatted: Font color: Black

## 895 **5 Conclusion**

896 The fully interactive web-based Ecological Platform for Assimilating Data (EcoPAD)  
897 (v1.0) into models aims to promote data-model integration towards predictive ecology through  
898 bringing the complex ecosystem model and data assimilation techniques accessible to different  
899 audience. It is supported by meta-databases of biogeochemical variables, libraries of modules of  
900 process models, toolbox of inversion techniques and the scalable scientific workflow. Through  
901 these components, it automates data management, model simulation, data assimilation,  
902 ecological forecasting, and result visualization, providing an open, convenient, transparent,  
903 flexible, scalable, traceable and readily portable platform to systematically conduct data-model  
904 integration towards better ecological forecasting.

Formatted: Font color: Black

905 We illustrated several of its functionalities through the Spruce and Peatland Responses  
906 Under Climatic and Environmental change (SPRUCE) experiment. The iterative forecasting  
907 approach from EcoPAD-SPRUCE through the prediction-question-discussion-adjustment-

908 prediction cycle and extensive communication between model and data creates a new paradigm  
909 to best inform forecasting. In addition to forecasting, EcoPAD [\(v1.0\)](#) enables interactive web-  
910 based approach to conduct model simulation, estimate model parameters or state variables,  
911 quantify uncertainty of estimated parameters and projected states of ecosystems, evaluate model  
912 structures, and assess sampling strategies. Altogether, EcoPAD-SPRUCE creates a smart  
913 interactive model-experiment (ModEx) system from which experimenters can know what an  
914 ecosystem's response might be at the beginning of their experiments, constrain models through  
915 collected measurements, predict ecosystem's response in the rest of the experiments, adjust  
916 measurements to better understand their system, periodically update projections and improve  
917 models, the data assimilation system, and field experiments.

918 Specifically to forecasting carbon dynamics, EcoPAD-SPRUCE revealed that better  
919 forecasting relies on improvements in model structure, parameterization and accurate external  
920 forcing. Accurate external forcing is critical for short-term flux-based carbon predictions while  
921 right process understanding, parameterization and historical information are essential for long-  
922 term pool-based predictions. In addition, EcoPAD [\(v1.0\)](#) provides an avenue to disentangle  
923 different sources of uncertainties in carbon cycling studies and to provide reliable forecasts with  
924 accountable uncertainties.

925

926 **Code availability:**

927 EcoPAD [\(v1.0\)](#) portal is available at [https://ecolab.nau.edu/ecopad\\_portal/](https://ecolab.nau.edu/ecopad_portal/) and code is provided  
928 at the GitHub repository (<https://github.com/ou-ecolab>).

929 **Data availability:**

930 Relevant data for this manuscript is available at the SPRUCE project webpage

931 (<https://mnspruce.ornl.gov/>) and the EcoPAD ([v1.0](#)) web portal

932 ([https://ecolab.nau.edu/ecopad\\_portal/](https://ecolab.nau.edu/ecopad_portal/)). Additional data can be requested from the

933 corresponding author.

934 **Competing interests:**

935 The authors declare that they have no conflict of interest.

936 **Acknowledgement:**

937 SPRUCE components of this work (PJH, DMR) are based upon work supported by the U.S.

938 Department of Energy, Office of Science, Office of Biological and Environmental Research. Oak

939 Ridge National Laboratory is managed by UT-Battelle, LLC, for the U.S. Department of Energy

940 under contract DE-AC05-00OR22725.

941

942 **Literature Cited**

943 Ahlstrom, A., G. Schurgers, A. Arneth, and B. Smith (2012), Robustness and uncertainty in  
944 terrestrial ecosystem carbon response to CMIP5 climate change projections,

945 *Environmental Research Letters*, 7(4), doi:10.1088/1748-9326/7/4/044008

946 Anderson, J., T. Hoar, K. Raeder, H. Liu, N. Collins, R. Torn, and A. Avellano (2009), The data  
947 assimilation research testbed A Community Facility, *Bulletin of the American*

948 *Meteorological Society*, 90(9), 1283-1296, doi:10.1175/2009bams2618.1

949 Baldocchi, D., E. Falge, L. H. Gu, R. Olson, D. Hollinger, S. Running, P. Anthoni, C. Bernhofer,

950 K. Davis, R. Evans, J. Fuentes, A. Goldstein, G. Katul, B. Law, X. H. Lee, Y. Malhi, T. Meyers, W.

951 Munger, W. Oechel, K. T. P. U, K. Pilegaard, H. P. Schmid, R. Valentini, S. Verma, T. Vesala, K.

952 Wilson, and S. Wofsy (2001), FLUXNET: A new tool to study the temporal and spatial

953 variability of ecosystem-scale carbon dioxide, water vapor, and energy flux densities,

954 *Bulletin of the American Meteorological Society*, 82(11), 2415-2434, doi:10.1175/1520-

955 0477(2001)082<2415:fantts>2.3.co;2

956 Ball, J. T., I. E. Woodrow, and J. A. Berry (1987), A model predicting stomatal conductance

957 and its contribution to the control of photosynthesis under different environmental

958 conditions, in *Progress in Photosynthesis Research*, edited by J. Biggens, pp. 221– 224,

959 Martinus Nijhoff, Zoetermeer, Netherlands.

960 Bastiaanssen, W. G. M., and S. Ali (2003), A new crop yield forecasting model based on

961 satellite measurements applied across the Indus Basin, Pakistan, *Agriculture Ecosystems &*

962 *Environment*, 94(3), 321-340, doi:10.1016/s0167-8809(02)00034-8

963 Beckage, B., L. J. Gross, and S. Kauffman (2011), The limits to prediction in ecological  
964 systems, *Ecosphere*, 2(11), doi:10.1890/es11-00211.1  
965 Bloom, A. A., J. F. Exbrayat, I. R. van der Velde, L. Feng, and M. Williams (2016), The decadal  
966 state of the terrestrial carbon cycle: Global retrievals of terrestrial carbon allocation, pools,  
967 and residence times, *Proceedings of the National Academy of Sciences of the United States*  
968 *of America*, 113(5), 1285-1290, doi:10.1073/pnas.1515160113  
969 Botkin, D. B., H. Saxe, M. B. Araujo, R. Betts, R. H. W. Bradshaw, T. Cedhagen, P. Chesson, T. P.  
970 Dawson, J. R. Etterson, D. P. Faith, S. Ferrier, A. Guisan, A. S. Hansen, D. W. Hilbert, C. Loehle,  
971 C. Margules, M. New, M. J. Sobel, and D. R. B. Stockwell (2007), Forecasting the effects of  
972 global warming on biodiversity, *Bioscience*, 57(3), 227-236, doi:10.1641/b570306  
973 Braswell, B. H., W. J. Sacks, E. Linder, and D. S. Schimel (2005), Estimating diurnal to annual  
974 ecosystem parameters by synthesis of a carbon flux model with eddy covariance net  
975 ecosystem exchange observations, *Global Change Biology*, 11(2), 335-355,  
976 doi:10.1111/j.1365-2486.2005.00897.x  
977 Clark, J. S., S. R. Carpenter, M. Barber, S. Collins, A. Dobson, J. A. Foley, D. M. Lodge, M.  
978 Pascual, R. Pielke, W. Pizer, C. Pringle, W. V. Reid, K. A. Rose, O. Sala, W. H. Schlesinger, D. H.  
979 Wall, and D. Wear (2001), Ecological forecasts: An emerging imperative, *Science*,  
980 293(5530), 657-660, doi:10.1126/science.293.5530.657  
981 Clark, J. S., M. Lewis, J. S. McLachlan, and J. HilleRisLambers (2003), Estimating population  
982 spread: What can we forecast and how well?, *Ecology*, 84(8), 1979-1988, doi:10.1890/01-  
983 0618  
984 Cook, B. I., T. M. Smith, and M. E. Mann (2005), The North Atlantic Oscillation and regional  
985 phenology prediction over Europe, *Global Change Biology*, 11(6), 919-926,  
986 doi:10.1111/j.1365-2486.2005.00960.x  
987 Corbet, S. A., N. M. Saville, M. Fussell, O. E. PrysJones, and D. M. Unwin (1995), The  
988 competition box: A graphical aid to forecasting pollinator performance, *Journal of Applied*  
989 *Ecology*, 32(4), 707-719, doi:10.2307/2404810  
990 Coreau, A., G. Pinay, J. D. Thompson, P. O. Cheptou, and L. Mermet (2009), The rise of  
991 research on futures in ecology: rebalancing scenarios and predictions, *Ecology Letters*,  
992 12(12), 1277-1286, doi:10.1111/j.1461-0248.2009.01392.x  
993 Coreau, A., S. Treyer, P. O. Cheptou, J. D. Thompson, and L. Mermet (2010), Exploring the  
994 difficulties of studying futures in ecology: what do ecological scientists think?, *Oikos*,  
995 119(8), 1364-1376, doi:10.1111/j.1600-0706.2010.18195.x  
996 Craft, C., J. Clough, J. Ehman, S. Joye, R. Park, S. Pennings, H. Y. Guo, and M. Machmuller  
997 (2009), Forecasting the effects of accelerated sea-level rise on tidal marsh ecosystem  
998 services, *Frontiers in Ecology and the Environment*, 7(2), 73-78, doi:10.1890/070219  
999 Cressie, N., C. A. Calder, J. S. Clark, J. M. V. Hoef, and C. K. Wikle (2009), Accounting for  
1000 uncertainty in ecological analysis: the strengths and limitations of hierarchical statistical  
1001 modeling, *Ecological Applications*, 19(3), 553-570, doi:10.1890/07-0744.1  
1002 Crozier, L. G., R. W. Zabel, and A. F. Hamlett (2008), Predicting differential effects of climate  
1003 change at the population level with life-cycle models of spring Chinook salmon, *Global*  
1004 *Change Biology*, 14(2), 236-249, doi:10.1111/j.1365-2486.2007.01497.x  
1005 Dietze, M. C., D. S. Lebauer, and R. Kooper (2013), On improving the communication  
1006 between models and data, *Plant Cell and Environment*, 36(9), 1575-1585,  
1007 doi:10.1111/pce.12043

1008 Diez, J. M., I. Ibanez, A. J. Miller-Rushing, S. J. Mazer, T. M. Crimmins, M. A. Crimmins, C. D.  
1009 Bertelsen, and D. W. Inouye (2012), Forecasting phenology: from species variability to  
1010 community patterns, *Ecology Letters*, 15(6), 545-553, doi:10.1111/j.1461-  
1011 0248.2012.01765.x  
1012 Ellison, A. M. (2004), Bayesian inference in ecology, *Ecology Letters*, 7(6), 509-520,  
1013 doi:10.1111/j.1461-0248.2004.00603.x  
1014 Ellison, A. M. (2010), Repeatability and transparency in ecological research, *Ecology*, 91(9),  
1015 2536-2539, doi:10.1890/09-0032.1  
1016 Farquhar, G. D., S. V. Caemmerer, and J. A. Berry (1980), A biochemical-model of  
1017 photosynthetic CO<sub>2</sub> assimilation in leaves of C<sub>3</sub> species, *Planta*, 149(1), 78-90,  
1018 doi:10.1007/bf00386231  
1019 Fordham, D. A., H. R. Akcakaya, M. B. Araujo, J. Elith, D. A. Keith, R. Pearson, T. D. Auld, C.  
1020 Mellin, J. W. Morgan, T. J. Regan, M. Tozer, M. J. Watts, M. White, B. A. Wintle, C. Yates, and B.  
1021 W. Brook (2012), Plant extinction risk under climate change: are forecast range shifts alone  
1022 a good indicator of species vulnerability to global warming?, *Global Change Biology*, 18(4),  
1023 1357-1371, doi:10.1111/j.1365-2486.2011.02614.x  
1024 Gao, C., H. Wang, E. S. Weng, S. Lakshmirarahan, Y. F. Zhang, and Y. Q. Luo (2011),  
1025 Assimilation of multiple data sets with the ensemble Kalman filter to improve forecasts of  
1026 forest carbon dynamics, *Ecological Applications*, 21(5), 1461-1473,  
1027 Hampton, S. E., C. A. Strasser, J. J. Tewksbury, W. K. Gram, A. E. Budden, A. L. Batcheller, C. S.  
1028 Duke, and J. H. Porter (2013), Big data and the future of ecology, *Frontiers in Ecology and*  
1029 *the Environment*, 11(3), 156-162, doi:10.1890/120103  
1030 Hanson, P. J., J. S. Riggs, W. R. Nettles, J. R. Phillips, M. B. Krassovski, L. A. Hook, L. Gu, A. D.  
1031 Richardson, D. M. Aubrecht, D. M. Ricciuto, J. M. Warren, and C. Barbier (2017), Attaining  
1032 whole-ecosystem warming using air and deep-soil heating methods with an elevated CO<sub>2</sub>  
1033 atmosphere, *Biogeosciences*, 14, 861-883, doi:10.5194/bg-14-861-2017  
1034 Hararuk, O., J. Y. Xia, and Y. Q. Luo (2014), Evaluation and improvement of a global land  
1035 model against soil carbon data using a Bayesian Markov chain Monte Carlo method, *Journal*  
1036 *of Geophysical Research-Biogeosciences*, 119(3), 403-417, doi:10.1002/2013jg002535  
1037 Hare, J. A., M. A. Alexander, M. J. Fogarty, E. H. Williams, and J. D. Scott (2010), Forecasting  
1038 the dynamics of a coastal fishery species using a coupled climate-population model,  
1039 *Ecological Applications*, 20(2), 452-464, doi:10.1890/08-1863.1  
1040 Huang, Y., J. Jiang, S. Ma, D. Ricciuto, P. J. Hanson, and Y. Luo (2017), Soil thermal dynamics,  
1041 snow cover and frozen depth under five temperature treatments in an ombrotrophic bog:  
1042 Constrained forecast with data assimilation, *Journal of Geophysical Research:*  
1043 *Biogeosciences*, doi:10.1002/2016JG003725  
1044 Jiang, J., Y. Huang, S. Ma, M. Stacy, Z. Shi, D. M. Ricciuto, P. J. Hanson, and Y. Luo (2018),  
1045 Forecasting responses of a northern peatland carbon cycle to elevated CO<sub>2</sub> and a gradient  
1046 of experimental warming, *Journal of Geophysical Research: Biogeosciences*,  
1047 doi:10.1002/2017jg004040  
1048 Johnson, B. R., T. U. Kampe, and M. Kuester (2010), Development of airborne remote  
1049 sensing instrumentations for NEON, paper presented at SPIE Optical Engineering+  
1050 Applications, International Society for Optics and Photonics.  
1051 Kearney, M. R., B. A. Wintle, and W. P. Porter (2010), Correlative and mechanistic models of  
1052 species distribution provide congruent forecasts under climate change, *Conservation*  
1053 *Letters*, 3(3), 203-213, doi:10.1111/j.1755-263X.2010.00097.x

1054 Keenan, T. F., E. Davidson, A. M. Moffat, W. Munger, and A. D. Richardson (2012), Using  
1055 model-data fusion to interpret past trends, and quantify uncertainties in future projections,  
1056 of terrestrial ecosystem carbon cycling, *Global Change Biology*, 18(8), 2555-2569,  
1057 doi:10.1111/j.1365-2486.2012.02684.x  
1058 Kobori, H., J. L. Dickinson, I. Washitani, R. Sakurai, T. Amano, N. Komatsu, W. Kitamura, S.  
1059 Takagawa, K. Koyama, T. Ogawara, and A. J. Miller-Rushing (2016), Citizen science: a new  
1060 approach to advance ecology, education, and conservation, *Ecological Research*, 31(1), 1-  
1061 19, doi:10.1007/s11284-015-1314-y  
1062 Kolka, R. K., S. D. Sebestyen, E. S. Verry, and K. N. Brooks (2011), *Peatland biogeochemistry*  
1063 *and watershed hydrology at the Marcell Experimental Forest*, 488 pp., CRC Press Boca Raton  
1064 Lebauer, D. S., D. Wang, K. T. Richter, C. C. Davidson, and M. C. Dietze (2013), Facilitating  
1065 feedbacks between field measurements and ecosystem models, *Ecological Monographs*,  
1066 83(2), 133-154, doi:10.1890/12-0137.1  
1067 Liang, J. Y., D. J. Li, Z. Shi, J. M. Tiedje, J. Z. Zhou, E. A. G. Schuur, K. T. Konstantinidis, and Y. Q.  
1068 Luo (2015), Methods for estimating temperature sensitivity of soil organic matter based on  
1069 incubation data: A comparative evaluation, *Soil Biology & Biochemistry*, 80, 127-135,  
1070 doi:10.1016/j.soilbio.2014.10.005  
1071 Ludascher, B., I. Altintas, C. Berkley, D. Higgins, E. Jaeger, M. Jones, E. A. Lee, J. Tao, and Y.  
1072 Zhao (2006), Scientific workflow management and the Kepler system, *Concurrency and*  
1073 *Computation-Practice & Experience*, 18(10), 1039-1065, doi:10.1002/cpe.994  
1074 Luo, Y. Q., and J. F. Reynolds (1999), Validity of extrapolating field CO<sub>2</sub> experiments to  
1075 predict carbon sequestration in natural ecosystems, *Ecology*, 80(5), 1568-1583,  
1076 doi:10.1890/0012-9658(1999)080[1568:voefce]2.0.co;2  
1077 Luo, Y. Q., S. Q. Wan, D. F. Hui, and L. L. Wallace (2001), Acclimatization of soil respiration to  
1078 warming in a tall grass prairie, *Nature*, 413(6856), 622-625, doi:10.1038/35098065  
1079 Luo, Y. Q., J. Melillo, S. L. Niu, C. Beier, J. S. Clark, A. T. Classen, E. Davidson, J. S. Dukes, R. D.  
1080 Evans, C. B. Field, C. I. Czimczik, M. Keller, B. A. Kimball, L. M. Kueppers, R. J. Norby, S. L.  
1081 Pelini, E. Pendall, E. Rastetter, J. Six, M. Smith, M. G. Tjoelker, and M. S. Torn (2011a),  
1082 Coordinated approaches to quantify long-term ecosystem dynamics in response to global  
1083 change, *Global Change Biology*, 17(2), 843-854, doi:10.1111/j.1365-2486.2010.02265.x  
1084 Luo, Y. Q., K. Ogle, C. Tucker, S. F. Fei, C. Gao, S. LaDeau, J. S. Clark, and D. S. Schimel (2011b),  
1085 Ecological forecasting and data assimilation in a data-rich era, *Ecological Applications*,  
1086 21(5), 1429-1442,  
1087 Luo, Y. Q., A. Ahlstrom, S. D. Allison, N. H. Batjes, V. Brovkin, N. Carvalhais, A. Chappell, P.  
1088 Ciais, E. A. Davidson, A. C. Finzi, K. Georgiou, B. Guenet, O. Hararuk, J. W. Harden, Y. J. He, F.  
1089 Hopkins, L. F. Jiang, C. Koven, R. B. Jackson, C. D. Jones, M. J. Lara, J. Y. Liang, A. D. McGuire,  
1090 W. Parton, C. H. Peng, J. T. Randerson, A. Salazar, C. A. Sierra, M. J. Smith, H. Q. Tian, K. E. O.  
1091 Todd-Brown, M. Torn, K. J. van Groenigen, Y. P. Wang, T. O. West, Y. X. Wei, W. R. Wieder, J.  
1092 Y. Xia, X. Xu, X. F. Xu, and T. Zhou (2016), Toward more realistic projections of soil carbon  
1093 dynamics by Earth system models, *Global Biogeochemical Cycles*, 30(1), 40-56,  
1094 doi:10.1002/2015gb005239  
1095 Luo, Y. Q. (2017), Transient dynamics of terrestrial carbon storage: mathematical  
1096 foundation and its applications,  
1097 Ma, S., J. Jiang, Y. Huang, D. Ricciuto, P. J. Hanson, and Y. Luo (2017), Data-constrained  
1098 projections of methane fluxes in a Northern Minnesota Peatland in response to elevated  
1099 CO<sub>2</sub> and warming (Accepted), *Journal of Geophysical Research: Biogeosciences*,

1100 Medlyn, B. E., F. W. Badeck, D. G. G. De Pury, C. V. M. Barton, M. Broadmeadow, R.  
 1101 Ceulemans, P. De Angelis, M. Forstreuter, M. E. Jach, S. Kellomaki, E. Laitat, M. Marek, S.  
 1102 Philippot, A. Rey, J. Strassmeyer, K. Laitinen, R. Liozon, B. Portier, P. Roberntz, K. Wang,  
 1103 and P. G. Jarvis (1999), Effects of elevated CO<sub>2</sub> on photosynthesis in European forest  
 1104 species: a meta-analysis of model parameters, *Plant Cell and Environment*, 22(12), 1475-  
 1105 1495, doi:10.1046/j.1365-3040.1999.00523.x  
 1106 Melillo, J. M., S. D. Frey, K. M. DeAngelis, W. J. Werner, M. J. Bernard, F. P. Bowles, G. Pold, M.  
 1107 A. Knorr, and A. S. Grandy (2017), Long-term pattern and magnitude of soil carbon  
 1108 feedback to the climate system in a warming world, *Science*, 358(6359), 101-105,  
 1109 doi:10.1126/science.aan2874  
 1110 Michener, W. K., and M. B. Jones (2012), Ecoinformatics: supporting ecology as a data-  
 1111 intensive science, *Trends in Ecology & Evolution*, 27(2), 85-93,  
 1112 doi:10.1016/j.tree.2011.11.016  
 1113 Miller-Rushing, A., R. Primack, and R. Bonney (2012), The history of public participation in  
 1114 ecological research, *Frontiers in Ecology and the Environment*, 10(6), 285-290,  
 1115 doi:10.1890/110278  
 1116 Moorcroft, P. R. (2006), How close are we to a predictive science of the biosphere?, *Trends*  
 1117 *in Ecology & Evolution*, 21(7), 400-407, doi:10.1016/j.tree.2006.04.009  
 1118 Mouquet, N., Y. Lagadeuc, V. Devictor, L. Doyen, A. Duputie, D. Eveillard, D. Faure, E. Garnier,  
 1119 O. Gimenez, P. Huneman, F. Jabot, P. Jarne, D. Joly, R. Julliard, S. Kefi, G. J. Kergoat, S. Lavorel,  
 1120 L. Le Gall, L. Meslin, S. Morand, X. Morin, H. Morlon, G. Pinay, R. Pradel, F. M. Schurr, W.  
 1121 Thuiller, and M. Loreau (2015), REVIEW: Predictive ecology in a changing world, *Journal of*  
 1122 *Applied Ecology*, 52(5), 1293-1310, doi:10.1111/1365-2664.12482  
 1123 Niu, S. L., Y. Q. Luo, M. C. Dietze, T. F. Keenan, Z. Shi, J. W. Li, and F. S. Chapin (2014), The  
 1124 role of data assimilation in predictive ecology, *Ecosphere*, 5(5), doi:10.1890/es13-00273.1  
 1125 Ong, J. B. S., M. I. C. Chen, A. R. Cook, H. C. Lee, V. J. Lee, R. T. P. Lin, P. A. Tambyah, and L. G.  
 1126 Goh (2010), Real-Time Epidemic Monitoring and Forecasting of H1N1-2009 Using  
 1127 Influenza-Like Illness from General Practice and Family Doctor Clinics in Singapore, *Plos*  
 1128 *One*, 5(4), doi:10.1371/journal.pone.0010036  
 1129 Osterweil, L. J., L. A. Clarke, A. M. Ellison, E. Boose, R. Podorozhny, and A. Wise (2010), Clear  
 1130 and Precise Specification of Ecological Data Management Processes and Dataset  
 1131 Provenance, *Ieee Transactions on Automation Science and Engineering*, 7(1), 189-195,  
 1132 doi:10.1109/tase.2009.2021774  
 1133 Parton, W. J., J. W. B. Stewart, and C. V. Cole (1988), Dynamics of c, n, p and s in grassland  
 1134 soils - a model, *Biogeochemistry*, 5(1), 109-131, doi:10.1007/bf02180320  
 1135 Parton, W. J., J. A. Morgan, G. M. Wang, and S. Del Grosso (2007), Projected ecosystem  
 1136 impact of the Prairie Heating and CO<sub>2</sub> Enrichment experiment, *New Phytologist*, 174(4),  
 1137 823-834, doi:10.1111/j.1469-8137.2007.02052.x  
 1138 Perretti, C. T., S. B. Munch, and G. Sugihara (2013), Model-free forecasting outperforms the  
 1139 correct mechanistic model for simulated and experimental data, *Proceedings of the*  
 1140 *National Academy of Sciences of the United States of America*, 110(13), 5253-5257,  
 1141 doi:10.1073/pnas.1216076110  
 1142 Petchey, O. L., M. Pontarp, T. M. Massie, S. Kefi, A. Ozgul, M. Weilenmann, G. M. Palamara, F.  
 1143 Altermatt, B. Matthews, J. M. Levine, D. Z. Childs, B. J. McGill, M. E. Schaepman, B. Schmid, P.  
 1144 Spaak, A. P. Beckerman, F. Pennekamp, and I. S. Pearse (2015), The ecological forecast

1145 horizon, and examples of its uses and determinants, *Ecology Letters*, 18(7), 597-611,  
1146 doi:10.1111/ele.12443

1147 Peylin, P., C. Bacour, N. MacBean, S. Leonard, P. Rayner, S. Kuppel, E. Koffi, A. Kane, F.  
1148 Maignan, F. Chevallier, P. Ciais, and P. Prunet (2016), A new stepwise carbon cycle data  
1149 assimilation system using multiple data streams to constrain the simulated land surface  
1150 carbon cycle, *Geoscientific Model Development*, 9(9), 3321-3346, doi:10.5194/gmd-9-  
1151 3321-2016

1152 Purves, D., J. Scharlemann, M. Harfoot, T. Newbold, D. P. Tittensor, J. Hutton, and S. Emmott  
1153 (2013), Time to model all life on Earth, *Nature*, 493(7432), 295-297,

1154 Robertson, G. P., S. L. Collins, D. R. Foster, N. Brokaw, H. W. Ducklow, T. L. Gragson, C. Gries,  
1155 S. K. Hamilton, A. D. McGuire, and J. C. Moore (2012), Long-term ecological research in a  
1156 human-dominated world, *BioScience*, 62(4), 342-353,

1157 Schaefer, K., C. R. Schwalm, C. Williams, M. A. Arain, A. Barr, J. M. Chen, K. J. Davis, D.  
1158 Dimitrov, T. W. Hilton, D. Y. Hollinger, E. Humphreys, B. Poulter, B. M. Raczka, A. D.  
1159 Richardson, A. Sahoo, P. Thornton, R. Vargas, H. Verbeeck, R. Anderson, I. Baker, T. A. Black,  
1160 P. Bolstad, J. Q. Chen, P. S. Curtis, A. R. Desai, M. Dietze, D. Dragoni, C. Gough, R. F. Grant, L. H.  
1161 Gu, A. Jain, C. Kucharik, B. Law, S. G. Liu, E. Lokipitiya, H. A. Margolis, R. Matamala, J. H.  
1162 McCaughey, R. Monson, J. W. Munger, W. Oechel, C. H. Peng, D. T. Price, D. Ricciuto, W. J.  
1163 Riley, N. Roulet, H. Q. Tian, C. Tonitto, M. Torn, E. S. Weng, and X. L. Zhou (2012), A model-  
1164 data comparison of gross primary productivity: Results from the North American Carbon  
1165 Program site synthesis, *Journal of Geophysical Research-Biogeosciences*, 117,  
1166 doi:10.1029/2012jg001960

1167 Schimel, D. (2011), The era of continental-scale ecology, *Frontiers in Ecology and the*  
1168 *Environment*, 9(6), 311-311,

1169 Schindler, D. E., and R. Hilborn (2015), Prediction, precaution, and policy under global  
1170 change, *Science*, 347(6225), 953-954, doi:10.1126/science.1261824

1171 Scholze, M., T. Kaminski, P. Rayner, W. Knorr, and R. Giering (2007), Propagating  
1172 uncertainty through prognostic carbon cycle data assimilation system simulations, *Journal*  
1173 *of Geophysical Research-Atmospheres*, 112(D17), doi:10.1029/2007jd008642

1174 Shi, Z., X. Xu, O. Hararuk, L. F. Jiang, J. Y. Xia, J. Y. Liang, D. J. Li, and Y. Q. Luo (2015),  
1175 Experimental warming altered rates of carbon processes, allocation, and carbon storage in  
1176 a tallgrass prairie, *Ecosphere*, 6(11), doi:10.1890/es14-00335.1

1177 Shi, Z., Y. H. Yang, X. H. Zhou, E. S. Weng, A. C. Finzi, and Y. Q. Luo (2016), Inverse analysis of  
1178 coupled carbon-nitrogen cycles against multiple datasets at ambient and elevated CO<sub>2</sub>,  
1179 *Journal of Plant Ecology*, 9(3), 285-295, doi:10.1093/jpe/rtv059

1180 Sitch, S., C. Huntingford, N. Gedney, P. E. Levy, M. Lomas, S. L. Piao, R. Betts, P. Ciais, P. Cox,  
1181 P. Friedlingstein, C. D. Jones, I. C. Prentice, and F. I. Woodward (2008), Evaluation of the  
1182 terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using  
1183 five Dynamic Global Vegetation Models (DGVMs), *Global Change Biology*, 14(9), 2015-  
1184 2039, doi:10.1111/j.1365-2486.2008.01626.x

1185 Steppe, K., J. S. von der Crone, and D. J. W. Pauw (2016), TreeWatch.net: A Water and  
1186 Carbon Monitoring and Modeling Network to Assess Instant Tree Hydraulics and Carbon  
1187 Status, *Frontiers in Plant Science*, 7, doi:10.3389/fpls.2016.00993

1188 Stumpf, R. P., M. C. Tomlinson, J. A. Calkins, B. Kirkpatrick, K. Fisher, K. Nierenberg, R.  
1189 Currier, and T. T. Wynne (2009), Skill assessment for an operational algal bloom forecast  
1190 system, *Journal of Marine Systems*, 76(1-2), 151-161, doi:10.1016/j.jmarsys.2008.05.016

1191 Sugihara, G., R. May, H. Ye, C. H. Hsieh, E. Deyle, M. Fogarty, and S. Munch (2012), Detecting  
1192 Causality in Complex Ecosystems, *Science*, 338(6106), 496-500,  
1193 doi:10.1126/science.1227079  
1194 Thomas, R. Q., E. B. Brooks, A. L. Jersild, E. Ward, R. H. Wynne, T. J. Albaugh, H. Dinon-  
1195 Aldridge, H. E. Burkhart, J. Domec, T. R. Fox, C. A. Gonzalez-Benecke, T. A. Martin, A.  
1196 Noormets, D. A. Sampson, and R. O. Teskey (2017), Leveraging 35 years of *Pinus taeda*  
1197 research in the southeastern US to constrain forest carbon cycle predictions: regional data  
1198 assimilation using ecosystem experiments, *Biogeosciences*, 14, 3525-3547,  
1199 Vitolo, C., Y. Elkhatib, D. Reusser, C. J. A. Macleod, and W. Buytaert (2015), Web technologies  
1200 for environmental Big Data, *Environmental Modelling & Software*, 63, 185-198,  
1201 doi:10.1016/j.envsoft.2014.10.007  
1202 Walker, A. P., P. J. Hanson, M. G. De Kauwe, B. E. Medlyn, S. Zaehle, S. Asao, M. Dietze, T.  
1203 Hickler, C. Huntingford, C. M. Iversen, A. Jain, M. Lomas, Y. Q. Luo, H. McCarthy, W. J. Parton,  
1204 I. C. Prentice, P. E. Thornton, S. S. Wang, Y. P. Wang, D. Warlind, E. S. Weng, J. M. Warren, F. I.  
1205 Woodward, R. Oren, and R. J. Norby (2014), Comprehensive ecosystem model-data  
1206 synthesis using multiple data sets at two temperate forest free-air CO<sub>2</sub> enrichment  
1207 experiments: Model performance at ambient CO<sub>2</sub> concentration, *Journal of Geophysical*  
1208 *Research-Biogeosciences*, 119(5), 937-964, doi:10.1002/2013jg002553  
1209 Wallenstein, M. D., and E. K. Hall (2012), A trait-based framework for predicting when and  
1210 where microbial adaptation to climate change will affect ecosystem functioning,  
1211 *Biogeochemistry*, 109(1-3), 35-47, doi:10.1007/s10533-011-9641-8  
1212 Wang, Y. P., and R. Leuning (1998), A two-leaf model for canopy conductance,  
1213 photosynthesis and partitioning of available energy I: Model description and comparison  
1214 with a multi-layered model, *Agricultural and Forest Meteorology*, 91(1-2), 89-111,  
1215 doi:10.1016/s0168-1923(98)00061-6  
1216 Wang, Y. P., C. M. Trudinger, and I. G. Enting (2009), A review of applications of model-data  
1217 fusion to studies of terrestrial carbon fluxes at different scales, *Agricultural and Forest*  
1218 *Meteorology*, 149(11), 1829-1842, doi:10.1016/j.agrformet.2009.07.009  
1219 Wang, Y. P., R. M. Law, and B. Pak (2010), A global model of carbon, nitrogen and  
1220 phosphorus cycles for the terrestrial biosphere, *Biogeosciences*, 7(7), 2261-2282,  
1221 doi:10.5194/bg-7-2261-2010  
1222 Ward, E. J., E. E. Holmes, J. T. Thorson, and B. Collen (2014), Complexity is costly: a meta-  
1223 analysis of parametric and non-parametric methods for short-term population forecasting,  
1224 *Oikos*, 123(6), 652-661, doi:10.1111/j.1600-0706.2014.00916.x  
1225 Weng, E. S., and Y. Q. Luo (2008), Soil hydrological properties regulate grassland ecosystem  
1226 responses to multifactor global change: A modeling analysis, *Journal of Geophysical*  
1227 *Research-Biogeosciences*, 113(G3), doi:10.1029/2007jg000539  
1228 Weng, E. S., and Y. Q. Luo (2011), Relative information contributions of model vs. data to  
1229 short- and long-term forecasts of forest carbon dynamics, *Ecological Applications*, 21(5),  
1230 1490-1505,  
1231 Weng, E. S., Y. Q. Luo, C. Gao, and R. Oren (2011), Uncertainty analysis of forest carbon sink  
1232 forecast with varying measurement errors: a data assimilation approach, *Journal of Plant*  
1233 *Ecology*, 4(3), 178-191, doi:10.1093/jpe/rtr018  
1234 Wieder, W. R., G. B. Bonan, and S. D. Allison (2013), Global soil carbon projections are  
1235 improved by modelling microbial processes, *Nature Climate Change*, 3(10), 909-912,  
1236 doi:10.1038/nclimate1951

1237 Xu, T., L. White, D. F. Hui, and Y. Q. Luo (2006), Probabilistic inversion of a terrestrial  
1238 ecosystem model: Analysis of uncertainty in parameter estimation and model prediction,  
1239 *Global Biogeochemical Cycles*, 20(2), doi:10.1029/2005gb002468  
1240 Zhou, T., and Y. Q. Luo (2008), Spatial patterns of ecosystem carbon residence time and  
1241 NPP-driven carbon uptake in the conterminous United States, *Global Biogeochemical*  
1242 *Cycles*, 22(3), doi:10.1029/2007gb002939  
1243 Zhou, X. H., T. Zhou, and Y. Q. Luo (2012), Uncertainties in carbon residence time and NPP-  
1244 driven carbon uptake in terrestrial ecosystems of the conterminous USA: a Bayesian  
1245 approach, *Tellus Series B-Chemical and Physical Meteorology*, 64,  
1246 doi:10.3402/tellusb.v64i0.17223

1247

1248 **Tables**

1249 Table 1. Discussion stimulated by EcoPAD-SPRUCE forecasting among modellers and  
 1250 experimenters on how to improve predictions of the relative contribution of different pathways  
 1251 of methane emissions

	Discussion
1	No strong bubbles are noted at field and a non-observation constrained modelling study at a similar site from another project concluded minor ebullition contribution, which are at odds with TECO result.
2	CH <sub>4</sub> :CO <sub>2</sub> ratio might explain the discrepancy. The other modelling study assumed that decomposed C is mainly turned into CO <sub>2</sub> and a smaller fraction is turned into CH <sub>4</sub> . The large CH <sub>4</sub> :CO <sub>2</sub> ratio at this site may result in higher CH <sub>4</sub> flux. It seems that the most “flexible” term is ebullition because any “excess” (above saturation) CH <sub>4</sub> is immediately released to ebullition, while the plant transport term is constrained by vegetation data.
3	Experimental researches on the relative contribution to methane emission from three different pathways are rare.
4	Current available observations include net surface flux of methane from the large collars, incubation data that should represent methane sources within the profile, and gas/DOC profile data that can indicate active zones within the peat profile. What are additional data needed to constrain relative contribution of different pathways?
5	I had always thought that peatlands don’t bubble much, but the super-sensitive GPS measurements found movements of the surface of the GLAP peatlands consistent with degassing events, and subsurface radar images did show layers that were interpreted as bubble-layers.
6	Pore water gas data, perhaps N <sub>2</sub> or Ar may shed some light on the relative importance of ebullition.
7	It is really hard to accurately distinguish the three pathways. It has to rely on multiple approaches. Particularly for the SPRUCE site, the vegetation cover varies, vegetation species varies. How many channels each species has affect the transport? Meanwhile, the presence of plant (even not vascular plant) will lead to more gas transport, but as bubbles, rather than plant-mediated transport.
8	It depends on model structure and algorithm to simulate diffusion, vascular, and ebullition. Most models assume a threshold to allow ebullition. Diffusion is treated in similar ways as ebullition in some models (most one layer or two layers models). For the multiple layers models, the diffusion occurs from bottom to top mm by mm, layer by layer, therefore, the gas diffusion from top layer to atmosphere is considered the diffusion flux. If that is the case, the time step and wind speed and pressure matter (most models do not consider wind and pressure impacts). Plant transport is really dependent on the parameter for plant species, aerenchyma, etc. The gas transportability of plant is associated with biomass, NPP, or root biomass, seasonality of plant growth, etc. in models. All these differences might cause biases in the final flux.
9	With only the CH <sub>4</sub> emission data cannot constrain the relative contribution of three pathways. Concentration data in different soil layers may help constrain.
10	Diffusion coefficient calculation in TECO adopts the “three-porosity-model” which is ideal for mineral soil, but may not fit the organic soil. “Millington-Quirk model” for should be a better choice for peat soil.
11	The boundary condition should be taken care of, but it brings in more uncertainties including the wind speed and piston velocity, etc.,
12	CH <sub>4</sub> emissions captured in static chambers does not include the episodic ebullition events. So (1) the static chambers underestimate the total methane emission and (2) might need to exclude the ebullition pathway when using the observation data to constrain the CH <sub>4</sub> emission. But this point seems haven't been paid attention to in other models.

1252

1253 **Figure Legends**

1254 **Figure 1** Schema of approaches to forecast future ecological responses from common practice  
1255 (the upper panel) and the Ecological Platform for Assimilation of Data (EcoPAD (v1.0))  
1256 (bottom panel). The common practice makes use of observations to develop or calibrate models  
1257 to make predictions while the EcoPAD (v1.0) approach advances the common practice through  
1258 its fully interactive platform. EcoPAD (v1.0) consists of four major components:  
1259 experiment/data, model, data assimilation and the scientific workflow (green arrows or lines).  
1260 Data and model are iteratively integrated through its data assimilation systems to improve  
1261 forecasting. And its near-real time forecasting results are shared among research groups through  
1262 its web interface to guide new data collections. The scientific workflow enables web-based data  
1263 transfer from sensors, model simulation, data assimilation, forecasting, result analysis,  
1264 visualization and reporting, encouraging broad user-model interactions especially for the  
1265 experimenters and the general public with limited background in modelling. Images from the  
1266 SPRUCE field experiments (<https://mnspruce.ornl.gov/>) are used to represent data collection and  
1267 the flowchart of TECO model is used to delegate ecological models.

1268 **Figure 2** The data assimilation system inside the Ecological Platform for Assimilation of Data  
1269 (EcoPAD (v1.0)) towards better forecasting of terrestrial carbon dynamics

1270 **Figure 3** The scientific workflow of EcoPAD (v1.0). The workflow wraps ecological models  
1271 and data assimilation algorithms with the docker containerization platform. Users trigger  
1272 different tasks through the Representational State Transfer (i.e., RESTful) application-  
1273 programming interface (API). Tasks are managed through the asynchronous task queue, Celery.  
1274 Tasks can be executed concurrently on a single or more worker servers across different scalable

1275 IT infrastructures. MongoDB is a database software that takes charge of data management in  
1276 EcoPAD [\(v1.0\)](#) and RabbitMQ is a message broker.  
1277

1278 **Figure 4.** Schema of interactive communication between modellers and experimenters through  
1279 the prediction-question-discussion-adjustment-prediction cycle to improve ecological  
1280 forecasting. The schema is inspired by an episode of experimenter-modeller communication  
1281 stimulated by the EcoPAD-SPRUCE platform. The initial methane model constrained by static  
1282 chamber methane measurements was used to predict relative contributions of three methane  
1283 emission pathways (i.e., ebullition, plant mediated transportation (PMT) and diffusion) to the  
1284 overall methane fluxes under different warming treatments (+ 0 °C, +2.25 °C, +4.5 °C, +6.75 °C  
1285 and +9 °C). The initial results indicated a dominant contribution from ebullition especially under  
1286 +9 °C which was doubted by experimenters. The discrepancy stimulated communications  
1287 between modellers and experimenters with detailed information listed in Table 1. After extensive  
1288 discussion, the model structure was adjusted and field observations were re-evaluated. And a  
1289 second round of forecasting yielded more reliable predictions.

1290 **Figure 5.** Posterior distribution of the ratio of CH<sub>4</sub>:CO<sub>2</sub> (panel a) and the temperature sensitivity  
1291 of methane production (Q<sub>10\_CH4</sub>, panel b) under 5 warming treatments.

1292 **Figure 6.** Updated vs. un-updated forecasting of gross primary production (GPP, panels a,c) and  
1293 soil organic C content (SoilC, panels b,d). The upper panels show 3 series of forecasting with  
1294 updated vs. stochastically generated weather forcing. [“updated” means the real meteorology](#)  
1295 [forcing monitored from field weather station](#). Cyan indicates forecasting with 100 stochastically  
1296 generated weather forcing from January 2015 to December 2024 (S1); red corresponds to  
1297 updated forecasting with two stages, that is, updating with measured weather forcing from

1298 January 2015 to July 2016 followed by forecasting with 100 stochastically generated weather  
1299 forcing from August 2016 to December 2024 (S2); and blue shows updated forecasting with  
1300 measured weather forcing from January 2015 to December 2016 followed by forecasting with  
1301 100 stochastically generated weather forcing from January 2017 to December 2024 (S3). The  
1302 bottom panels display mismatches between updated forecasting (S2,3) and the original un-  
1303 updated forecasting (S1). Red displays the difference between S2 and S1 (S2-S1) and blue shows  
1304 discrepancy between S3 and S1 (S3-S1). Dashed green lines indicate the start of forecasting with  
1305 stochastically generated weather forcing. Note that the left 2 panels are plotted on yearly time-  
1306 scale and the right 2 panels show results on monthly time-scale.


1307

1308

1309

1310 **Figure 1**

1311




1312

1313

1314 **Figure 2**

1315



1316

1317


1318

1319


1320

1321 **Figure 3**

1322



1323




1324

1325

1326

1327 **Figure 4**

1328




1329

1330

1331 **Figure 5**

1332



1333


1334

1335

1336

1337

1338 **Figure 6**



1339

1340